

CS101 (Fall 2012): Problem Set #3 Bayesian Networks

Chuan

December 2, 2012

1 Problem 1

- a. (c)
- b. (a)
- c. (a)
- d. The CPT for the G_{child} :

G_{mother}	G_{father}	$P(G_{child} = l)$	$P(G_{child} = r)$
l	l	$1 - m$	m
l	r	0.5	0.5
r	l	0.5	0.5
r	r	m	$1 - m$

e.

$$\begin{aligned}
 P(G_{child} = l) &= \sum_{G_{mother}, G_{father}} P(G_{child} = l, G_{father}, G_{mother}) \\
 &= P(G_{child} = l, G_{father} = l, G_{mother} = l) + P(G_{child} = l, G_{father} = l, G_{mother} = r) + \\
 &\quad P(G_{child} = l, G_{father} = r, G_{mother} = l) + P(G_{child} = l, G_{father} = r, G_{mother} = r) \\
 &= P(G_{child} = l \mid G_{father} = l, G_{mother} = l)P(G_{father} = l, G_{mother} = l) + \\
 &\quad P(G_{child} = l \mid G_{father} = l, G_{mother} = r)P(G_{father} = l, G_{mother} = r) + \\
 &\quad P(G_{child} = l \mid G_{father} = r, G_{mother} = l)P(G_{father} = r, G_{mother} = l) + \\
 &\quad P(G_{child} = l \mid G_{father} = r, G_{mother} = r)P(G_{father} = r, G_{mother} = r) \\
 &= (1 - m)q^2 + 0.5q(1 - q) + 0.5(1 - q)q + m(1 - q)^2 \\
 &= (1 - m)q^2 + q(1 - q) + m(1 - q)^2
 \end{aligned}$$

f. According to e., we get:

$$\begin{aligned}P(G_{child} = l) &= P(G_{father}) = P(G_{mother}) = q \\(1 - m)q^2 + q(1 - q) + m(1 - q)^2 &= q \\q &= 0.5\end{aligned}$$

As we know about the handedness in humans, a variety of studies suggest that 70-90% of the world population is right-handed(1). But according to the hypothesis, the distribution of handedness is same as the gene, which means 50% of the world population is right-handed. This is not true. So the hypothesis is wrong.

References

- [1] *Handedness. From Wikipedia, the free encyclopedia.*