


Assessment	Test	Domain	Skill	Difficulty
SAT	Reading and Writing	Information and Ideas	Command of Evidence	Hard

Question



The population of the coral *Lophelia pertusa* declined significantly around 9,000 years ago in the Alboran Sea and around 11,000 years ago near the Mauritanian coast. Using the ratio of manganese to calcium, which inversely correlates with ocean oxygenation levels, marine scientist Rodrigo da Costa Portilho-Ramos and colleagues evaluated whether oxygenation played a role in the declines of *L. pertusa*. The researchers concluded that oxygenation may have been important in the Alboran Sea but not near the Mauritanian coast, since _____

Which choice most effectively uses data from the graph to complete the statement?

- Answer**
- A. a substantial increase in oxygenation in the Alboran Sea corresponded with the local decline in *L. pertusa*, but the opposite relationship between oxygenation and *L. pertusa* was found near the Mauritanian coast.
 - B. *L. pertusa* declined in the Alboran Sea during a period of substantial local decline in oxygenation, but *L. pertusa* declined near the Mauritanian coast during a period of little local change in oxygenation.
 - C. oxygenation in the Alboran Sea was higher before the decline in *L. pertusa* than after the decline, whereas oxygenation near the Mauritanian coast was relatively low both before and after the decline in *L. pertusa*.
 - D. oxygenation in the Alboran Sea tended to be substantially higher than oxygenation near the Mauritanian coast during the period studied.

Rationale

Choice B is the best answer because it effectively uses data from the graph to complete the statement about Rodrigo da Costa Portilho-Ramos and colleagues' conclusion. The graph shows the ratio of manganese to calcium in *L. pertusa* coral samples from the Alboran Sea and the Mauritanian coast. The graph reflects time in approximate years before present: in other words, the greater the number in years noted on the graph's horizontal axis, the farther that moment is in the past. The text indicates that the researchers tested the samples to determine whether oxygenation played a role in the decline of *L. pertusa*. The text goes on to note that a change in the ratio of manganese to calcium would signal an inverse, or opposite, change in oxygenation. According to the graph, the ratio of manganese to calcium in samples from the Alboran Sea increased from about 30 micromoles per mole 10,000 years ago to about 80 micromoles per mole 8,000 years ago, which means that oxygenation decreased between 10,000 and 8,000 years ago. Meanwhile, there was almost no discernible change in the ratio of manganese to calcium in samples from the Mauritanian coast between 12,000 and 10,000 years ago. According to the text, the population of *L. pertusa* declined significantly around 9,000 years ago in the Alboran Sea and around 11,000 years ago near the Mauritanian coast. Thus, the increase in the ratio of manganese to calcium around 9,000 years ago in the Alboran Sea coincides with the decline in the *L. pertusa* population, suggesting an association between the decrease in oxygenation and the decline in population of the coral. No such relationship is suggested around 11,000 years ago near the Mauritanian coast. So, oxygenation likely played a role in the *L. pertusa* decline in the Alboran Sea but not in the coral's decline near the Mauritanian coast.

Choice A is incorrect because it asserts the opposite of what the graph indicates regarding oxygenation in the Alboran Sea, and it misrepresents what the graph indicates about oxygenation near the Mauritanian coast. The graph indicates that at the time of the decline in *L. pertusa* (approximately 9,000 years ago), the samples from the Alboran Sea contained a ratio of manganese to calcium that was increasing. According to the text, this ratio inversely correlates with ocean oxygenation levels, so if the ratio was increasing, oxygenation was decreasing, not substantially increasing. Furthermore, the graph shows that the ratio of manganese to calcium remained relatively stable in coral samples from the Mauritanian coast during the period studied, which suggests that there was no discernible relationship between oxygenation and the coral's population decline in that location, not that there was a substantial decrease in oxygenation corresponding to the coral's decline. Choice C is incorrect. Although the graph suggests that the level of oxygenation in the Alboran Sea was higher before the decline in *L. pertusa* than after—because the ratio of manganese to calcium inversely correlates with ocean oxygenation levels and this ratio was lower before the decline than after—the graph doesn't support the claim that oxygenation near the Mauritanian coast was consistently low before and after the coral's decline there. Rather, the graph indicates that relative to coral samples from the Alboran Sea, the ratio of manganese to calcium in samples from near the Mauritanian coast was consistently low, which suggests that oxygenation levels were relatively high both before and after the decline of *L. pertusa*. Choice D is incorrect because it states the opposite of what the graph indicates: the graph shows that throughout the period studied, the ratio of manganese to calcium was higher in coral samples from the Alboran Sea than it was in samples from near the Mauritanian coast. Since the text indicates that the ratio of manganese to calcium inversely correlates with ocean oxygenation levels, oxygenation in the Alboran Sea was therefore lower than, not higher than, oxygenation near the Mauritanian coast during the period studied. Moreover, even if choice D did accurately represent the graph, it wouldn't effectively complete the statement since a comparison of the ocean oxygenation levels at the two locations is not relevant to the claim that a decline in oxygenation levels was associated with the decline of *L. pertusa* in the Alboran Sea but not near the Mauritanian coast.