

Question

Delta 15-N Values in Seagrass Samples from Four Sites on the Yucatan, 2016–2017

Site	February 2016	October 2016	February 2017	October 2017
Akumal Bay	no data available	3.3	2.0	6.3
Mahahual	0.7	no data available	2.5	3.4
Tulum	6.1	5.9	2.3	5.5
Xahuayxol	0.9	0.3	–0.9	1.4

Because water from natural, uncontaminated sources is less enriched with the stable nitrogen isotope ^{15}N than wastewater from human activities is, the presence of such wastewater in nature can be detected by examining delta 15-N values (a measure of the ratio of ^{15}N to ^{14}N) in plants. Karla A. Camacho-Cruz and colleagues assessed delta 15-N values in the seagrass *Thalassia testudinum* from sites on Mexico’s Yucatan peninsula with intermediate tourism development, including Akumal Bay and Tulum, and low tourism development, including Mahahual and Xahuayxol, throughout 2016 and 2017. The data suggest that the intermediate-tourism sites experienced influxes of human wastewater. However, the researchers concluded that this happened intermittently.

Which choice best describes data from the table that support the underlined conclusion?

Answer

- A. Although delta 15-N values were generally higher in Akumal Bay and Tulum than in Mahahual and Xahuayxol, the values were lower in Akumal Bay than in Mahahual and Xahuayxol in February 2017.
- B. Delta 15-N values reached their lowest level in February 2017 in both Akumal Bay and Tulum, but no data were available for Akumal Bay in February 2016, when the values reached their highest level in Tulum.
- C. Although all sites showed considerable variation in delta 15-N values, the values remained relatively constant in Akumal Bay from October 2016 to February 2017 and in Tulum from February 2016 to October 2016.
- D. In Akumal Bay and Tulum, delta 15-N values fluctuated considerably across the three measurements made from October 2016 to October 2017.