6 Probability 3

6.1 Expected value (Mean)

The **mean** of a discrete random variable is:

$$E(Y) = \sum_{i=1}^{n} y_i P(Y = y_i)$$

where y_i's are the values that the variable takes on and the sum is taken over all n possible values that Y can take.

Rules: If X and Y are two random variables and Z=X+Y // linear operation //

$$E(Z) = E(X+Y) = E(X) + E(Y)$$

$$E(aX+b) = aE(X) + b$$

6.2 Variance

The **variance** of a discrete random variable is

$$Var(Y) = E((Y - E(Y)^2) = \sum_{i=1}^{n} (y_i - E(Y))^2 P(Y = y_i)$$

In addition

$$Var(Y) = E(Y^2) - E(Y)^2$$

and the standard deviation is

$$SD_Y = \sqrt{Var(Y)}$$

The **covariance** between two random variables is / a measure of association

$$Cov(X,Y) = E(XY) - E(X)E(Y) = E[(X - E(X))(Y - E(y))]$$

and the coefficient of correlation is

$$\rho_{X,Y} = \frac{Cov(X,Y)}{SD_X SD_Y}$$

In many contexts-such as molecular biology-we count events (e.g., codons, DNA reads, CG diagrams), producing discrete variables, unlike continuous measures such as mass or intensity. Some variables are binary (e.g., mutation: yes/no), while others have multiple levels (e.g., genotypes AA, Aa, aa) or even many

$$\mathrm{var}(x) = \mathrm{cov}(x, x)$$
$$= E[x^2] - E[x]^2$$

$$\mathrm{cov}(x, y) = E[xy] - E[x]E[y] \leftarrow$$

$$\mathrm{var}(ax+b) = \mathrm{cov}(ax+b, ax+b)$$

$$= \mathrm{cov}(ax, ax) + \cancel{\mathrm{cov}(ax, b)}^{0} +$$
$$\cancel{\mathrm{cov}(b, ax)}^{0} + \cancel{\mathrm{cov}(b, b)}^{0}$$

Note: $\mathrm{cov}(b, b) = 0$

$$\mathrm{cov}(b, ax) = E[abx] - E[b]E[ax]$$
$$= ab\,E[x] - ba\,E[x]$$
$$= 0$$

$$\mathrm{cov}(ax, ax) = E[a^2x^2] - E[ax]E[ax]$$
$$= a^2E[x^2] - a^2E[x]^2$$
$$= a^2\,\mathrm{var}(x)$$

$$\mathrm{var}(ax) = a^2\,\mathrm{var}(x)$$

10^2

categories (e.g., bacterial types or 64 codons).

To summarize categorical data, we tally the frequency of each level. In R, these are stored as factors, which make tabulation straightforward. For example, here we capture the different blood genotypes for 19 subjects in a vector which we tabulate.

```
genotype = c("AA","AO","BB","AO","OO","AO","AA","BO","BO",
             "AO","BB","AO","BO","AB","OO","AB","BB","AO","AO")
table(genotype)
```

```
genotype
AA AB AO BB BO OO
 2  2  7  3  3  2
```

If the order in which the data are observed does not matter, we call the random variable **exchangeable**. In this case, the vector of frequencies is sufficient to capture all the relevant information in the data, providing an efficient way to summarize it.

6.2.1 Bernoulli trials

The **Bernoulli distribution** is the simplest probability model, based on a so-called Bernoulli random variable, which can take only two outcomes: failure or success. The key idea is that you have a parameter p, which represents the probability of success, and the complementary event is called probability of failure $1 - p$:

Probability distribution:

$\boxed{X \sim Be(p)}$

$$P(X = 1) = p$$ ← one parameter p
$$P(X = 0) = 1 - p$$

For a fair coin, p = 0.5. For a biased coin that lands heads 70% of the time, p = 0.7.

> **Note**
>
> **Example:** Coin tosses
> Suppose we want to simulate a sequence of 15 fair tosses using R. To get the outcome of 15 tosses of a fair coin (p = 0.50), we write:

samples

```
rbinom(15, prob = 0.5, size = 1)
```

```
[1] 0 0 1 0 0 0 0 0 0 0 1 1 0 0 0
```

Now observe the outcome when using a biased coin that lands heads 70% of the time:

set.seed(12) ← makes sure you use the same "random" generator

```
rbinom(15, prob = 0.7, size = 1)
```

```
[1] 1 0 1 1 1 0 0 1 1 1 0 1 1 1 1
```

$X \sim Be(p) \qquad x = \begin{cases} 0 & w.p\ 1-p \\ 1 & w.p\ p \end{cases}$

$E[X] = (0)(1-p) + (1)p$

$= p$

$V[X] = E[X^2] - E[X]^2$

$= p - p^2 = p(1-p)$

$P(x=0) = 1-p$

$P(x=1) = p$

$x \sim Bin(n,p)$

$E[X] = \sum_{i=0}^{n} i\, P(x=i)$

$P(x=0) = (1-p)^n$

$\rightarrow P(x=1) = \binom{n}{1} p(1-p)^{n-1} = \frac{n!}{(n-1)!} p(1-p)^{n-1} = np(1-p)^{n-1}$

$P(x=k) = \binom{n}{k} p^k (1-p)^{n-k}$

$P(x=n) = p^n$

$\underbrace{0101}_{n}$

0's or 1's and i 1's.

$\binom{n}{i} = nC_i = \frac{n!}{i!(n-i)!}$

$n! = n(n-1)(n-2)\ldots(1)$

$$\binom{5}{2}$$

$$= \frac{5!}{2!\,3!}$$

$$= \frac{5 \times \cancel{4}^{2} \times \cancel{3} \times \cancel{2} \times 1}{(\cancel{2} \times 1)(\cancel{3} \times \cancel{2} \times 1)}$$

$$= 10$$

11000
10100
10010
10001
01100
01010
01001
00110
⋮
00011

$\binom{n}{i}$ = # sets of n elements with i of one type and $n-i$ of the other type

Question. Note that when you call the function above several times, you get different outcomes. Why is that?

Properties

The **mean**, or **expected value**, of a Bernoulli distribution is given by p, and the **variance** is defined as $p(1-p)$, so the variance is maximized when p = 0.5, and it goes to 0 as p approaches 0 or 1. The Bernoulli distribution is very important because it represents the building block of many other distributions.

6.2.2 Binomial success counts = sum of bernoulli vars.

In the previous example, if we are only interested in the number of heads ($X = 1$), we just take the sum of the cells in the output vector, then reporting a single number instead of a binary vector. In R, we can do this as follows:

```
rbinom(1, prob = 0.7, size = 15)
```

$X \sim Bin(n, p)$

trials / experiments.

success probability.

```
[1] 10
```

The sum of multiple independent (or exchangeable) Bernoulli trials is called the **Binomial distribution** (in the example above, how many heads in 15 flips). That is, the number of successes in 15 Bernoulli trials with a probability of success of 0.7 is called a **binomial** random variable, or a random variable that follows the B(15, 0.7) distribution. We can generate binomial samples in R as follows:

```
set.seed(384569724)
rbinom(1, prob = 0.7, size = 15)
```

```
[1] 11
```

Question. Repeat this function call 10 times. What seems to be the most common outcome?

Solution. The most frequent value is 10.5≈ 11. In fact, the theoretical proportion of times that we expect 10.5 to appear is the value of the probability that $X = 4$ if X follows B(15,0.7).

The complete **probability mass distribution** is available by typing:

```
probabilities = dbinom(0:15, prob = 0.7, size = 15)
round(probabilities, 2)
```

```
 [1] 0.00 0.00 0.00 0.00 0.00 0.00 0.01 0.03 0.08 0.15 0.21
0.22 0.17 0.09 0.03
[16] 0.00
```

A barplot for this distribution in R, is as follows:

```
barplot(probabilities, names.arg = 0:15, col = "black", main = '
```



Mathematically, the theoretical distribution of the binomial distribution is

Probability distribution:

$$P(X = k) = \binom{n}{k} p^k (1-p)^{n-k}, \quad k = 0, 1, \ldots, n,$$

where n is the number of trials, k is the number of successes, p is the probability of success, and $\binom{n}{k} = \frac{n!}{k!(n-k)!}$ is the binomial coefficient.

Properties

The **mean** of a Binomial distribution is given by $n \cdot p$ (in the figure, the highest bar around $x \approx 11$, or $n(=15) \cdot p(=0.7)$, and the **variance** is defined as $np(1-p) = (15)(0.7)(0.3) = 3.15$.

6.3 Poisson distributions

When the probability of success p is small and the number of trials n is large, the Binomial distribution can be approximated by a **Poisson distribution** with parameter $\lambda = np$, which provides a simple model for describing the data.

Probability distribution:

$$P(X = k) = \frac{e^{-\lambda}\lambda^k}{k!}, \quad k = 0, 1, 2, \ldots,$$

where k is the number of events, λ is the average number of events per interval, and $e \approx 2.718$ is the Euler's number.

Properties