



# A Chiral Symmetric Dirac Equation Watson & Musielak (Int. J. Mod. Phys. A, 2020)

Julian L. Avila-Martinez   Laura Y. Herrera-Martinez   Sebastian Rodriguez-Garcia

Física, Universidad Distrital Francisco José de Caldas

## What is Chirality?

### Chiral Symmetric Dirac Equation

$$\left(i\gamma^\mu\partial_\mu - me^{-i2\alpha\gamma^5}\right)\psi = 0 \quad (1)$$

## Chiral Angle and Mass

### What is the Chiral Angle?

The DECS can be re-written (eq. (1)):

$$\left(i\gamma^\mu\partial_\mu - M - \tilde{M}\gamma^5\right)\psi = 0 \quad (2)$$

Where:

- $M = m \cos 2\alpha$  is the standard **scalar mass**.
- $\tilde{M} = -im \sin 2\alpha$  is a **pseudoscalar mass**. The chiral angle  $\alpha$  mixes the scalar and pseudoscalar mass terms.

### Generating Mass: A Two-Field Higgs Model

These  $M$  and  $\tilde{M}$  terms can arise from Yukawa couplings to two Higgs-like fields: a scalar  $\phi_1$  and a pseudoscalar  $\phi_2$ .

$$\mathcal{L}_Y \approx -\frac{\lambda_1 v_1}{\sqrt{2}}\bar{\psi}\psi - \frac{\lambda_2 v_2}{\sqrt{2}}\bar{\psi}\gamma^5\psi \quad (3)$$

This *fixes* the parameters  $m$  and  $\alpha$ :

$$m = \sqrt{\frac{\lambda_1^2 v_1^2 - \lambda_2^2 v_2^2}{2}} \quad (4)$$

$$\alpha = \frac{i}{4} \ln \left( \frac{\lambda_1 v_1 + \lambda_2 v_2}{\lambda_1 v_1 - \lambda_2 v_2} \right) \quad (5)$$

## Neutrinos and Dark Matter

### Neutrino Failure

The paper proposes its main application in explaining “anomalously small” neutrino masses. It suggests the scalar ( $M$ ) and the pseudoscalar ( $\tilde{M}$ ) mass contributions, which arise from  $\phi_1$  and  $\phi_2$  respectively, nearly cancel each other out.

$$m_\nu = \frac{1}{2\sqrt{2}}(\lambda_1 v_1 - \lambda_2 v_2) \quad (6)$$

However, the paper’s derivation contains a fundamental contradiction. It states that the neutrino field has no right-chiral component (Eq. 26:  $\frac{1}{2}(1 + \gamma^5)\nu = 0$ ). As is well-known, a Dirac mass term (both scalar and pseudoscalar) requires both chiral components to be non-zero.

### Dark Matter Candidate

- The pseudoscalar Higgs field,  $\phi_2$ , is proposed as a Dark Matter candidate.
- It couples to Standard Model particles only via this pseudoscalar Yukawa coupling, primarily interacting with neutrinos.
- This makes it massive, long-lived, and “dark,” satisfying the requirements for a WIMP-like particle.

## References

- [1] V. Bargmann and E. P. Wigner. “Group Theoretical Discussion Of Relativistic Wave Equations”. In: *Part I: Particles and Fields. Part II: Foundations of Quantum Mechanics*. Springer Berlin Heidelberg, 1997, pp. 82–94. ISBN: 9783662092033. DOI: 10.1007/978-3-662-09203-3\_6.
- [2] T. B. Watson and Z. E. Musielak. “Chiral Symmetry In Dirac Equation And Its Effects On Neutrino Masses And Dark Matter”. In: *International Journal of Modern Physics A* 35.30 (Oct. 2020), p. 2050189. ISSN: 1793-656X. DOI: 10.1142/s0217751x20501894.
- [3] Timothy Watson and Zdzislaw Musielak. “Chiral Dirac Equation And Its Spacetime And Cpt Symmetries”. In: *Symmetry* 13.9 (Sept. 2021), p. 1608. ISSN: 2073-8994. DOI: 10.3390/sym13091608.
- [4] Timothy Blake Watson. “Chirality In Quantum Field Theory And Its Role In The Standard Model And Beyond”. PhD thesis. University of Texas at Arlington, 2022. URL: [https://mavmatrix.uta.edu/physics\\_dissertations/122](https://mavmatrix.uta.edu/physics_dissertations/122).