

Polarimetric signatures of a potential binary supermassive black hole in Mrk 231.


BIEDERMANN Julie

Astronomical Observatory of Strasbourg, France
29 June - 4 July 2026

Introduction - Markarian 231 (Mrk 231)

Mrk 231 - unique spectrum


- **Quasar**
  - ➡ z = 0.042 (~ 175 Mpc)
  - ➡ Ultra luminous infrared galaxy
- **Unique UV-optical spectrum**
  - ➡ Cut off ~ 2400 Å



Mrk 231 UV-IR spectrum, *Leighly et al. (2014)*

Mrk 231 - unique spectrum

- **Quasar**
  - ➡ z = 0.042 (~ 175 Mpc)
  - ➡ Ultra luminous infrared galaxy
- **Unique UV-optical spectrum**
  - ➡ Cut off ~ 2400 Å




Mrk 231 UV-IR spectrum, *Leighly et al. (2014)*

Introduction - Markarian 231 (Mrk 231)

Mrk 231 - unique polarimetry

- **Quasar**
  - ➡ z = 0.042 (~ 175 Mpc)
  - ➡ Ultra luminous infrared galaxy
- **Unique UV-optical spectrum**
  - ➡ Cut off ~ 2400 Å
- **Unique UV- optical polarization properties**
  - ➡ Wavelength-dependent polarization **degree**
  - ➡ Rotation of the polarization **angle**

➡ **Two sources ?**



Spectropolarimetric data of Mrk 231, HST/FOS, *Smith et al. (1995)*


Introduction - Markarian 231 (Mrk 231)

Mrk 231 - Hypothesis for a supermassive black holes binary (SMBH) - Yan and al (2015)

- Binary SMBH in the core of Mrk 231
  - ➡ To explain the cut off in the UV total flux spectrum

- **GOAL**
  - ➡ Reproduce the Spectral Energy Distribution (SED)
  - ➡ Analyse the response of the model in polarization

➡ **Never studied !**




Accretion disk diagram of a supermassive black hole binary system, *Yan et al. (2015)*

Modeling Spectral Energy distribution - Simple SMBH

SED for 1 SMBH

- Geometrically thin and optically thick disk model
- Multi temperature blackbody radiation
- Outer radius - critical accretion rate $\dot{M}_+$
  Disk becomes unstable : $\dot{M} > \dot{M}_+$ *Lacosta (2016)*




LOG-LOG critical accretion rate as a function of the normalized radius
Biedermann et al. (2025)

Modeling Spectral Energy distribution - Simple SMBH

SED for 1 SMBH

- Geometrically thin and optically thick disk model
- Multi temperature blackbody radiation
- Outer radius - critical accretion rate $\dot{M}_+$
  Disk becomes unstable : $\dot{M} > \dot{M}_+$ *Lacosta (2016)*
- SED for a simple SMBH

$$\boldsymbol{\theta} \equiv \begin{pmatrix} M_* \\ \alpha \\ p \\ \theta \end{pmatrix} \begin{matrix} \longleftarrow \text{SMBH mass} \\ \longleftarrow \text{Viscosity} \\ \longleftarrow \text{Accretion parameter} \\ \longleftarrow \text{Inclination angle} \end{matrix}$$



LOG-LOG Flux a function of the wavelength $M_* = 1.5 \times 10^8 M_\odot$
Biedermann et al. (2025)

Modeling Spectral Energy distribution - Binary SMBH

SED for 2 SMBHs


➡ Let's add a secondary SMBH

Primary larger SMBH

- Outer radius - critical accretion rate $\dot{M}_{+}$
- Inner radius - ~~ISCO~~ → Hill radius

Secondary smaller SMBH

- Outer radius - Roche lobe radius
- Inner radius - ISCO radius



Accretion disk diagram of a supermassive black hole binary system, inspired from *Yan et al. (2015)*

Modeling Spectral Energy distribution - Binary SMBH

SED for 2 SMBHs

➡ Let's add a secondary SMBH

Primary larger SMBH

- Outer radius - critical accretion rate $\dot{M}$
- Inner radius - ~~ISCO~~ → Hill radius

Secondary smaller SMBH

- Outer radius - Roche lobe radius
- Inner radius - ISCO radius


➡ SED for a binary SMBH

Roche lobe fraction

Semi major axis

Primary SMBH Secondary SMBH

$$\boldsymbol{\theta} \equiv \{M_{*,p}, \alpha_p, p_p, \theta_p \; ; \; M_{*,s}, \alpha_s, p_s, \theta_s \; ; \; f_{r,s}, a_{BBH}\}$$




LOG-LOG Flux a function of the wavelength for a binary SED
Biedermann et al. (2025)

Modeling Spectral Energy distribution - Total SED of the quasar

Total SED

- Archival data
  - Flux with corresponding polarimetric observations

- Reproduce the observations
  - Binary SED




Modeling Spectral Energy distribution - Total SED of the quasar

Total SED

- Archival data
  - Flux with corresponding polarimetric observations

- Reproduce the observations
  - Binary SED
  - Host galaxy - *Polletta et al (2007)*
  - Torus - *Siebenmorgen et al (2015)*



Modeling Spectral Energy distribution - Total SED of the quasar


Total SED

➡ Archival data
- Flux with corresponding polarimetric observations

➡ Reproduce the observations
- Binary SED
- Host galaxy - *Polletta et al (2007)*
- Torus - *Siebenmorgen et al (2015)*

➡ Best fit total Spectral Energy Distribution (SED)

➡ **A step further : what is the response of the model in polarization ?**




Foreword - What is polarization ?

➡ Orientation of the electric field $\vec{E}$

➡ Stokes formalism

$$\vec{S} = \begin{pmatrix} I \\ Q \\ U \\ V \end{pmatrix}$$

I : Total intensity
Q : Linear polarization $0° \,|\, 90°$
U : Linear polarization $\pm 45°$
V : Circular polarization



➡ Polarization adds 2 additional informations to the intensity

- Polarization degree $P\,[\%] = \dfrac{\sqrt{Q^2 + U^2 + V^2}}{I}$
- Polarization angle $\theta = \dfrac{1}{2} arctan\left(\dfrac{U}{Q}\right)$

Polarimetry

Polarimetric analysis

➡ For each component of the SED, a best fit polarization value is assigned and the Stokes fluxes are summed.


Polarimetry - Degree of polarization

Polarimetric analysis

➡ For each component of the SED, a best fit polarization value is assigned and the Stokes fluxes are summed.

Polarization degree

$$p = \frac{\sqrt{Q^2 + U^2}}{I}$$



Degree of polarization as a function of Log wavelength
Biedermann et al. (2025)

	Primary Black hole	Secondary Black hole	Host galaxy	Torus
P (%)	23.0 ± 2.7	3.1 ± 0.3	0	0.9 ± 0.1


Polarimetry - Angle of polarization

Polarimetric analysis

➡ For each component of the SED, a best fit polarization value is assigned and the Stokes fluxes are summed.

Polarization degree $p = \frac{\sqrt{Q^2 + U^2}}{I}$

Polarization angle $\theta = \frac{1}{2} arctan\left(\frac{U}{Q}\right)$



Angle of polarization as a function of Log wavelength
Biedermann et al. (2025)

	Primary Black hole	Secondary Black hole	Host galaxy	Torus
P (%)	23.0 ± 2.7	3.1 ± 0.3	0	0.9 ± 0.1
PA (°)	95.9 ± 0.2	133.8 ± 1.5	0	112.1 ± 2.1

Polarimetry - Polarized flux

Polarimetric analysis

➡ For each component of the SED, a best fit polarization value is assigned and the Stokes fluxes are summed.

Polarization degree


$$p = \frac{\sqrt{Q^2 + U^2}}{I}$$

Polarization angle

$$\theta = \frac{1}{2} arctan\left(\frac{U}{Q}\right)$$

Polarized flux

➡ Strong prediction for a second bump in FUV



LOG-LOG polarized Flux as a function of the wavelength
Biedermann et al. (2025)

	Primary Black hole	Secondary Black hole	Host galaxy	Torus
P (%)	23.0 ± 2.7	3.1 ± 0.3	0	0.9 ± 0.1
PA (°)	95.9 ± 0.2	133.8 ± 1.5	0	112.1 ± 2.1

Polarimetry - Polarized flux

Polarimetric analysis

➡ For each component of the SED, a best fit polarization value is assigned and the Stokes fluxes are summed.

Polarization degree

$$p = \frac{\sqrt{Q^2 + U^2}}{I}$$


Polarization angle

$$\theta = \frac{1}{2} arctan\left(\frac{U}{Q}\right)$$

Polarized flux

➡ Strong prediction for a second bump in FUV

Model reproduces three independent measurements



LOG-LOG polarized Flux as a function of the wavelength
Biedermann et al. (2025)

	Primary Black hole	Secondary Black hole	Host galaxy	Torus
P (%)	23.0 ± 2.7	3.1 ± 0.3	0	0.9 ± 0.1
PA (°)	95.9 ± 0.2	133.8 ± 1.5	0	112.1 ± 2.1

In Summary - Probing binary SMBH in quasar through spectropolarimetry

- ➡ Polarization is a strong tool to probe the inner structure of quasars
- ➡ In this work, we reconstruct the total SED of Mrk 231 using a simple binary SMBH model
- ➡ The binary SMBH model is in agreement with our spectropolarimetric data for Mrk 231
  - It reproduces well three independent measurements (polarized flux, polarization degree, polarization angle)
- ➡ Spectropolarimetric signatures: a second bump in the polarized flux and breaks in the total flux (UV)
- ➡ It opens promising perspectives for future UV spectropolarimeters, such as the POLLUX like instrument proposed for the Habitable Worlds Observatory

THANK YOU

FOR YOUR ATTENTION

Questions ?

Summary - Probing binary SMBH in quasars through spectropolarimetry

➡ Recreate the total SED of Mrk 231 using a simple binary SMBH model

➡ Binary SMBH model in agreement with our spectropolarimetric data

➡ Next step to test the binary hypothesis

✓ To probe the variation in time of Lyα profile

✓ To confirm/reject the presence of a second bump in FUV most easily detectable in polarized light

($\lambda_{rest} \sim 1460 - 2350$ Å)




➡ Future high-sensitivity UV spectropolarimeters such as the instrument like POLLUX proposed for the Habitable Worlds Observatory

THANK YOU

FOR YOUR ATTENTION

Questions ?

BACK SLIDE - Pollux for HWO



Habitable Worlds Observatory Telescope (HWO) *Artist's concept NASA*


➡ HWO : futur space mission composed of 4 instruments

- ✓ Coronagraph ~ 200 – 1800 [nm]
- ✓High resolution imager ~ 200 – 2500 [nm]
- ✓UV multi object spectrograph ~ 100 – 1000 [nm]
- ✓ ?

➡ POLLUX →candidate European instrument

- ✓ UV spectrometer with polarimetric capabilities
- ✓ High resolution spectrograph (R>70000)
- ✓ Continuous and simultaneous coverage from FUV to the NIR
- ✓Three pollux science working groups (Stars, (Exo)planets, cosmos ecosystems)

BACK SLIDE - Binary configuration



BACK SLIDE - Instability criterion


- Theoretical radius - Toomre parameter indicates a local gravitational instability in the disc
  - Disk become unstable : $\dot{M} > \dot{M}_{+}$ *Lacosta (2016)*

$$\dot{M}_{+} = 8.07 \times 10^{15} \alpha_{0.1}^{-0.01} \left(\frac{r}{10^{10}[cm]} \right)^{2.64} \left(\frac{M_*}{M_\odot} \right)^{-0.89}$$

$$\dot{M} = \alpha p \dot{M}_{Eddington}$$

- Black hole masse : M_*
- Viscosity : α
- Accretion parameter : p
- Radius of the accretion disk : r
- Eddington accretion rate : $\dot{M}_{Eddington}$

- $\dot{M}$ is linked to thermal instabilities



BACK SLIDE - Radius

❖Inner radius for the primary SMBH

$$R_{int,p} = \frac{a_{BBH}}{1+q} + a_{BBH}\left(\frac{M_{*,s}}{3M_{*,p}}\right)^{\frac{1}{3}}$$


Yan et al (2015)

❖Outer radius for the secondary SMBH

$$R_{out,s} = f_{r,s} \times \left(\frac{0.49 a_{BBH} q^{\frac{2}{3}}}{0.6 q^{\frac{2}{3}} + log(1+q^{\frac{1}{3}})}\right)$$

Yan et al (2015)

$$R_{int,s} = r_s$$



Roche lobe fraction

$$\boldsymbol{\theta} \equiv \{\underbrace{M_{*,p}, \alpha_p, p_p, \theta_p}_{\text{Primary SMBH}} ; \underbrace{M_{*,s}, \alpha_s, p_s, \theta_s}_{\text{Secondary SMBH}} ; f_{r,s}, a_{BBH}\}$$

Semi major axis