

Graphs in Machine Learning

Daniele Calandriello

DeepMind Paris, France

Collaborators: Achraf Azize,
Michal Valko

Partially based on material by: Gary Miller,
Mikhail Belkin, Branislav Kveton,
Doyle & Schnell, Daniel Spielman



Previous Lecture

- ▶ spectral graph theory
- ▶ Laplacians and their properties
 - ▶ symmetric and asymmetric normalization
 - ▶ random walks
- ▶ geometry of the data and the connectivity
- ▶ spectral clustering

This Lecture

- ▶ manifold learning with Laplacians eigenmaps
- ▶ inductive and transductive semi-supervised learning
- ▶ graph-based semi-supervised learning
- ▶ graph based manifold regularization

$\mathbb{R}^d \rightarrow \mathbb{R}^m$

manifold learning

...discworld

Manifold Learning: Recap

problem: definition reduction/manifold learning

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d find $\{\mathbf{y}_i\}_{i=1}^N$ in \mathbb{R}^m , where $m \ll d$.

Manifold Learning: Recap

problem: definition reduction/manifold learning

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d find $\{\mathbf{y}_i\}_{i=1}^N$ in \mathbb{R}^m , where $m \ll d$.

- ▶ What do we know about the **dimensionality reduction**
 - ▶ representation/visualization (2D or 3D)
 - ▶ an old example: globe to a map
 - ▶ feature extraction

Manifold Learning: Recap

problem: definition reduction/manifold learning

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d find $\{\mathbf{y}_i\}_{i=1}^N$ in \mathbb{R}^m , where $m \ll d$.

- ▶ What do we know about the **dimensionality reduction**
 - ▶ representation/visualization (2D or 3D)
 - ▶ an old example: globe to a map
 - ▶ feature extraction
 - ▶ often assuming $\mathcal{M} \subset \mathbb{R}^d$

Manifold Learning: Recap

problem: definition reduction/manifold learning

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d find $\{\mathbf{y}_i\}_{i=1}^N$ in \mathbb{R}^m , where $m \ll d$.

- ▶ What do we know about the **dimensionality reduction**
 - ▶ representation/visualization (2D or 3D)
 - ▶ an old example: globe to a map
 - ▶ feature extraction
 - ▶ often assuming $\mathcal{M} \subset \mathbb{R}^d$
 - ▶ linear vs. nonlinear dimensionality reduction


Manifold Learning: Linear vs. Non-linear

What do we know about linear vs. nonlinear methods?

Manifold Learning: Linear vs. Non-linear

What do we know about linear vs. nonlinear methods?




no linear or that
properly separates → we want non linear proj



PCA projection

LLE projection

IsoMap projection




color by dimension

not good


is better, more
non linear

We want 2D projection
to correctly clustering



Manifold Learning: Preserving (just) local distances



A manifold is a space that is non-linear but locally linear.



Manifold Learning: Preserving (just) local distances




euclidian distance $\leftarrow d(\mathbf{y}_i, \mathbf{y}_j) = d(\mathbf{x}_i, \mathbf{x}_j)$ only if $d(\mathbf{x}_i, \mathbf{x}_j)$ is small




+ →

Manifold Learning: Preserving (just) local distances



projection sent
from one



$$d(\mathbf{y}_i, \mathbf{y}_j) = d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{only if} \quad d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{is small}$$



1-D

$$\min \sum_{ij} w_{ij}(y_i - y_j)^2$$

Manifold Learning: Preserving (just) local distances





$$d(\mathbf{y}_i, \mathbf{y}_j) = d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{only if} \quad d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{is small}$$

$$\text{1-D} \quad \min \sum_{ij} w_{ij}(y_i - y_j)^2$$

Looks familiar?

Manifold Learning: Preserving (just) local distances





$$d(\mathbf{y}_i, \mathbf{y}_j) = d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{only if} \quad d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{is small}$$

$$\text{1-D} \quad \min \sum_{ij} w_{ij}(y_i - y_j)^2 \quad \approx \min \sum_{ij} w_{ij}(f_i - f_j)^2$$

Looks familiar?

Manifold Learning: Preserving (just) local distances



$$d(\mathbf{y}_i, \mathbf{y}_j) = d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{only if} \quad d(\mathbf{x}_i, \mathbf{x}_j) \quad \text{is small}$$

1-D

$$\min \sum_{ij} w_{ij}(y_i - y_j)^2 \approx \min \sum_{ij} w_{ij}(f_i - f_j)^2$$

Looks familiar?

m-D

$$\min_{\mathbf{y}} \sum_{ij} w_{ij} \|\mathbf{y}_i - \mathbf{y}_j\|^2$$

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

The meaning of the constraints is similar as for spectral clustering:

$\mathbf{f}^T \mathbf{D} \mathbf{f} = 1$ is for

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

The meaning of the constraints is similar as for spectral clustering:

$\mathbf{f}^T \mathbf{D} \mathbf{f} = 1$ is for scaling

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

The meaning of the constraints is similar as for spectral clustering:

$\mathbf{f}^T \mathbf{D} \mathbf{f} = 1$ is for scaling

$\mathbf{f}^T \mathbf{D} \mathbf{1} = 0$ is to

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

The meaning of the constraints is similar as for spectral clustering:

$\mathbf{f}^T \mathbf{D} \mathbf{f} = 1$ is for scaling

$\mathbf{f}^T \mathbf{D} \mathbf{1} = 0$ is to not get \mathbf{v}_1

Manifold Learning: Laplacian Eigenmaps to 1D

Laplacian Eigenmaps 1D objective

$$\min_{\mathbf{f}} \mathbf{f}^T \mathbf{L} \mathbf{f} \quad \text{s.t.} \quad f_i \in \mathbb{R}, \quad \mathbf{f}^T \mathbf{D} \mathbf{1} = 0, \quad \mathbf{f}^T \mathbf{D} \mathbf{f} = 1$$

The meaning of the constraints is similar as for spectral clustering:

$\mathbf{f}^T \mathbf{D} \mathbf{f} = 1$ is for scaling

$\mathbf{f}^T \mathbf{D} \mathbf{1} = 0$ is to not get \mathbf{v}_1

What is the solution?

Manifold Learning: Laplacian Eigenmaps

Step 1: Solve generalized eigenproblem:

$$\mathbf{Lf} = \lambda \mathbf{Df}$$

Manifold Learning: Laplacian Eigenmaps

Step 1: Solve generalized eigenproblem:

$$\mathbf{Lf} = \lambda \mathbf{Df}$$

Step 2: Assign m new coordinates:

$$\mathbf{x}_i \mapsto (f_2(i), \dots, f_{m+1}(i)) = \mathbf{y}_i$$

Manifold Learning: Laplacian Eigenmaps

Step 1: Solve generalized eigenproblem:

$$\mathbf{Lf} = \lambda \mathbf{Df}$$

Step 2: Assign m new coordinates:

$$\mathbf{x}_i \mapsto (f_2(i), \dots, f_{m+1}(i)) = \mathbf{y}_i$$

Note₁: we need to get $m + 1$ smallest eigenvectors

Manifold Learning: Laplacian Eigenmaps

Step 1: Solve generalized eigenproblem:

$$\mathbf{Lf} = \lambda \mathbf{Df}$$

Step 2: Assign m new coordinates:


$$\mathbf{x}_i \mapsto (f_2(i), \dots, f_{m+1}(i)) = \mathbf{y}_i$$

Note₁: we need to get $m + 1$ smallest eigenvectors

Note₂: \mathbf{f}_1 is useless

http://web.cse.ohio-state.edu/~mbelkin/papers/LEM_NC_03.pdf

Manifold Learning: Example



<http://www.mathworks.com/matlabcentral/fileexchange/>



36141-laplacian-eigenmap---diffusion-map---manifold-learning

SSL

semi-supervised learning



...our running example for learning
with graphs

Semi-supervised learning: How is it possible?



Take unlabeled data
and try
to associate
unlabeled
data to
labelled data

Semi-supervised learning: How is it possible?



This is how children learn! hypothesis

Semi-supervised learning (SSL) *labelled data*

SSL problem: definition

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d and $\{y_i\}_{i=1}^{n_l}$, with $n_l \ll N$, find $\{y_i\}_{i=n_l+1}^N$ (**transductive**) or
find f predicting y well beyond that (**inductive**). *Total nb of data*

Semi-supervised learning (SSL)

SSL problem: definition

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d and $\{y_i\}_{i=1}^{n_l}$, with $n_l \ll N$, find $\{y_i\}_{i=n_l+1}^N$ (**transductive**) or find f predicting y well beyond that (**inductive**). \hookrightarrow unlabeled data

Some facts about SSL

- ▶ assumes that the unlabeled data is useful

Semi-supervised learning (SSL)

SSL problem: definition

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d and $\{y_i\}_{i=1}^{n_l}$, with $n_l \ll N$, find $\{y_i\}_{i=n_l+1}^N$ (**transductive**) or find f predicting y well beyond that (**inductive**).

Some facts about SSL

- ▶ assumes that the unlabeled data is useful
- ▶ works with data geometry assumptions
 - ▶ cluster assumption — low-density separation
 - ▶ smoothness assumptions, generative models, ...
 - ▶ manifold assumption

Semi-supervised learning (SSL)

SSL problem: definition

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d and $\{y_i\}_{i=1}^{n_l}$, with $n_l \ll N$, find $\{y_i\}_{i=n_l+1}^N$ (**transductive**) or find f predicting y well beyond that (**inductive**).

Some facts about SSL

- ▶ assumes that the unlabeled data is useful
- ▶ works with data geometry assumptions
 - ▶ cluster assumption — low-density separation
 - ▶ smoothness assumptions, generative models, ...
 - ▶ manifold assumption
- ▶ now it helps now, now it does not (sic)
 - ▶ provable cases when it helps

Semi-supervised learning (SSL)

SSL problem: definition

Given $\{\mathbf{x}_i\}_{i=1}^N$ from \mathbb{R}^d and $\{y_i\}_{i=1}^{n_l}$, with $n_l \ll N$, find $\{y_i\}_{i=n_l+1}^N$ (**transductive**) or find f predicting y well beyond that (**inductive**).

Some facts about SSL

- ▶ assumes that the unlabeled data is useful
- ▶ works with data geometry assumptions
 - ▶ cluster assumption — low-density separation
 - ▶ smoothness assumptions, generative models, ...
 - ▶ manifold assumption
- ▶ now it helps now, now it does not (sic)
 - ▶ provable cases when it helps
- ▶ inductive or transductive/out-of-sample extension

<http://olivier.chapelle.cc/ssl-book/discussion.pdf>

SSL: Overview: Self-Training

SSL: Self-Training

Input: $\mathcal{L} = \{\mathbf{x}_i, y_i\}_{i=1}^{n_l}$ and $\mathcal{U} = \{\mathbf{x}_i\}_{i=n_l+1}^N$

Repeat:

- ▶ train f using \mathcal{L}
- ▶ apply f to (some) \mathcal{U} and add them to \mathcal{L}

SSL: Overview: Self-Training

SSL: Self-Training

Input: $\mathcal{L} = \{\mathbf{x}_i, y_i\}_{i=1}^{n_l}$ and $\mathcal{U} = \{\mathbf{x}_i\}_{i=n_l+1}^N$

Repeat:

- ▶ train f using \mathcal{L}
- ▶ apply f to (some) \mathcal{U} and add them to \mathcal{L}

What are the properties of self-training?

SSL: Overview: Self-Training

SSL: Self-Training

Input: $\mathcal{L} = \{\mathbf{x}_i, y_i\}_{i=1}^{n_l}$ and $\mathcal{U} = \{\mathbf{x}_i\}_{i=n_l+1}^N$


Repeat:

- ▶ train f using \mathcal{L}
- ▶ apply f to (some) \mathcal{U} and add them to \mathcal{L}


What are the properties of self-training?

- ▶ its a wrapper method
- ▶ heavily depends on the internal classifier
- ▶ some theory exist for specific classifiers
- ▶ nobody uses it anymore (\neq self supervised)
- ▶ errors propagate (unless the clusters are well separated)


SSL: Self-Training (Good Case)




(a) Iteration 1



(b) Iteration 25




(c) Iteration 74




(d) Final labeling of all instances


SSL: Self-Training (Bad Case)




(a)



(b)



(c)



(d)

SSL(\mathcal{G})

semi-supervised learning with
graphs and harmonic functions


...our running example for learning with graphs

SSL with Graphs: Prehistory


Blum/Chawla: Learning from Labeled and Unlabeled Data using Graph Mincuts

<http://www.aladdin.cs.cmu.edu/papers/pdfs/y2001/mincut.pdf>

*following some insights from vision research in 1980s




SSL with Graphs: MinCut



MinCut SSL: an idea similar to MinCut clustering


SSL with Graphs: MinCut



MinCut SSL: an idea similar to MinCut clustering

Where is the link?


SSL with Graphs: MinCut



MinCut SSL: an idea similar to MinCut clustering

Where is the link? connected classes, not necessarily compact

SSL with Graphs: MinCut




MinCut SSL: an idea similar to MinCut clustering

Where is the link? connected classes, not necessarily compact

What is the formal statement?

SSL with Graphs: MinCut




MinCut SSL: an idea similar to MinCut clustering

Where is the link? connected classes, not necessarily compact

What is the formal statement? We look for $f(\mathbf{x}) \in \{\pm 1\}$

$$\text{cut} = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \Omega(f)$$

SSL with Graphs: MinCut



MinCut SSL: an idea similar to MinCut clustering


Where is the link? connected classes, not necessarily compact

What is the formal statement? We look for $f(\mathbf{x}) \in \{\pm 1\}$

$$\text{cut} = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \Omega(f)$$

Why $(f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$ and not $|f(\mathbf{x}_i) - f(\mathbf{x}_j)|$?

SSL with Graphs: MinCut



MinCut SSL: an idea similar to MinCut clustering

Where is the link? connected classes, not necessarily compact

What is the formal statement? We look for $f(\mathbf{x}) \in \{\pm 1\}$

$$\text{cut} = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \Omega(f)$$

Why $(f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$ and not $|f(\mathbf{x}_i) - f(\mathbf{x}_j)|$? It does not matter.

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_I+n_U} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Clustering was unsupervised, here we have supervised data.

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_I+n_U} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Clustering was unsupervised, here we have supervised data.

Recall the general objective-function framework:

$$\min_{\mathbf{w}, b} \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(\mathbf{f})$$

orig loss *min cut loss*

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Clustering was unsupervised, here we have supervised data.

Recall the general objective-function framework:

$$\min_{\mathbf{w}, b} \sum_i^{n_l} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(\mathbf{f})$$

It would be nice if we match the prediction on labeled data:

$$V(\mathbf{x}, y, f(\mathbf{x}))$$

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Clustering was unsupervised, here we have supervised data.

Recall the general objective-function framework:

$$\min_{\mathbf{w}, b} \sum_i^{n_l} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(\mathbf{f})$$

It would be nice if we match the prediction on labeled data:

$$V(\mathbf{x}, y, f(\mathbf{x})) = \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2$$

SSL with Graphs: MinCut

We look for $f(\mathbf{x}) \in \{\pm 1\}$ to minimize the cut $\Omega(\mathbf{f})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_I+n_U} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Clustering was unsupervised, here we have supervised data.

Recall the general objective-function framework:

$$\min_{\mathbf{w}, b} \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(\mathbf{f})$$

It would be nice if we match the prediction on labeled data:

$$V(\mathbf{x}, y, f(\mathbf{x})) = \infty \sum_{i=1}^{n_I} (f(\mathbf{x}_i) - y_i)^2$$

SSL with Graphs: MinCut

Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

SSL with Graphs: MinCut

Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

SSL with Graphs: MinCut

Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it?

SSL with Graphs: MinCut

Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

SSL with Graphs: MinCut

Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \frac{1}{2} \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

Are we happy?

SSL with Graphs: MinCut


Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

Are we happy?



SSL with Graphs: MinCut


Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \frac{1}{2} \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

Are we happy?



There are six solutions.

SSL with Graphs: MinCut


Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

Are we happy?



There are six solutions. All equivalent.

SSL with Graphs: MinCut


Final objective function:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

This is an integer program :(

Can we solve it? It still just MinCut.

Are we happy?



There are six solutions. All equivalent.

We need a better way to reflect the confidence.

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Locally, form extra constraints to get a unique solution

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Idea 1: Look for a **unique** solution.

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Idea 1: Look for a **unique** solution.

Idea 2: Find a smooth one.

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Idea 1: Look for a **unique** solution.

Idea 2: Find a smooth one. (**harmonic** solution) ↗ much in physics to characterize
 $\text{In} + \text{Out} = 0$. Appears when have energy flow.

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Idea 1: Look for a **unique** solution.

Idea 2: Find a smooth one. (**harmonic** solution)

Harmonic SSL

1): As before, we constrain f to match the supervised data:

$$f(\mathbf{x}_i) = y_i \quad \forall i \in \{1, \dots, n_I\}$$

SSL with Graphs: Harmonic Functions

Zhu/Ghahramani/Lafferty: Semi-Supervised Learning Using Gaussian Fields and Harmonic Functions (ICML 2013) <http://mlg.eng.cam.ac.uk/zoubin/papers/zgl.pdf>

*a seminal paper that convinced people to use graphs for SSL

Idea 1: Look for a **unique** solution.

Idea 2: Find a smooth one. (**harmonic** solution)

Harmonic SSL

1): As before, we constrain f to match the supervised data:

$$f(\mathbf{x}_i) = y_i \quad \forall i \in \{1, \dots, n_l\}$$

2): We enforce the solution f to be harmonic:

$$f(\mathbf{x}_i) = \frac{\sum_{j \sim i} f(\mathbf{x}_j) w_{ij}}{\sum_{j \sim i} w_{ij}} \quad \forall i \in \{n_l + 1, \dots, n_u + n_l\}$$

SSL with Graphs: Harmonic Functions

The harmonic solution is obtained from the mincut one ...

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

SSL with Graphs: Harmonic Functions

The harmonic solution is obtained from the mincut one ...

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

...if we just relax the integer constraints to be real ...

$$\min_{\mathbf{f} \in \mathbb{R}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

SSL with Graphs: Harmonic Functions

The harmonic solution is obtained from the mincut one ...

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

...if we just relax the integer constraints to be real ...

$$\min_{\mathbf{f} \in \mathbb{R}^{n_l+n_u}} \infty \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

...or equivalently (note that $f(\mathbf{x}_i) = f_i$) ...

$$\min_{\mathbf{f} \in \mathbb{R}^{n_l+n_u}} \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

$$s.t. \quad y_i = f(\mathbf{x}_i) \quad \forall i = 1, \dots, n_l$$

SSL with Graphs: Harmonic Functions

Properties of the relaxation from ± 1 to \mathbb{R}

- ▶ there is a closed form solution for f
- ▶ this solution is unique
- ▶ globally optimal

SSL with Graphs: Harmonic Functions

Properties of the relaxation from ± 1 to \mathbb{R}


- ▶ there is a closed form solution for f
- ▶ this solution is unique
- ▶ globally optimal
- ▶ $f(\mathbf{x}_i)$ may not be discrete
 - ▶ but we can threshold it

SSL with Graphs: Harmonic Functions


Properties of the relaxation from ± 1 to \mathbb{R}

- ▶ there is a closed form solution for f
- ▶ this solution is unique
- ▶ globally optimal
- ▶ $f(x_i)$ may not be discrete
 - ▶ but we can threshold it
- ▶ electric-network interpretation
- ▶ random-walk interpretation

SSL with Graphs: Harmonic Functions




(a) The electric network interpretation




(b) The random walk interpretation

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation




(b) The random walk interpretation


Random walk interpretation:

- 1) start from the vertex you want to label and randomly walk

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation




(b) The random walk interpretation

Random walk interpretation:


1) start from the vertex you want to label and randomly walk

$$2) P(j|i) = \frac{w_{ij}}{\sum_k w_{ik}}$$

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation




(b) The random walk interpretation

Random walk interpretation:


1) start from the vertex you want to label and randomly walk

$$2) P(j|i) = \frac{w_{ij}}{\sum_k w_{ik}} \quad \equiv \quad \mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$$

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation




(b) The random walk interpretation


Random walk interpretation:

- 1) start from the vertex you want to label and randomly walk
- 2) $P(j|i) = \frac{w_{ij}}{\sum_k w_{ik}} \quad \equiv \quad \mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$
- 3) finish when a labeled vertex is hit

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation



(b) The random walk interpretation

Random walk interpretation:

1) start from the vertex you want to label and randomly walk


$$2) P(j|i) = \frac{w_{ij}}{\sum_k w_{ik}} \quad \equiv \quad \mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$$

3) finish when a labeled vertex is hit


absorbing random walk

$$f_i =$$

SSL with Graphs: Harmonic Functions



(a) The electric network interpretation



(b) The random walk interpretation

Random walk interpretation:

1) start from the vertex you want to label and randomly walk

$$2) P(j|i) = \frac{w_{ij}}{\sum_k w_{ik}} \quad \equiv \quad \mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$$

3) finish when a labeled vertex is hit

absorbing random walk

f_i = probability of reaching a positive labeled vertex

SSL with Graphs: Harmonic Functions

How to compute HS?

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration/propagation

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration/propagation

Step 1: Set $f(\mathbf{x}_i) = y_i$ for $i = 1, \dots, n_I$

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration/propagation

Step 1: Set $f(\mathbf{x}_i) = y_i$ for $i = 1, \dots, n_l$

Step 2: Propagate iteratively (only for unlabeled)

$$f(\mathbf{x}_i) \leftarrow \frac{\sum_{i \sim j} f(\mathbf{x}_j) w_{ij}}{\sum_{i \sim j} w_{ij}} \quad \forall i \in \{n_l + 1, \dots, n_u + n_l\}$$

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration/propagation

Step 1: Set $f(\mathbf{x}_i) = y_i$ for $i = 1, \dots, n_l$

Step 2: Propagate iteratively (only for unlabeled)

$$f(\mathbf{x}_i) \leftarrow \frac{\sum_{i \sim j} f(\mathbf{x}_j) w_{ij}}{\sum_{i \sim j} w_{ij}} \quad \forall i \in \{n_l + 1, \dots, n_u + n_l\}$$

Pros:

SSL with Graphs: Harmonic Functions

How to compute HS? **Option A:** iteration/propagation

Step 1: Set $f(\mathbf{x}_i) = y_i$ for $i = 1, \dots, n_l$

Step 2: Propagate iteratively (only for unlabeled)

$$f(\mathbf{x}_i) \leftarrow \frac{\sum_{i \sim j} f(\mathbf{x}_j) w_{ij}}{\sum_{i \sim j} w_{ij}} \quad \forall i \in \{n_l + 1, \dots, n_u + n_l\}$$

Pros:

- ▶ this will converge to the harmonic solution
- ▶ we can set the initial values for unlabeled nodes arbitrarily
- ▶ an interesting option for large-scale data

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

Define $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n_l+n_u})) = (f_1, \dots, f_{n_l+n_u})$

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

Define $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n_l+n_u})) = (f_1, \dots, f_{n_l+n_u})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

Define $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n_l+n_u})) = (f_1, \dots, f_{n_l+n_u})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

\mathbf{L} is a $(n_l + n_u) \times (n_l + n_u)$ matrix:

$$\mathbf{L} = \begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix}$$

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

Define $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n_l+n_u})) = (f_1, \dots, f_{n_l+n_u})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

\mathbf{L} is a $(n_l + n_u) \times (n_l + n_u)$ matrix:

$$\mathbf{L} = \begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix}$$

How to compute this **constrained** minimization problem?

SSL with Graphs: Harmonic Functions

How to compute HS? **Option B:** Closed form solution

Define $\mathbf{f} = (f(\mathbf{x}_1), \dots, f(\mathbf{x}_{n_l+n_u})) = (f_1, \dots, f_{n_l+n_u})$

$$\Omega(\mathbf{f}) = \sum_{i,j=1}^{n_l+n_u} w_{ij} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2 = \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

\mathbf{L} is a $(n_l + n_u) \times (n_l + n_u)$ matrix:

$$\mathbf{L} = \begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix}$$

How to compute this **constrained** minimization problem?

Yes, Lagrangian multipliers are an option, but . . .

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{II} & \mathbf{L}_{Iu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_I \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

\mathbf{f}_I is constrained to be \mathbf{y}_I and for \mathbf{f}_u

$$\mathbf{L}_{ul}\mathbf{f}_I + \mathbf{L}_{uu}\mathbf{f}_u = \mathbf{0}_u$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

\mathbf{f}_l is constrained to be \mathbf{y}_l and for \mathbf{f}_u

$$\mathbf{L}_{ul}\mathbf{f}_l + \mathbf{L}_{uu}\mathbf{f}_u = \mathbf{0}_u$$

...from which we get

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{ul}\mathbf{f}_l)$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

\mathbf{f}_l is constrained to be \mathbf{y}_l and for \mathbf{f}_u

$$\mathbf{L}_{ul}\mathbf{f}_l + \mathbf{L}_{uu}\mathbf{f}_u = \mathbf{0}_u$$

...from which we get

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{ul}\mathbf{f}_l) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{ul}\mathbf{f}_l).$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

\mathbf{f}_l is constrained to be \mathbf{y}_l and for \mathbf{f}_u

$$\mathbf{L}_{ul}\mathbf{f}_l + \mathbf{L}_{uu}\mathbf{f}_u = \mathbf{0}_u$$

...from which we get

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{ul}\mathbf{f}_l) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{ul}\mathbf{f}_l). \quad \mathbf{L}_{ul} = \mathbf{0} - \mathbf{W}_{ul}$$

SSL with Graphs: Harmonic Functions

Let us compute **harmonic** solution using **harmonic** property!

$$(\mathbf{L}\mathbf{f})_u = \mathbf{0}_u$$

In matrix notation

$$\begin{bmatrix} \mathbf{L}_{ll} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

\mathbf{f}_l is constrained to be \mathbf{y}_l and for \mathbf{f}_u

$$\mathbf{L}_{ul}\mathbf{f}_l + \mathbf{L}_{uu}\mathbf{f}_u = \mathbf{0}_u$$

...from which we get

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{ul}\mathbf{f}_l) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{ul}\mathbf{f}_l). \quad \mathbf{L}_{ul} = \mathbf{0} - \mathbf{W}_{ul}$$

Note that \mathbf{f}_u does not depend on \mathbf{L}_{ll} .

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I)$$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{uI}\mathbf{f}_I)$$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{uI}\mathbf{f}_I)$$

Note $\mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$ and $\mathbf{L}_{uu}^{-1} = (\mathbf{D}_{uu}(\mathbf{I} - \mathbf{P}_{uu}))^{-1} = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{D}_{uu}^{-1}$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{uI}\mathbf{f}_I)$$

Note $\mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$ and $\mathbf{L}_{uu}^{-1} = (\mathbf{D}_{uu}(\mathbf{I} - \mathbf{P}_{uu}))^{-1} = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{D}_{uu}^{-1}$

$$\mathbf{f}_u = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{P}_{uI}\mathbf{f}_I.$$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{ul}\mathbf{f}_l) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{ul}\mathbf{f}_l)$$

Note $\mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$ and $\mathbf{L}_{uu}^{-1} = (\mathbf{D}_{uu}(\mathbf{I} - \mathbf{P}_{uu}))^{-1} = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{D}_{uu}^{-1}$

$$\mathbf{f}_u = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{P}_{ul}\mathbf{f}_l.$$

Split the equation into +ve & -ve part:

$$f_i = (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{ul}\mathbf{f}_l$$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{uI}\mathbf{f}_I)$$

Note $\mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$ and $\mathbf{L}_{uu}^{-1} = (\mathbf{D}_{uu}(\mathbf{I} - \mathbf{P}_{uu}))^{-1} = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{D}_{uu}^{-1}$

$$\mathbf{f}_u = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{P}_{uI}\mathbf{f}_I.$$

Split the equation into +ve & -ve part:

$$\begin{aligned} f_i &= (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uI}\mathbf{f}_I \\ &= \underbrace{\sum_{j:y_j=1} (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uj}}_{p_i^{(+)}} - \underbrace{\sum_{j:y_j=-1} (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uj}}_{p_i^{(-)}} \end{aligned}$$

SSL with Graphs: Harmonic Functions

Can we see that this calculates the probability of a random walk?

$$\mathbf{f}_u = \mathbf{L}_{uu}^{-1}(-\mathbf{L}_{uI}\mathbf{f}_I) = \mathbf{L}_{uu}^{-1}(\mathbf{W}_{uI}\mathbf{f}_I)$$

Note $\mathbf{P} = \mathbf{D}^{-1}\mathbf{W}$ and $\mathbf{L}_{uu}^{-1} = (\mathbf{D}_{uu}(\mathbf{I} - \mathbf{P}_{uu}))^{-1} = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{D}_{uu}^{-1}$

$$\mathbf{f}_u = (\mathbf{I} - \mathbf{P}_{uu})^{-1}\mathbf{P}_{uI}\mathbf{f}_I.$$

Split the equation into +ve & -ve part:

$$\begin{aligned} f_i &= (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uI}\mathbf{f}_I \\ &= \underbrace{\sum_{j:y_j=1} (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uj}}_{p_i^{(+1)}} - \underbrace{\sum_{j:y_j=-1} (\mathbf{I} - \mathbf{P}_{uu})_{iu}^{-1}\mathbf{P}_{uj}}_{p_i^{(-1)}} \\ &= p_i^{(+1)} - p_i^{(-1)} \end{aligned}$$

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)}$$

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

We add a **sink** to the graph.

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

We add a **sink** to the graph.

sink = artificial label node with value 0

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

We add a **sink** to the graph.

sink = artificial label node with value 0

We connect it to every other vertex.

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

We add a **sink** to the graph.

sink = artificial label node with value 0

We connect it to every other vertex.

What will this do to our predictions?

SSL with Graphs: Regularized Harmonic Functions

$$f_i = p_i^{(+1)} - p_i^{(-1)} \implies f_i = \underbrace{|f_i|}_{\text{confidence}} \times \underbrace{\text{sgn}(f_i)}_{\text{label}}$$

What if a nasty outlier sneaks in?

The prediction for the outlier can be hyperconfident :(

How to control the confidence of the inference?

Allow the random walk to **die**!

We add a **sink** to the graph.

sink = artificial label node with value 0

We connect it to every other vertex.

What will this do to our predictions?

depends on the weight on the edges

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

Formally, to get the harmonic solution on the graph with sink ...

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

Formally, to get the harmonic solution on the graph with sink ...

$$\begin{bmatrix} \mathbf{L}_{ll} + \gamma_G \mathbf{I}_{n_l} & \mathbf{L}_{lu} & -\gamma_G \mathbf{1}_{n_l \times 1} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} + \gamma_G \mathbf{I}_{n_u} & -\gamma_G \mathbf{1}_{n_u \times 1} \\ -\gamma_G \mathbf{1}_{1 \times n_l} & -\gamma_G \mathbf{1}_{1 \times n_u} & n \gamma_G \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \\ 0 \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \\ \dots \end{bmatrix}$$

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

Formally, to get the harmonic solution on the graph with sink ...

$$\begin{bmatrix} \mathbf{L}_{II} + \gamma_G \mathbf{I}_{n_I} & \mathbf{L}_{IU} & -\gamma_G \mathbf{1}_{n_I \times 1} \\ \mathbf{L}_{UL} & \mathbf{L}_{UU} + \gamma_G \mathbf{I}_{n_U} & -\gamma_G \mathbf{1}_{n_U \times 1} \\ -\gamma_G \mathbf{1}_{1 \times n_I} & -\gamma_G \mathbf{1}_{1 \times n_U} & n \gamma_G \end{bmatrix} \begin{bmatrix} \mathbf{f}_I \\ \mathbf{f}_U \\ 0 \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \\ \dots \end{bmatrix}$$

$$\mathbf{L}_{UL} \mathbf{f}_I + (\mathbf{L}_{UU} + \gamma_G \mathbf{I}_{n_U}) \mathbf{f}_U = \mathbf{0}_u$$

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

Formally, to get the harmonic solution on the graph with sink ...

$$\begin{bmatrix} \mathbf{L}_{ll} + \gamma_G \mathbf{I}_{n_l} & \mathbf{L}_{lu} & -\gamma_G \mathbf{1}_{n_l \times 1} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} + \gamma_G \mathbf{I}_{n_u} & -\gamma_G \mathbf{1}_{n_u \times 1} \\ -\gamma_G \mathbf{1}_{1 \times n_l} & -\gamma_G \mathbf{1}_{1 \times n_u} & n \gamma_G \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \\ 0 \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \\ \dots \end{bmatrix}$$

$$\mathbf{L}_{ul} \mathbf{f}_l + (\mathbf{L}_{uu} + \gamma_G \mathbf{I}_{n_u}) \mathbf{f}_u = \mathbf{0}_u$$

...which is the same if we disregard the last column and row ...

$$\begin{bmatrix} \mathbf{L}_{ll} + \gamma_G \mathbf{I}_{n_l} & \mathbf{L}_{lu} \\ \mathbf{L}_{ul} & \mathbf{L}_{uu} + \gamma_G \mathbf{I}_{n_u} \end{bmatrix} \begin{bmatrix} \mathbf{f}_l \\ \mathbf{f}_u \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

SSL with Graphs: Regularized Harmonic Functions

How do we represent the sink in \mathbf{L} explicitly?

Formally, to get the harmonic solution on the graph with sink ...

$$\begin{bmatrix} \mathbf{L}_{II} + \gamma_G \mathbf{I}_{n_I} & \mathbf{L}_{IU} & -\gamma_G \mathbf{1}_{n_I \times 1} \\ \mathbf{L}_{UL} & \mathbf{L}_{UU} + \gamma_G \mathbf{I}_{n_U} & -\gamma_G \mathbf{1}_{n_U \times 1} \\ -\gamma_G \mathbf{1}_{1 \times n_I} & -\gamma_G \mathbf{1}_{1 \times n_U} & n \gamma_G \end{bmatrix} \begin{bmatrix} \mathbf{f}_I \\ \mathbf{f}_U \\ 0 \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \\ \dots \end{bmatrix}$$

$$\mathbf{L}_{UL} \mathbf{f}_I + (\mathbf{L}_{UU} + \gamma_G \mathbf{I}_{n_U}) \mathbf{f}_U = \mathbf{0}_u$$

...which is the same if we disregard the last column and row ...

$$\begin{bmatrix} \mathbf{L}_{II} + \gamma_G \mathbf{I}_{n_I} & \mathbf{L}_{IU} \\ \mathbf{L}_{UL} & \mathbf{L}_{UU} + \gamma_G \mathbf{I}_{n_U} \end{bmatrix} \begin{bmatrix} \mathbf{f}_I \\ \mathbf{f}_U \end{bmatrix} = \begin{bmatrix} \dots \\ \mathbf{0}_u \end{bmatrix}$$

...and therefore we simply add γ_G to the diagonal of $\mathbf{L}!$

SSL with Graphs: Regularized Harmonic Functions

How do we compute this **regularized** random walk?

SSL with Graphs: Regularized Harmonic Functions

How do we compute this **regularized** random walk?

$$\mathbf{f}_u = (\mathbf{L}_{uu} + \gamma_g \mathbf{I})^{-1} (\mathbf{W}_{uI} \mathbf{f}_I)$$

SSL with Graphs: Regularized Harmonic Functions

How do we compute this **regularized** random walk?

$$\mathbf{f}_u = (\mathbf{L}_{uu} + \gamma_g \mathbf{I})^{-1} (\mathbf{W}_{uI} \mathbf{f}_I)$$



How does γ_g influence HS?

SSL with Graphs: Regularized Harmonic Functions

How do we compute this **regularized** random walk?

$$\mathbf{f}_u = (\mathbf{L}_{uu} + \gamma_g \mathbf{I})^{-1} (\mathbf{W}_{ul} \mathbf{f}_l)$$

How does γ_g influence HS?





SSL with Graphs: Regularized Harmonic Functions

How do we compute this **regularized** random walk?

$$\mathbf{f}_u = (\mathbf{L}_{uu} + \gamma_g \mathbf{I})^{-1} (\mathbf{W}_{ul} \mathbf{f}_l)$$

How does γ_g influence HS?



What happens to sneaky outliers?

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

$$\min_{\mathbf{f} \in \mathbb{R}^{n_I+n_u}} \frac{1}{2} \sum_{i=1}^{n_I} (f(\mathbf{x}_i) - y_i)^2 + \lambda \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

$$\min_{\mathbf{f} \in \mathbb{R}^{n_I+n_U}} \frac{1}{2} \sum_{i=1}^{n_I} (f(\mathbf{x}_i) - y_i)^2 + \lambda \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

What if we do not really believe that $f(\mathbf{x}_i) = y_i, \forall i$?

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

$$\min_{\mathbf{f} \in \mathbb{R}^{n_I+n_u}} \frac{1}{2} \sum_{i=1}^{n_I} (f(\mathbf{x}_i) - y_i)^2 + \lambda \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

What if we do not really believe that $f(\mathbf{x}_i) = y_i, \forall i?$

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C} (\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

$$\min_{\mathbf{f} \in \mathbb{R}^{n_l+n_u}} \frac{1}{2} \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

What if we do not really believe that $f(\mathbf{x}_i) = y_i, \forall i?$

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C} (\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

\mathbf{C} is diagonal with $C_{ii} = \begin{cases} c_l & \text{for labeled examples} \\ c_u & \text{otherwise.} \end{cases}$

SSL with Graphs: Soft Harmonic Functions

Regularized HS objective with $\mathbf{Q} = \mathbf{L} + \gamma_g \mathbf{I}$:

$$\min_{\mathbf{f} \in \mathbb{R}^{n_l+n_u}} \frac{1}{2} \sum_{i=1}^{n_l} (f(\mathbf{x}_i) - y_i)^2 + \lambda \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

What if we do not really believe that $f(\mathbf{x}_i) = y_i, \forall i$?

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C} (\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

\mathbf{C} is diagonal with $C_{ii} = \begin{cases} c_l & \text{for labeled examples} \\ c_u & \text{otherwise.} \end{cases}$

$\mathbf{y} \equiv$ pseudo-targets with $y_i = \begin{cases} \text{true label} & \text{for labeled examples} \\ 0 & \text{otherwise.} \end{cases}$

SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C} (\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

Closed form **soft harmonic solution**:

SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Closed form **soft harmonic solution**:



$$\mathbf{f}^* = (\mathbf{C}^{-1}\mathbf{Q} + \mathbf{I})^{-1}\mathbf{y}$$

SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Closed form **soft harmonic solution**:

$$\mathbf{f}^* = (\mathbf{C}^{-1}\mathbf{Q} + \mathbf{I})^{-1}\mathbf{y}$$





SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Closed form **soft harmonic solution**:

$$\mathbf{f}^* = (\mathbf{C}^{-1}\mathbf{Q} + \mathbf{I})^{-1}\mathbf{y}$$





What are the differences between hard and soft?

SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Closed form **soft harmonic solution**:

$$\mathbf{f}^* = (\mathbf{C}^{-1}\mathbf{Q} + \mathbf{I})^{-1}\mathbf{y}$$



What are the differences between hard and soft?



Not much different in practice.

SSL with Graphs: Soft Harmonic Functions

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^n} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Closed form **soft harmonic solution**:

$$\mathbf{f}^* = (\mathbf{C}^{-1}\mathbf{Q} + \mathbf{I})^{-1}\mathbf{y}$$



What are the differences between hard and soft?

Not much different in practice.

Provable generalization guarantees for the soft one.

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point.

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\mathbf{f}_2^* - \mathbf{f}_1^*$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\mathbf{f}_2^* - \mathbf{f}_1^* = \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|A\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|A\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|A\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|\mathbf{A}\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \lambda_M(\mathbf{Q})$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|\mathbf{A}\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|\mathbf{A}\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Think about **stability** of this solution.

Consider two datasets differing in exactly one *labeled* point. $\mathcal{C}_1 = \mathbf{C}_1^{-1}\mathbf{Q} + \mathbf{I}$ and $\mathcal{C}_2 = \mathbf{C}_2^{-1}\mathbf{Q} + \mathbf{I}$

What is the maximal difference in the solutions?

$$\begin{aligned}\mathbf{f}_2^* - \mathbf{f}_1^* &= \mathcal{C}_2^{-1}\mathbf{y}_2 - \mathcal{C}_1^{-1}\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} - \mathcal{C}_2^{-1})\mathbf{y}_1 \\ &= \mathcal{C}_2^{-1}(\mathbf{y}_2 - \mathbf{y}_1) - (\mathcal{C}_1^{-1} [(\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1})\mathbf{Q}] \mathcal{C}_2^{-1})\mathbf{y}_1\end{aligned}$$

Note that $\mathbf{v} \in \mathbb{R}^{N \times 1}$, $\lambda_m(A)\|\mathbf{v}\|_2 \leq \|A\mathbf{v}\|_2 \leq \lambda_M(A)\|\mathbf{v}\|_2$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)\lambda_m(\mathcal{C}_1)}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)\lambda_m(\mathcal{C}_1)}$$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)\lambda_m(\mathcal{C}_1)}$$

Using $\lambda_m(\mathcal{C}) \geq \frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C})} + 1$

SSL with Graphs: Stability Bounds

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\lambda_m(\mathcal{C}_2)\lambda_m(\mathcal{C}_1)}$$

Using $\lambda_m(\mathcal{C}) \geq \frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C})} + 1$

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_2 \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$.

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right)\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right)\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\lambda_m(\mathbf{Q}) + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{Q})}{(\lambda_m(\mathbf{Q}) + 1)^2} \right]$$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\lambda_m(\mathbf{Q}) + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{Q})}{(\lambda_m(\mathbf{Q}) + 1)^2} \right]$$

\mathbf{Q} is reg. \mathbf{L} : $\lambda_m(\mathbf{Q}) = \lambda_m(\mathbf{L}) + \gamma_g$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\lambda_m(\mathbf{Q}) + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{Q})}{(\lambda_m(\mathbf{Q}) + 1)^2} \right]$$

\mathbf{Q} is reg. \mathbf{L} : $\lambda_m(\mathbf{Q}) = \lambda_m(\mathbf{L}) + \gamma_g$ and $\lambda_M(\mathbf{Q}) = \lambda_M(\mathbf{L}) + \gamma_g$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\lambda_m(\mathbf{Q}) + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{Q})}{(\lambda_m(\mathbf{Q}) + 1)^2} \right]$$

\mathbf{Q} is reg. \mathbf{L} : $\lambda_m(\mathbf{Q}) = \lambda_m(\mathbf{L}) + \gamma_g$ and $\lambda_M(\mathbf{Q}) = \lambda_M(\mathbf{L}) + \gamma_g$

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\gamma_g + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{L}) + \gamma_g}{\gamma_g^2 + 1} \right]$$

SSL with Graphs: Stability Bounds

$$\|\mathbf{f}_2^* - \mathbf{f}_1^*\|_\infty \leq \beta \leq \frac{\|\mathbf{y}_2 - \mathbf{y}_1\|_2}{\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1} + \frac{\lambda_M(\mathbf{Q})\|\mathbf{C}_1^{-1} - \mathbf{C}_2^{-1}\|_2 \cdot \|\mathbf{y}_1\|_2}{\left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_2)} + 1\right) \left(\frac{\lambda_m(\mathbf{Q})}{\lambda_M(\mathbf{C}_1)} + 1\right)}$$

Now, let us plug in the values for our problem.

Take $c_l = 1$ and $c_l > c_u$. We have $|y_i| \leq 1$ and $|f_i^*| \leq 1$.

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\lambda_m(\mathbf{Q}) + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{Q})}{(\lambda_m(\mathbf{Q}) + 1)^2} \right]$$

\mathbf{Q} is reg. \mathbf{L} : $\lambda_m(\mathbf{Q}) = \lambda_m(\mathbf{L}) + \gamma_g$ and $\lambda_M(\mathbf{Q}) = \lambda_M(\mathbf{L}) + \gamma_g$

$$\beta \leq 2 \left[\frac{\sqrt{2}}{\gamma_g + 1} + \sqrt{2n_l} \frac{1 - c_u}{c_u} \frac{\lambda_M(\mathbf{L}) + \gamma_g}{\gamma_g^2 + 1} \right]$$

This algorithm is β -stable!

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance**

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS?

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS?

What if $N \rightarrow \infty$?

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions?

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g 2) amplified commute distance

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g 2) amplified commute distance 3) \mathbf{L}^p

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g 2) amplified commute distance 3) $\mathbf{L}^p \dots$

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g 2) amplified commute distance 3) $\mathbf{L}^p \dots$

The goal of these solutions:

SSL with Graphs: Regularized Harmonic Functions

Larger implications of random walks

random walk relates to **commute distance** which should satisfy

- (*) Vertices in the **same** cluster of the graph have a **small** commute distance, whereas two vertices in **different** clusters of the graph have a **large** commute distance.

Do we have this property for HS? What if $N \rightarrow \infty$?

Luxburg/Radl/Hein: *Getting lost in space: Large sample analysis of the commute distance*

<http://www.informatik.uni-hamburg.de/ML/contents/people/luxburg/publications/>

LuxburgRadlHein2010_PaperAndSupplement.pdf

Solutions? 1) γ_g 2) amplified commute distance 3) $\mathbf{L}^p \dots$

The goal of these solutions: **make them remember!**

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives?

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $x_{n_l+n_u+1}$ arrives?

also called out-of-sample extension

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $x_{n_l+n_u+1}$ arrives?

also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $x_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive!**

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive**!

Allow to be defined everywhere: $f : \mathcal{X} \mapsto \mathbb{R}$

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive**!

Allow f to be defined everywhere: $f : \mathcal{X} \mapsto \mathbb{R}$

Allow $f(\mathbf{x}_i) \neq y_i$.

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive**!

Allow to be defined everywhere: $f : \mathcal{X} \mapsto \mathbb{R}$

Allow $f(\mathbf{x}_i) \neq y_i$. **Why?**

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive**!

Allow to be defined everywhere: $f : \mathcal{X} \mapsto \mathbb{R}$

Allow $f(\mathbf{x}_i) \neq y_i$. **Why?** To deal with noise.

SSL with Graphs: Out of sample extension

Both **MinCut** and **HFS** only inferred the labels on unlabeled data.

They are **transductive**.

What if a new point $\mathbf{x}_{n_l+n_u+1}$ arrives? also called out-of-sample extension

Option 1) Add it to the graph and recompute HFS.

Option 2) Make the algorithms **inductive**!

Allow f to be defined everywhere: $f : \mathcal{X} \mapsto \mathbb{R}$

Allow $f(\mathbf{x}_i) \neq y_i$. **Why?** To deal with noise.

Solution: **Manifold Regularization**

SSL with Graphs: Manifold Regularization

General (S)SL objective:

$$\min_f \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(f)$$

Want to control f , also for the out-of-sample data, i.e., **everywhere**.

SSL with Graphs: Manifold Regularization

General (S)SL objective:

$$\min_f \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(f)$$

Want to control f , also for the out-of-sample data, i.e., **everywhere**.

$$\Omega(f) = \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f} + \lambda_1$$

SSL with Graphs: Manifold Regularization

General (S)SL objective:

$$\min_f \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(f)$$

Want to control f , also for the out-of-sample data, i.e., **everywhere**.

$$\Omega(f) = \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f} + \lambda_1 \int_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x})^2 d\mathbf{x}$$

SSL with Graphs: Manifold Regularization

General (S)SL objective:

$$\min_f \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda \Omega(f)$$

Want to control f , also for the out-of-sample data, i.e., **everywhere**.

$$\Omega(f) = \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f} + \lambda_1 \int_{\mathbf{x} \in \mathcal{X}} f(\mathbf{x})^2 d\mathbf{x}$$

For general **kernels**:

$$\min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_I} V(\mathbf{x}_i, y_i, f(\mathbf{x}_i)) + \lambda_1 \|f\|_{\mathcal{K}}^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

SSL with Graphs: Manifold Regularization

$$f^* = \arg \min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_I} V(x_i, y_i, f) + \lambda_1 \|f\|_{\mathcal{K}}^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

SSL with Graphs: Manifold Regularization

$$f^* = \arg \min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_l} V(\mathbf{x}_i, y_i, f) + \lambda_1 \|f\|_{\mathcal{K}}^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Representer theorem for manifold regularization

The minimizer f^* has a **finite** expansion of the form

$$f^*(\mathbf{x}) = \sum_{i=1}^{n_l+n_u} \alpha_i \mathcal{K}(\mathbf{x}, \mathbf{x}_i)$$

SSL with Graphs: Manifold Regularization

$$f^* = \arg \min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_I} V(\mathbf{x}_i, y_i, f) + \lambda_1 \|f\|_{\mathcal{K}}^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Representer theorem for manifold regularization

The minimizer f^* has a **finite** expansion of the form

$$f^*(\mathbf{x}) = \sum_{i=1}^{n_I+n_u} \alpha_i \mathcal{K}(\mathbf{x}, \mathbf{x}_i)$$

$$V(\mathbf{x}, y, f) = (y - f(\mathbf{x}))^2$$

LapRLS Laplacian Regularized Least Squares

SSL with Graphs: Manifold Regularization

$$f^* = \arg \min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_l} V(\mathbf{x}_i, y_i, f) + \lambda_1 \|f\|_{\mathcal{K}}^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Representer theorem for manifold regularization

The minimizer f^* has a **finite** expansion of the form

$$f^*(\mathbf{x}) = \sum_{i=1}^{n_l+n_u} \alpha_i \mathcal{K}(\mathbf{x}, \mathbf{x}_i)$$

$$V(\mathbf{x}, y, f) = (y - f(\mathbf{x}))^2$$

LapRLS Laplacian Regularized Least Squares

$$V(\mathbf{x}, y, f) = \max(0, 1 - yf(\mathbf{x}))$$

LapSVM Laplacian Support Vector Machines

SSL with Graphs: Laplacian SVMs

$$f^* = \arg \min_{f \in \mathcal{H}_K} \sum_i^{n_I} \max(0, 1 - y f(\mathbf{x})) + \gamma_A \|f\|_K^2 + \gamma_I \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

SSL with Graphs: Laplacian SVMs




$$f^* = \arg \min_{f \in \mathcal{H}_K} \sum_i^{n_I} \max(0, 1 - y f(\mathbf{x})) + \gamma_A \|f\|_K^2 + \gamma_I \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Allows us to learn a function in **RKHS**, i.e., **RBF** kernels.


SSL with Graphs: Laplacian SVMs

$$f^* = \arg \min_{f \in \mathcal{H}_{\mathcal{K}}} \sum_i^{n_I} \max(0, 1 - y f(\mathbf{x})) + \gamma_A \|f\|_{\mathcal{K}}^2 + \gamma_I \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Allows us to learn a function in **RKHS**, i.e., **RBF** kernels.



SSL with Graphs: Laplacian SVMs



Checkpoint 1

Semi-supervised learning with graphs:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} (\infty) \sum_{i=1}^{n_l} w_{ij} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Checkpoint 1

Semi-supervised learning with graphs:

$$\min_{\mathbf{f} \in \{\pm 1\}^{n_l+n_u}} (\infty) \sum_{i=1}^{n_l} w_{ij} (f(\mathbf{x}_i) - y_i)^2 + \lambda \sum_{i,j=1}^{n_l+n_u} (f(\mathbf{x}_i) - f(\mathbf{x}_j))^2$$

Regularized harmonic Solution:

$$\mathbf{f}_u = (\mathbf{L}_{uu} + \gamma_g \mathbf{I})^{-1} (\mathbf{W}_{ul} \mathbf{f}_l)$$

Checkpoint 2

Unconstrained regularization in general:

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q}\mathbf{f}$$

Checkpoint 2

Unconstrained regularization in general:

$$\mathbf{f}^* = \min_{\mathbf{f} \in \mathbb{R}^N} (\mathbf{f} - \mathbf{y})^\top \mathbf{C}(\mathbf{f} - \mathbf{y}) + \mathbf{f}^\top \mathbf{Q} \mathbf{f}$$

Out of sample extension: Laplacian SVMs

$$f^* = \arg \min_{f \in \mathcal{H}_K} \sum_i^{n_I} \max(0, 1 - y f(\mathbf{x})) + \lambda_1 \|f\|_K^2 + \lambda_2 \mathbf{f}^\top \mathbf{L} \mathbf{f}$$

Daniele Calandriello

dcalandriello@google.com

ENS Paris-Saclay, MVA 2022/2023

<https://sites.google.com/view/daniele-calandriello/>