

## Exercises 4

$i$ : school

$j$ : student

$\theta$ : school level means

$$y_{ij} \sim \mathcal{N}(\theta_i, \sigma^2)$$

$$\begin{aligned} \text{(A)} \quad \mathcal{L}(\theta_i) &= \prod_{j=1}^n f(y_{ij} | \theta_i, \sigma^2) \\ &= \prod_{j=1}^n \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{1}{2\sigma^2}(y_{ij} - \theta_i)^2\right) \\ &= \left(\frac{1}{\sigma\sqrt{2\pi}}\right)^n \prod_{j=1}^n \exp\left(-\frac{1}{2\sigma^2}(y_{ij} - \theta_i)^2\right) \\ &= \left(\frac{1}{\sigma\sqrt{2\pi}}\right)^n \exp\left(-\frac{1}{2\sigma^2} \sum_{j=1}^n (y_{ij} - \theta_i)^2\right) \\ &= \left(\frac{1}{\sigma\sqrt{2\pi}}\right)^n \exp\left[-\frac{1}{2\sigma^2} (\bar{y}^2 n - 2\theta_i \bar{y} n + n\theta_i^2)\right] \end{aligned}$$

$$\begin{aligned} \frac{\partial \log \mathcal{L}(\theta_i)}{\partial \theta_i} &= 0 \\ \frac{\partial}{\partial \theta_i} \left[ -\frac{1}{2\sigma^2} (\bar{y}^2 n - 2\theta_i \bar{y} n + n\theta_i^2) \right] &= 0 \\ -2\bar{y} n + 2\theta_i n &= 0 \\ \theta_i &= \bar{y} \end{aligned}$$

For all schools, we have

$$\theta_{MLE} = (\bar{y}_1, \bar{y}_2, \dots, \bar{y}_P)$$

(b) This is because fewer samples will result in a poorer estimate of the true mean of the school

(c)

Fit model using Gibbs sampling:

$$y_{ij} \sim \mathcal{N}(\theta_i, \sigma^2) \quad | \rightarrow P(y_{ij} | -)$$

$$\begin{aligned}\theta_i &\sim \mathcal{N}(\mu, \tau^2 \sigma^2) \\ \mu &\sim \text{Flat}(\mu) \\ \sigma^2 &\sim \text{Jeffrey's prior} \\ \tau^2 &\sim \text{IG}(1/2, 1/2)\end{aligned}$$

$$\begin{aligned}P(\theta_i | -) \\ P(\mu | -) \\ P(\sigma^2 | -) \\ P(\tau^2 | -)\end{aligned}$$

$$\text{Note } \bar{y}_i \sim \mathcal{N}(\theta_i, \frac{\sigma^2}{n})$$

$$* P(\theta_i | \bar{y}_i, \mu, \sigma^2, \tau^2)$$

$$\propto P(\bar{y}_i | -) P(-)$$

$$\propto P(\bar{y}_i | -) P(\theta_i | \tau^2, \sigma^2, \mu) \underbrace{P(\tau^2) P(\sigma^2) P(\mu)}_{\tau^2 \perp \sigma^2 \perp \mu}$$

$$\propto P(\bar{y}_i | -) P(\theta_i | \tau^2, \sigma^2, \mu)$$

$$\propto \exp\left(-\frac{1}{2\sigma^2/n} (\bar{y}_i - \theta_i)^2\right) \times \exp\left(-\frac{1}{2\sigma^2\tau^2} (\theta_i - \mu)^2\right)$$

$$\propto \exp\left(-\frac{1}{2\sigma^2/n} (\bar{y}_i^2 - 2\bar{y}_i\theta_i + \theta_i^2) - \frac{1}{2\sigma^2\tau^2} (\theta_i^2 - 2\theta_i\mu + \mu^2)\right)$$

$$\propto \exp\left(-\frac{1}{2\sigma^2\tau^2/n} (-2\bar{y}_i\theta_i\tau^2 + \theta_i^2\tau^2 + \theta_i^2 n^{-1} - 2\theta_i\mu n^{-1})\right)$$

$$\propto \exp\left(-\frac{n}{2\sigma^2\tau^2} (-2\theta_i(\bar{y}_i\tau^2 + \mu n^{-1}) + \theta_i^2(\tau^2 + n^{-1}))\right)$$

$$\propto \exp\left(-\frac{n}{2\sigma^2\tau^2} \left(\theta_i^2 - 2\theta_i \left(\frac{\bar{y}_i\tau^2 + \mu n^{-1}}{\tau^2 + n^{-1}}\right)\right) (\tau^2 + n^{-1})\right)$$

$$\propto \exp\left(-\frac{n(\tau^2 + n^{-1})}{2\sigma^2\tau^2} \left(\theta_i - \left(\frac{\bar{y}_i\tau^2 + \mu n^{-1}}{\tau^2 + n^{-1}}\right)^2\right)\right)$$

$$\text{This is a } \mathcal{N}\left(\frac{\bar{y}_i\tau^2 + \mu n^{-1}}{\tau^2 + n^{-1}}, \frac{\sigma^2\tau^2}{\tau^2 + n^{-1}}\right)$$

$$\left[ \frac{1}{\tau^2} + \frac{1}{\sigma^2} \right]$$


$$\propto \frac{\bar{y} \tau^2 n_i + N}{\tau^2 n_i + 1}$$

$$* P(N|-)$$

$$\propto P(N|\theta_i)$$

$$\propto P(\theta_i|N, \sigma^2, \tau^2) P(N)$$

$$\propto \prod_{i=1}^P \exp\left(-\frac{1}{2\tau^2\sigma^2} (\theta_i - N)^2\right)$$



$$\propto \exp\left(-\frac{1}{2\tau^2\sigma^2} \sum_P (\theta_i - N)^2\right)$$

$$\propto \exp\left(-\frac{1}{2\tau^2\sigma^2} \sum_P (\theta_i^2 - 2\theta_i N + N^2)\right)$$

$$\propto \exp\left(-\frac{1}{2\tau^2\sigma^2} (P\bar{\theta}^2 - 2\bar{\theta}NP + PN^2)\right)$$

$$\propto \exp\left(-\frac{P}{2\tau^2\sigma^2} (-2\bar{\theta}N + N^2)\right)$$

$$\propto \exp\left(-\frac{P}{2\tau\sigma^2} (\bar{\theta} - N)^2\right)$$

This is a normal  $N(\bar{\theta}, \frac{\sigma^2\tau^2}{P})$

\*

$$P(\tau^2|-) \propto P(\tau^2|\theta) \propto P(\theta|\tau^2) P(\tau^2)$$

$$\propto (\tau\sigma)^{-P} \exp\left(-\frac{1}{2}(\theta - N)^T (\tau^2\sigma^2\mathbf{I})^{-1} (\theta - N)\right) (\tau^2)^{-\frac{1}{2}-1} \exp\left(-\frac{1}{2}\tau^{-2}\right)$$

$$\propto (\tau^2)^{-\frac{P}{2}-\frac{1}{2}-1} \exp\left(-\frac{1}{2}(\theta - N)^T (\tau^2\sigma^2\mathbf{I})^{-1} (\theta - N) + \tau^{-2}\right)$$

$$\propto (\tau^2)^{-\frac{P+1}{2}-1} \exp\left(-\frac{1}{2}\tau^{-2} \left[(\theta - N)^T (\sigma^2\mathbf{I})^{-1} (\theta - N) + 1\right]\right)$$

This is an  $IG\left(\frac{P+1}{2}, \frac{(\theta - N)^T (\sigma^2\mathbf{I})^{-1} (\theta - N) + 1}{2}\right)$

$$* P(\sigma^2|-) \propto P(\sigma^2|\theta_i, y_i)$$

$$\propto P(\sigma^2|\theta, y) \propto P(y|\theta) P(\theta) P(\sigma^2)$$

First find  $P(y|\theta)$

$$\propto \prod_{i=1}^p \prod_{j=1}^{n_i} \sigma^{-1} \exp\left(-\frac{1}{2\sigma^2} (\theta_i - y_{ij})^2\right)$$

$$\propto \sigma^{-N} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^p \sum_{j=1}^{n_i} (\theta_i - y_{ij})^2\right)$$

$$\text{so } p(\sigma^2 | -) \propto (\sigma^2)^{-\frac{N}{2}} \exp\left(-\frac{1}{2\sigma^2} \sum_{i=1}^p \sum_{j=1}^{n_i} (\theta_i - y_{ij})^2\right)$$

$$(\sigma^2 \tau^2)^{-\frac{p}{2}} \exp\left(-\frac{1}{2} (\theta - N)^T (\sigma^2 \tau^2 I)^{-1} (\theta - N)\right) \times \frac{1}{\sigma^2}$$

$$\propto (\sigma^2)^{-\frac{N+p}{2}-1} \exp\left(-\frac{1}{2\sigma^2} ((\theta - N)^T (\tau^2 I)^{-1} (\theta - N) + \sum_{i=1}^p \sum_{j=1}^{n_i} (\theta_i - y_{ij})^2)\right)$$

This is an IG  $\left( \frac{N+p}{2}, \frac{(\theta - N)^T (\tau^2 I)^{-1} (\theta - N)}{2} + \frac{\sum_{i=1}^p \sum_{j=1}^{n_i} (\theta_i - y_{ij})^2}{2} \right)$

$$D) E(\theta_i | -) = \frac{\bar{y}_i \tau^2 + N n_i^{-1}}{\tau^2 + n_i^{-1}}$$

$$= \frac{\bar{y}_i \tau^2 n_i + N}{\tau^2 n_i + 1}$$

$$= \underbrace{\frac{1}{\tau^2 n_i + 1}}_{k_i} N + \underbrace{\frac{\tau^2 n_i}{\tau^2 n_i + 1}}_{1 - k_i} \bar{y}_i$$

$$k_i = \frac{1}{\tau^2 n_i + 1}$$

$$(E) \quad y_{ij} = \mu + \delta_i + \epsilon_{ij}$$

$$\delta_i \sim \mathcal{N}(0, \tau^2 \sigma^2)$$

$$\epsilon_{ij} \sim \mathcal{N}(0, \sigma^2)$$

$$y_{ij} \sim \mathcal{N}(\theta_i, \sigma^2)$$

$$\theta_i \sim \mathcal{N}(\mu, \tau^2, \sigma^2)$$

$$\theta_i = \mu + \delta_i$$

$$\delta_i \sim \mathcal{N}(0, \tau^2 \sigma^2)$$

$$\epsilon_{ij} \sim \mathcal{N}(0, \sigma^2)$$

$$y_{ij} = \mu + \delta_i + \epsilon_{ij}$$

$$y_{ij} = \mu + \delta_i + \epsilon_{ij}$$

$$\text{cov}(y_{ij}, y_{ik}) \quad j \neq k$$

$i \equiv \text{school}$   
 $j \equiv \text{student}$

between  
students

$$E[(\mu + \delta_i + \epsilon_{ij} - E(y_{ij}))$$

$$(\mu + \delta_i + \epsilon_{ik} - E(y_{ik}))]$$

$$= E[(\cancel{\mu} + \delta_i + \epsilon_{ij} - \cancel{\mu})(\cancel{\mu} + \delta_i + \epsilon_{ik} - \cancel{\mu})]$$

$$= E[(\delta_i + \epsilon_{ij})(\delta_i + \epsilon_{ik})]$$

$$= E[\delta_i^2 + \delta_i \epsilon_{ik} + \epsilon_{ij} \delta_i + \epsilon_{ij} \epsilon_{ik}]$$

$$= \text{var}(\delta_i) + \cancel{\text{cov}(\delta_i, \epsilon_{ik})} + \cancel{\text{cov}(\delta_i, \epsilon_{ij})} + \cancel{\text{cov}(\epsilon_{ij}, \epsilon_{ik})}$$

$$= \boxed{\sigma^2}$$

$$\text{cov}(y_{ij}, y_{i'k})$$

different  
schools

$$= E[(\cancel{\mu} + \delta_i + \epsilon_{ij} - \cancel{\mu})(\cancel{\mu} + \delta_{i'} + \epsilon_{i'k} - \cancel{\mu})]$$

$$= E[\delta_i \delta_{i'} + \delta_i \epsilon_{i'k} + \epsilon_{ij} \delta_{i'} + \epsilon_{ij} \epsilon_{i'k}]$$

$$= \boxed{0}$$

(F) No?

## Blood pressure

A) t-test

$$H_0: \mu_2 - \mu_1 = 0$$

$$H_1: \mu_2 - \mu_1 \neq 0$$

similar  
variances

if  $H_0$  is true  $\mu_2 - \mu_1 \sim t$  with

$$\mu_0 = 0, \quad s = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}}$$

$$df = n_1 + n_2 - 2$$

Reject  $H_0$  with  $\alpha = 0.05$

Not appropriate because there are several observations per person & people sampled different number of times.

B) Still reject  $H_0$ . Standard error is bigger because we have less



samples. it is still problematic that all subjects don't have the same number of tests.

(c)  $y_{ij} \sim N(\theta_i, \sigma^2)$  \* wrong, probably  
 $\theta_i \sim N(\mu + \beta x_i, \tau^2 \sigma^2)$

$j$  = measurement number

$i$  = person

flat prior on  $\beta \rightarrow$  doesn't change anything, replace  $N$  with

$P(\beta | \dots) \propto P(\beta | \theta_i)$   
 $\propto P(\theta_i | \beta, \dots) P(\beta)$   $\mu + \beta x_i$   
in conditionals

$\propto \prod_i^P \exp\left(-\frac{1}{2\sigma^2\tau^2}(\theta_i - (\mu + \beta x_i))^2\right)$

$\propto \exp\left(-\frac{1}{2\sigma^2\tau^2} \sum_i^P (\theta_i - (\mu + \beta x_i))^2\right)$

$\propto \exp\left(-\frac{1}{2\sigma^2\tau^2} \sum_i^P (\theta_i^2 - 2\theta_i(\mu + \beta x_i) + (\mu + \beta x_i)^2)\right)$

$\propto \exp\left(-\frac{1}{2\sigma^2\tau^2} \left(\cancel{\beta^2 \sum_i^P x_i^2} - 2 \sum_i^P \theta_i \beta x_i + \sum_i^P \mu^2 + 2\mu \sum_i^P \beta x_i + \beta^2 \sum_i^P x_i^2\right)\right)$



$$\propto \exp \left( \frac{-1}{2\sigma^2\tau^2} \left( -2\overline{\theta_i x_i} p \beta + 2N\beta p \bar{x}_i + \beta^2 \overline{x_i^2} p \right) \right)$$

$$\propto \exp \left( \frac{-1}{2\sigma^2\tau^2} \left( -2\beta (\overline{\theta_i x_i} p - Np\bar{x}_i) + \beta^2 \overline{x_i^2} p \right) \right)$$

$$\propto \exp \left( \frac{-\overline{x_i^2} p}{2\sigma^2\tau^2} \left( -2\beta \left( \frac{\overline{\theta_i x_i} p - Np\bar{x}_i}{\overline{x_i^2} p} \right) + \beta^2 \right) \right)$$

$$\propto \exp \left( -\frac{\overline{x_i^2} p}{2\sigma^2\tau^2} \left( \beta - \left( \frac{\overline{\theta_i x_i} p - Np\bar{x}_i}{p \overline{x_i^2}} \right)^2 \right) \right)$$

This is a  $\mathcal{N} \left( \frac{\overline{\theta x} - N \bar{x}}{\overline{x^2}}, \frac{\sigma^2\tau^2}{p \overline{x^2}} \right)$

$$P(N| \rightarrow) \propto P(N|\theta) \quad \downarrow \quad \frac{0.5(\theta_2 - N)}{0.5} \quad \downarrow \quad \frac{\sigma^2\tau^2}{1}$$

$$\propto P(\theta|N) P(N)$$

$$\propto \prod_p \exp \left( -\frac{1}{2\tau^2\sigma^2} (\theta_i - (N + \beta x_i))^2 \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} \sum (\theta_i - (N + \beta x_i))^2 \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} \sum (\theta_i^2 - 2(N + \beta x_i)\theta_i + (N + \beta x_i)^2) \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} \sum (\theta_i^2 - 2(N + \beta x_i)\theta_i + N^2 + 2\beta x_i N + \beta^2 x_i^2) \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} \sum (-2(N\theta_i - \beta x_i N) + N^2) \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} (-2(N\bar{\theta} - \beta \bar{x}N) + N^2) \right)$$

$$\propto \exp \left( -\frac{1}{2\tau^2\sigma^2} (-2N(\bar{\theta} - \beta \bar{x}) + N^2) \right)$$

$$\propto \exp \left( -\frac{P}{2\tau^2\sigma^2} (N - (\bar{\theta} - \beta \bar{x}))^2 \right)$$

$$\mathcal{N}(\bar{\theta} - \beta \bar{x}, \frac{\tau^2\sigma^2}{P})$$

$$P(\theta_i | -) \propto P(\theta_i | \bar{y}_i)$$

$$\propto P(\bar{y}_i | \theta_i) P(\theta_i)$$

$$\propto \exp \left( -\frac{n}{2\sigma^2} (\bar{y}_i - \theta_i)^2 \right)$$

$$\times \exp \left( -\frac{1}{2\sigma^2\tau^2} (\theta_i - (N + \beta x_i))^2 \right)$$

$$2 \exp \left( -\frac{n}{2\sigma^2} (\bar{y}_i^2 - 2\bar{y}_i\theta + \theta^2) \right. \\ \left. - \frac{1}{2\sigma^2\tau^2} (\theta^2 - 2\theta(n + \beta x_i) + (n + \beta x_i)^2) \right)$$

$$2 \exp \left( -\frac{n}{2\sigma^2\tau^2} (-2\bar{y}_i\theta + \theta^2\tau^2 + \theta^2 n_i^{-1} - 2\theta(n + \beta x_i)) \right)$$

$$2 \exp \left( \frac{-n}{2\sigma^2\tau^2} (-2\theta(\bar{y}_i\tau^2 + n_i^{-1}(n + \beta x_i)) + \theta^2(\tau^2 + n_i^{-1})) \right)$$

$$2 \exp \left( -\frac{n_i(\tau^2 + n_i^{-1})}{2\sigma^2\tau^2} \left( \theta - \frac{\bar{y}_i\tau^2}{\tau^2 + n_i^{-1}} + n_i^{-1}(n + \beta x_i) \right) \right)$$

$$\mathcal{N} \left( \frac{\bar{y}_i\tau^2 n_i + (n + \beta x_i)}{n_i\tau^2 + 1}, \frac{\sigma^2\tau^2}{\tau^2 n_i + 1} \right)$$

$$n_1 = n_2 + \beta$$

Q.4

$$\beta = \Delta n$$