Journal of Water and Health

Emerging pathogens and deliberate attacks on water supplies: A scenario planning workshop --Manuscript Draft--

Manuscript Number:	JWH-D-18-00248			
Full Title:	Emerging pathogens and deliberate attacks on water supplies: A scenario planning workshop			
Article Type:	Research Paper			
Corresponding Author:	Julii Brainard University of East Anglia Norwich Medical School UNITED KINGDOM			
Corresponding Author Secondary Information:				
Corresponding Author's Institution:	University of East Anglia Norwich Medical School			
Corresponding Author's Secondary Institution:				
First Author:	Julii Brainard			
First Author Secondary Information:				
Order of Authors:	Julii Brainard			
	Charlotte Christiane Hammer, MA, MSc			
	Maha Bouzid, PhD			
	Paul Raymond Hunter, PhD			
Order of Authors Secondary Information:				
Abstract:	Microbiological contamination of drinking water supplies is an ever-present concern for water utility managers. Most such threats are routine, well-recognised and described. Therefore, they can usually be prevented using standard protection measures. Incidents involving emerging pathogens and malicious attacks are inherently less predictable. In a multi-stage process over one day, participants with backgrounds in microbiology, medicine, infrastructure, data analysis, environmental or public health and facility management developed qualitative scenarios on potential threats posed by either an emergent pathogen in or a microbiological attack on drinking water supplies in a European country. Participants were guided via structured activities to identify key factors that would impact the magnitude and severity of such an emergency. Plausible variant states for each key factor were determined, and participants constructed sequences of events to create scenario outlines. Five scenarios in outline form are reported which incorporate genuine future events as well as pathogens of international concern. Common features that would exacerbate all scenarios were: underinvestment in public services, inadequate water quality testing and monitoring and lack of resources to keep water supplies safe. Participant evaluation of their scenario planning experience was broadly very positive and the scenario planning process was received as credible and relevant.			



Doctor Julii Brainard Norwich Medical School Norwich Research Park NR4 7TJ United Kingdom

Tel. 01603 591151 www.med.uea.ac.uk email j.brainard@uea.ac.uk

3 September 2018

Dear editors of the Journal of Water and Health,

Thank you for your invitation to resubmit our original Research Paper with title

Emerging pathogens and deliberate attacks on water supplies: A scenario planning workshop

We appreciate very much the comments from two previous reviewers on our article. We have taken into account all of their comments, which are addressed below.

This article is still an excellent fit for *JWH*. These scenario planning efforts addressed important security issues that attract significant concern. Our project integrated a multiplicity of perspectives from microbiology, medicine, risk-analysis, facility management, political science, public health and other areas of expertise to identify key aspects of risk with regard to unusual contamination events. The process was innately creative and encouraged minority viewpoints to see how these unusual situations can arise. The manuscript serves both to show how to undertake a similar workshop and provides evaluation of how well such an approach can meet the needs of people involved with protecting health and water safety.

We appreciate your consideration of our research and hope that you will find it suitable for publication.

We supply the names and addresses of three additional persons who can supply informed independent review if required, as follows:

Mr. Matthew Heberger, matth@sfei.org Thomas.Kistemann@ukb.uni-bonn.de

Juli S Brainard

Faithfully yours,

Dr. Julii Brainard (corresponding author, J.Brainard@uea.ac.uk) on behalf of all authors

RESPONSES TO REVIEWERS. Reviewer comments in purple, Author replies in black

Reviewer #1: The title does not clearly reflect that this is not an assessment of threats, nor is it an evaluation of response actions.

The last part of title was changed to 'A scenario planning workshop' to more clearly convey that this was a workshop event.

Rev#1 Overall the paper is an interesting recounting of the workshop,

Thank you for this comment

Rev#1 I found it difficult to think about replicating the process or learning lessons for other water systems. Seems more like a conference presentation than a research paper.

We hope that this revised manuscript seems improved and more valuable. The workshops steps were difficult to explain clearly. We added a figure (1) that may help. The Discussion section now explains more clearly what the benefits can be to participants and the organisations they work for. We hope that if nothing else, the article demonstrates what kind of outputs can be produced out of a one-day workshop with many experts. Readers will be able to judge for themselves if commitment to a similar workshop would be valuable to their organisation. We describe (lines 314-317) how the intermediate products (key factors for instance) may be useful for risk management in themselves. There are lessons to be learnt indirectly with regard to what factors these participants felt were most relevant to the possible impact of an attack or emerging disease in their water source. These exact points have been added to our manuscript (first paragraph of Discussion section)

Reviewer #2: 18-171

This article is worth being published, once added some lacking pieces of information.

Thank you for this comment

Rev#2 More about attendance How many were the participants to the meeting?

Reviewer is very right, we should have included this information. The numbers (31 participants) were added at the start of Results section, line 209.

What profession were represented (said in the summary, but not in the text body)?

Reviewer is very right, we should have included this information in the main text. The professional backgrounds and their types of employers as well as countries where they worked were added at the start of Results section, lines 210-215

It would be worth distinguishing the different professional skills involved and to highlight possible difference among the points of view concerning the factors participating to the impact.

There was only one notable set of contributions that could be linked to professional background; one of our microbiologists struggled with the emerging pathogen scenario. She argued that progress in genomic analysis would soon make it unlikely that an emerging pathogen would pose a huge threat simply by its metabolism or effects on human health being poorly understood. We allude to that perspective in the revised manuscript (lines 241-246) but also caveat that argument by saying that rapid and reliable detection relies on good expertise, regular and consistent water quality tests, and well-resourced laboratory testing (all of which it may be foolish to take for granted).

As well, all participants emphasized the lack of resources (staff, money) but did not detail where funding would be more useful. Did the participant agree on the priorities?

Unfortunately, we can't report on this in the detail that Reviewer would like because we didn't explore funding priorities or competing demands in enough detail. The subtext was probably that those working in the water industry (many from southern Europe which was hard-hit in the recent economic crisis) were keenly aware of the atmosphere of 'Austerity' in public service provision, and that they didn't want those economic pressures to undermine water safety. We shouldn't try to infer all of that formally, though.

Outbreak detection

Among the different steps of the investigation, the first one from the earliest signal occurrence to the gathering of the crisis committee may take a very variable time according to the former experience and training of the receptor.

We identified from our experience in outbreak investigation that the early collation of both health information (epidemiological approach) and water related information (weather and water operation data) is crucial to detect a water related outbreak, speed up the investigations, guide lab strategy to identify the involved pathogens and, above all, orientate decision making for people protection (boiling advice, cut of water distribution...). The interpretation of signals often differ among different professions, thus possibly creating confusion in communication and lowering the concurrent mobilization of actors.

All participants to the Aquavalens meeting underlined the importance of forming a crisis committee to avoid cacophony in communication, but the prior early meeting of water operational staff and local health authorities is not addressed.

Collaboration and efficient communication between water and health staff is certainly important as stated by the reviewer and evidenced by the key factors and variants identified by the workshop participants and presented in Table 2 & 3. Specifically, Emergency response (for ED) and Interagency cooperation (for AtK) encompass this aspect. However, the workshop participants did not emphasise a particular focus on early meeting/ preparedness plan involving simultaneously water utilities and health authorities, which perhaps (unfortunately) mirrors the situation in the real world. The manuscript is a report of the workshop process and findings and as stated previously, another set of attendants could take a different direction and perspective to scenario formulation and response. Forming a crisis committee was not what our participants chose as a key aspect of the scenario stories. We're reluctant to start listing all the factors and developments our participants didn't choose (would be a different paper), but we now reinforce (line 304) that the scenarios chosen by our participants are far from definitive.

Also, the possible shift of the responsibility for steering investigation from civil to military staff in the case of terrorist attack may also create problems.

Again, this is a valid point, but one that depends highly on context; perhaps it didn't seem applicable in the countries of origin that our participants had. As we said on lines 304-309, a different group of water experts would have chosen different key factors and imagined different official responses in each scenario.

Minor remarks:

Introduction: The exclusion of chemical weapons is questionable and should be briefly justified. Indeed, stakeholders involved in the early management and investigation stages are quite the same.

We completely agree, but we could only do so much in one day. The first paragraph in the Discussion section (lines 298-317) now mentions that the methods are very suitable to be adapted to address other risks, especially chemical or radiological.

Since the article will be read by different professional who are not daily used to organize joined investigations (water operators, public health institutes, microbiologists...) a bit more basic pedagogy should be used to improve mutual understanding of event.

We were a bit unsure what reviewer meant, but hope these changes will help. There is text (lines 107, 171-173) about the value of silent methods to make sure that all voices are heard, the choice of participants to also ensure a broad range of backgrounds. We added a figure (Figure 1, flowchart style) that explains the scenario building steps in phases (page 6).

Figure 1 is not very useful and should be removed.

We swapped the previous figure 1 for a different image (now Figure 2) that more clearly shows what is written on the post-its and how they are items that could be sensibly grouped together. This figure shows what we were trying to describe (results of silent sorting). Reviewer 1 said they weren't sure how the workshop was implemented and removing this figure will remove clear detail of what we did. The caption was changed slightly (line 190)

Appendix is also partly redundant with Table 4 and may be suppressed.

The current movement is towards more transparency about data collection methods; and seeing genuine evaluation forms can be useful to others (was useful to us when trying to design our own). We would like to keep this in the appendix.

L 196: Could you precise what kind of "policy changes" you mean? All policy changes do not disrupt the management of the crisis.

That is a very valid observation. A paragraph was added to the discussion to address (briefly) types of policies and their impacts (lines 244-248, 334-341)

Emerging pathogens and deliberate attacks on water
 supplies: A scenario planning workshop

- Julii Brainard*, Charlotte C. Hammer, Maha Bouzid, Paul R. Hunter and
- 6 the Aquavalens Consortium
- 8 Norwich Medical School, University of East Anglia
- 9 Norwich NR4 7TJ, Norfolk, United Kingdom
- *corresponding author: j.brainard@uea.ac.uk
- Short Title = Scenario planning for pathogens in and attacks on water

Abstract

Microbiological contamination of drinking water supplies is an ever-present concern for water utility managers. Most such threats are routine, well-recognised and described. Therefore, they can usually be prevented using standard protection measures. Incidents involving emerging pathogens and malicious attacks are inherently less predictable. In a multi-stage process over one day, participants with backgrounds in microbiology, medicine, infrastructure, data analysis, environmental or public health and facility management developed qualitative scenarios on potential threats posed by either an emergent pathogen in or a microbiological attack on drinking water supplies in a European country. Participants were guided via structured activities to identify key factors that would impact the magnitude and severity of such an emergency. Plausible variant states for each key factor were determined, and participants constructed sequences of events to create scenario outlines. Five scenarios in outline form are reported which incorporate genuine future events as well as pathogens of international concern. Common features that would exacerbate all scenarios were: underinvestment in public services, inadequate water quality testing and monitoring and lack of resources to keep water supplies safe. Participant evaluation of their scenario planning experience was broadly very positive and the scenario planning process was received as credible and relevant.

Keywords: scenario planning, water safety, emerging diseases, contamination, health protection

Introduction and aims

The prospect of microbiological contamination of drinking water supplies is an ever-present concern for managers of public water supplies. Most such threats are routine, well-recognised and described, and thus can usually be prevented using standard protection measures. Incidents involving emerging pathogens and attacks, however, are inherently less predictable. Either because by definition emerging pathogens are not well understood, or because of deliberate planning by attackers to hide their actions and maximise surprise. Vulnerability of municipal water supplies to such events is a topic of ongoing concern. (Haimes *et al.* 1998, Clark & Deininger 2000, Qiao *et al.* 2007, Gleick & Heberger 2014, Mutchek & Williams 2014). Documented plots to deliberately contaminate drinking water supplies in the period 1946-2015 were described by Brainard and Hunter (2016). Literature about emerging diseases in drinking water is more expansive and diverse and tends to be segregated by species, groups of similar microbes or parasites, outbreak context or types of control methods. For instance, a worldwide review of protozoa outbreaks is in Baldursson and Karanis (2011). Sinclair *et al.* (2009) undertook a structured literature review to search for and report on virus-disease outbreaks in recreational waters.

Emerging diseases (ED) refer to poorly understood, often relatively recently discovered pathogens for which our understanding of the risks posed to public health through environmental routes of transmission are not well understood. Concerns about ED and potential mitigation measures overlap greatly with concerns about and possible strategies to resist deliberate contamination of water supplies (such as for terrorist reasons). This report documents the methods and outcomes from a meeting designed to help prepare for both possible threats.

Scenarios have been described as "Stories that can help us recognise and adapt to changing aspects of our present environment. They form a method for articulating the different pathways that might exist and identify plausible steps to move down each of those possible paths" (Schwartz 1996). Scenario planning can have a positive impact on decision quality in response to uncertain and rapidly developing situations (Meissner & Wulf 2013). Scenario development is the first step towards identifying strategies that result in robust decision making in multi-faceted situations with high uncertainty about possible risks (Kwakkel *et al.* 2016). As part of the exercise, our participants learned many concepts typically involved in structured approaches to scenario construction. Drawing on methods used in other published scenario building events, our approach was pragmatic and adapted

so as to be completed in one day and potentially easily replicated in participants' own organisations and workplaces, if desired.

Methods

European professionals with expertise in water safety, water provision, water-borne diseases and envirionmental protection were invited to attend a workshop that was designed to create an active participatory meeting. The workshop format was a short plenary session followed by smaller group collaborative work (in parallel), with a final plenary session at the end of the afternoon. All participants were allocated to either the Emerging Diseases (ED) or Attack (AtK) groups by event organisers. A subject expert was purposefully placed in each group (acting as both participant and topic expert), in case of technical queries that the facilitators could not answer. Our sessions were normative-participatory in nature, which means they consolidated multiple theories and concepts for participant engagement (Steinmüller 1997).

Theoretical foundation

Scenarios are "hypothetical sequence[s] of events constructed for the purpose of focusing attention on causal processes and decision points" (Kahn & Wiener 1967). Thus, scenarios are hypothetical but still outlined and concrete (Wilson 1978). Scenario planning is suited to developing multiple alternatives of possible futures taking into consideration unlikely futures with unknown probability of actually occurring. These are relatively extreme events, so-called very rarely expected 'Black Swans'. Alternative scenarios can be used to explore possible tipping points and thus interventions that might disrupt multiple negative scenarios. Because the scenarios we sought to develop were anticipatory in nature, we used a participatory-normative approach to scenario planning. There is little to no formalisation in this type of scenario planning unlike in forecasting and trend exploration (Steinmüller 1997).

Given the short time-frame, our scenarios were brief outlines, rather than fully developed and written versions. They were semi-global, taking into consideration wider contextual issues but remaining close to the original topic. The scenarios were fully qualitative. We used an intentional interpretation of the scenario, which assumes that the scenario is not the text developed by the participants but rather the blueprint of the future developed by them.

Core concepts

- Key factors and key factor analysis: Key factors are those factors that define the outcome of an event. They need to be highly important (i.e. must have strong impact) and uncertain (i.e. not definite, such as the force of gravity on Earth).
- Silent sorting: applicable at stage of nomination and grouping of possible key factors. The process is done in silence to encourage multiplicity of perspectives.
- Key factor variants: A variant of a key factor is a variable status that the factor could take (i.e. if a key factor is climate, variants could be hot and dry, cold and wet, cold and dry, etc.)
- Backcasting: Backcasting is the process to link the (future) scenario with the present. Backcasting starts at the point of the scenario (in our case 2023) and describes events in reverse order, going backwards from future to the present. It is the opposite of forecasting and seeks to develop pivotal points that lead to a scenario becoming reality or not.

Implementation

Prior to the workshop, fixed conditions for each scenario were decided by the facilitators. These fixed conditions are parameters that cannot be changed during the scenario planning process, and serve to orient the outputs towards prespecified objectives. Fixed scenario conditions are listed in Table 1. Figure 1 summarises the steps that participants went through during the workshop.

Table 1 | Fixed conditions for building each scenario

Both Scenarios			
Included conditions	Year = 2023, In a water supply that participants are responsible		
	for. Within European area. Pathogen is unknown initially.		
Excluded possible factors	Any problem more important than whether the water is safe to		
	drink, such as concurrent nuclear war, plague-like disease, zombie		
	apocalypse, etc. At discretion of facilitator.		
	Respective scenario fixed conditions		
Emerging disease	Presence in water is not result of deliberate contamination		
	Has been linked to some deaths.		
Attack	Deliberate attack confirmed by evidence of break-in, intelligence		
	chatter, and linked hospital admissions.		

Session

Plenary

and Selection

Factor Identification

Key

Variant selection

Backcasting to write scenario

Figure 1 | Scenario Planning Workshop Methods

Objectives explained to participants Overview of methods and fixed conditions explained Division of participants into two groups: Attack and Emerging Disease

Separately in each of Attack and Emerging Disease groups:

Propose key factors = Important but uncertain aspects of scenario

Sort the key factors into groups (silently)

Suggest links between KF groups (silently)

(Silently) Suggest summary key factors (labelling groups of KF)

Vote for the 4-8 most important key factors

Within each of Attack/ED groups, divide into 4-6 smaller groups to address variant states for each key factor

Each subgroup: Decide on four possible variant states for the 4-8 most important key factors

Each participant: Selects one variant from each KF factor to go into three scenarios

Divide participants still within Attack/ED themes into three subgroups

Each scenario subgroup: Write a scenario by backcasting from future (ie., year 2023) when all variants are true, to the present year and conditions.

Incorporate feasible ad hoc details to explain how the future variant statuses arose, going back in time

Rejoin Plenary to share scenarios with others

Identification and prioritisation of key factors

For the workshop, we devised an approach that allowed scenarios to be devised quickly (work progressed from factor identification to simple scenario descriptions in one full working day).

Participants were randomly allocated to one of the two scenario groups (attack and emerging pathogen) with some adjustment made to ensure relatively equal distribution of skills and representatives from the same organisation(s). Within each group, participants were asked to identify all relevant key factors that are likely to influence the scenario they are working on. These were written on adhesive notes and stuck to a wall or whiteboard. Duplicate entries were removed and the participants were asked to check if they could think of any further key factors. Once all key factors were identified and agreed, participants were asked to look for relationships and interactions between these factors, with the option of creating groups, over-arching category labels, and relationship indicators such as arrows (Figure 2). Participants were allowed to nominate, remove, cluster or remove from clusters both their own key factors and those contributed by other participants. This was done in silence (silent sorting). Any arguments about the validity of a factor or its associations had to be resolved silently by moving or removing the respective factor. Silence deterred habitual deference to verbally or personality dominant colleagues and meant that minority voices and alternative perspectives were more likely to be heard.

The previous phase allowed the development of a complex network of interrelated key factors. The next stage was to identify the most important factors for each scenario. In the voting stage, each participant was asked to label what, in their opinion, were the four most important factors. The labels could be applied to a single factor or any combination of factors. A participant could also give all their votes to the same single factor, if they felt very strongly about it. The factors with the most votes were selected for the next step, which was generating variant states for each factor. A repeat round of voting was an option in the event of tied votes.

Factor variants

For each key factor selected, participants were asked to identify four plausible variant states. This was done in small groups (3-4 people). Examples were given so as to clarify the process. For instance, if disease transmission is a key factor, examples of variant states could be: very infectious, not very infectious, waterborne or transmitted by bodily fluids.

Scenario building

1 2

44 ⁴⁵ 195

2005657 **201**

After variants were devised, simple scenario building was demonstrated to participants. This was done by going person to person, with each person choosing one variant for each factor, making sure that the variants selected could plausibly happen together. After each variant was chosen, that same variant was not available for the other scenarios. Once all factor variants were chosen, they were combined to form a 'skeleton' for each scenario. Subsequently, small groups (4-6 persons) worked on backcasting for their specific scenario, starting in the projected year (2023) and working backwards to fill in the story of how the scenario might happen from our starting point in late 2017.

Participants' feedback

A feedback questionnaire was distributed to the participants to assess their opinions about the scenario planning workshop (Appendix 1).

Results

Participants

Thirty-one participants attended the workshop event in Barcelona, Spain, held on 7 November 2017. They had expertise in microbiology, medicine, infrastructure, data analysis, environmental and/or public health and facility management. Most worked for water companies, universities or government agencies. Most were professionally based in Spain, but there were participants concurrently working in seven other countries: Austria, Denmark, France, Netherlands, Norway, Portugal and UK. Fifteen experts were placed in the emerging pathogen group, and sixteen in the attack group.

Factors and variants

The key factors and variants chosen by the participants for the emerging pathogen (ED) and attack (AtK) scenarios are shown in Tables 2 and 3, respectively. The ED group identified eight key factors, while the AtK group identified five. For each key factor, four variants were successfully formulated. Three key factors were shared between the two scenarios, namely emergency response, communications and pathogen characteristics/ source. Interestingly, the scope of these factors was described somewhat differently between the two groups, which is very much allowed within scenario planning methods. The ED group members were quite concerned with pathogen characteristics (infectious dose and inactivation methods) and modes of transmission, while the AtK group considered pathogen characteristics as a mix of pathogen types. One variant for source of pathogen (variant 4) included infectious dose, persistence and incubation period. Similarly, communication strategies identified by the ED group were multi-faceted and revolved around the efficacy of a potential crisis committee, interaction with media and political influences. Conversely, the AtK group described only the outcome of whatever communication efforts were in place. The key factor variants were poor, perfect, disastrous, etc. This was also the case for the key factor emergency response. The AtK group derived variants such as none, incomplete and complete. The ED group categorised the

emergency response and defined three key areas: health systems, official communication and compliance. All three components were considered while deriving variants (Table 2 and 3). The AtK group considered access to healthcare as a stand-alone key factor. Finally, the AtK group identified areas that were exposed to the contaminated water as a key factor. This is particularly relevant considering the uncertainties about the scale of the malicious attack.

Outline Scenarios

When building the scenarios, one group found scenario construction too difficult; they were inclined to think that detection methods (genomic analysis, specifically) made their scenario too unlikely to happen. However, five of the scenarios (two emerging disease and three attack) were well enough developed to be reported here. It was commonly mentioned in all scenarios that lack of resources or policy changes that led to poor equipment maintenance or substandard monitoring of contamination or other reduced microbe control measures could considerably exacerbate the worst impacts of any threat scenario. All scenarios had to make plausible assumptions about decisions and policies adopted by health systems, governments and others. Policy changes could be to reduce frequency of testing, remove types of testing, remove chlorine from water. Some of the suggested details in all of the scenarios are sensitive, especially the specific mechanics of how an attack could happen. For this reason, and for brevity, we publish here only the timelines for each scenario, which start five years in the future and work backwards to present day. The variants used to build each scenario are listed in Boxes (1a-5a). Corresponding scenario timelines are listed in Boxes (1b-5b).

Evaluation

26/31 (84%) individuals returned completed questionnaires to the organisers. Feedback was broadly very positive (see Table 4).

23 24 42 43 44 52 ⁵⁶ **261**

Table 2 | Factors and variants chosen and for the emerging diseases scenarios

Key Factor	Variant 1	Variant 2	Variant 3	Variant 4	
Attack rate (AR), Morbidity (Mb) and Mortality (Mt)	AR: Just the already ill, young, very old Mb: Only a few people who are sick will have severe illness Mt: no fatalities, diseases treatable	transmission communicate requires qua few 1/10,000 gets ill requires qua who are sick Mb: Symptoms more Mb: Symptoms resevere severe but not lethal; severe, infection care at home monitoring 8 possible treatment		AR: high transmission rate Mb: Symptoms are very severe, fast progression Mt: Risk of death increases if no treatment within 72 hours	
Emergency response	rgency Health systems (HS) HS cannot cope with		HS are coping, official communications not effective, people expected to comply with instructions	High demand on HS, official communications will be very forceful, people are expected to have poor compliance	
Transmission	Waterborne only	Waterborne spread & person to person	Airborne & person to person & waterborne spread	Varies due to seasons or vulnerability of individuals	
Identification of outbreak and source	Source identified, containable as route is known	Source identified, but cannot be contained (route unknown)	Cannot identify source, situation can be contained	Source cannot be identified, route unknown, no containment	
Communica- tions	Crisis committee is organized with designated spokesperson, good liaison with political institutions and press	Crisis committee is organized with designated spokesperson, there is supervision of media and political influences, press conferences occur	Crisis committee is organized with designated spokesperson but no social media and official communication via radio or TV	No crisis committee is organized, so no spokesperson or monitoring / interaction with social media or press	
Human resources and contingency plan	Both Available and skilled	Both available but lack skills			
Analytical technologies	Fast, cheap, standardized and specific	Fast, expensive, specific, not standardized	Cheap, specific, slow, not standardized	None available but other AT could be adapted	
Characteris- tics of pathogen	Low infectious dose, inactivated by chlorination	not inactivated by inactivated by dose, not chlorination chlorination inactivated		High infectious dose, not inactivated by chlorination	

263

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40

264 265

Variant 1 Variant 2 Variant 3 Variant 4 **Key Factor** Inter-agency None Incomplete Complete, but Complete & cooperation personnel not personnel trained (emergency trained response plans) Mix of pathogens Virus Not detected Highly infective Source of eradicated or by normal low dose, uncommon in control pathogen persistent in Europe systems water, long incubation period **Areas Water** Small area, high City wide, 1 house, high City wide, medium Reached impact, medium impact, not impact, chlorinated, impact, 30% chlorinated, chlorinated, branched network reticulated branched not network chlorinated, network reticulated **Communications** Perfect Complete Good message, Poor targeting led communications in Disaster in all Strategy/ but at wrong to unnecessary Management of every way people public scare, total ways **General Public** chaos Access to Hospitals shutting Public access Hospitals are Violence or panic healthcare down due to hindered due overcrowded inside and around contamination, to many hospital staff and /or factors shortages, complications

Table 3 | Factors and variants chosen for the attack scenario

Box 1a | Variants used to construct scenario described in Box 1b

(Attack rate, morbidity, mortality) AR: Only vulnerable groups are susceptible (young, very old); Mb: Only a few people will have severe sickness; Mt: no fatalities expected, disease is treatable

(Emergency Response) Health systems cannot cope with demand, official communications are effective, can expect good public compliance with instructions

(Transmission) Varies due to season and vulnerability of individuals

(Identification of outbreak and source) Source identified, but cannot be contained as route unknown

(Communications) Crisis committee is organised. There is a designated spokesperson but no social media or official media communication via TV, radio, printed press

(Human resources) None (eg., due to strikes), plus no contingency plan, lack of skilled personnel

(Analytical technologies) AT are cheap, but not standardised, are time consuming, are specific

(Characteristics of pathogen) Low infectious dose, inactivated by chlorination

Box 1b | Emerging disease scenario (1)

Olympic Games Paris 2024

- 2023: Outbreak in France (1 year before the games) affecting persons already ill, limited severity and no fatalities. Pathogen: low infectious dose but inactivated by Chlorine
- 2022: Because of the Games, French government increases recreational areas with water fountains, playgrounds, more public drinking water fountains
- 2021: Summer with high temperatures, linked to increasing pollution after US pulled out of Paris Climate Change Convention
- 2020: EU allows decrease in Chlorine levels, French water supplies go Chlorine-free
- 2019: EU stops investing in new water safety tools and nobody is submitting related proposals, strikes because of low salaries, tax increases and pension age increase lead to health systems becoming vulnerable and no supervision of social media or official social media strategy
- 2018: Launches of new platforms and tools for rapid monitoring & pathogen detection but health authorities only recognise traditional/conventional methods -> new tools are not recognised as gold standard

Box 2a | Variants used to construct scenario described in Box 2b

(Attack rate, morbidity, mortality) AR: High transmission rate; Mb: Symptoms are very severe, fast progression; Mt: Risk of death increases if no treatment within 72 h

(Emergency Response) High demand on health services, official communications are very forceful, people are expected to have poor compliance

(Transmission) Airborne and person to person and waterborne

(Identification of outbreak and source) Cannot identify, but can be contained

(Communications) Organised crisis committee has designated, spokesperson official who answers to social media and there is agreement among political institutions who are included in the crisis committee. There are official communications with the press

(Human resources) There are resources available, contingency plan available, skills are available

(Analytical technologies) None suitable for pathogen, but other AT could be adapted

(Characteristics of pathogen) High infectious dose required, is NOT inactivated by chlorination

Box 2b | Emerging disease scenario (2)

Everywhere Unknown

- 2023: Large outbreak across EU produced by an Unknown microbe that is transmitted by air, person to person, waterborne. Infection dose is high but microbe is resistant to chlorine.
- 2022: Repairs are completed that happened after the 2021 overflows, to infrastructure for drinking water and waste water. However, this meant large investments that diverted resources away from research and development.
- 2021: Big overflows overwhelm drinking and waste water infrastructure (causing huge damage); concurrent critical economic situation.
- 2019; Crisis committee identified the problem (partly) and communicated to public health agencies how they should treat future cases.
- 2017: Sporadic cases of an unknown opportunistic microbe, virulence of which depends on environmental conditions (unknown).

Box 3a | Variants used to construct scenario described in Box 3b

(Inter-agency cooperation and emergency response plans) Complete, but personnel not trained

(Source of pathogen) Not detected by normal control systems

(Areas of water reached) City wide, medium impact, chlorinated and branched network

(Communications strategy & management of general public) Perfect communications in every way

(Access to Health Care) Hospitals are overcrowded

Box 3b | Attack scenario (3)

Disgruntled Employee

- 2023: City-wide outbreak with medium impact not detected by normal control systems during especially severe flu season which is overcrowding hospitals. However, there is good communication and a complete response plan (despite the lack of trained personnel)
- 2023: Bad guy makes attack plan, does not want to kill but expose company as unready.
 Gets pathogen from black market source.
- 2022: Bad guy gets fired from water supply Company
- 2020: New detection kits become available and are incorporated into response plans, no resources to train personnel with new kits
- 2017: Water supply company realises that traditional communications are no longer valid and make new plans using newer communications media

Box 4a | Variants used to construct scenario in Box 4b

(Inter-agency cooperation and emergency response plans) Incomplete inter-agency co-operation

(Source of pathogen) Mix of pathogens

(Areas of water reached) Small area, high impact, chlorinated, reticulated

(Communications strategy & management of general public) Complete disaster

(Access to Health Care) Hospitals are shutting down due to complications from staff shortages

Box 4b | Attack scenario (4)

Dystopia 2023

- 2023: Unknown actors contaminate water supply to the National Assembly building and surrounding area, using multiple different pathogens some of which are hard to identify. Appears politically motivated. Response is poorly managed with different agencies not cooperating. Inter-agency co-operation break-down because Health Dept. does not want to reveal internally known weaknesses in staffing levels. Security services (civil protection) didn't communicate the known risk to the water system to avoid revealing weaknesses in their system and their capabilities. There is high degree of inter-service rivalry in the background exasperated by a recent funding crisis. Multiple conflicting messages are sent by each agency to the media by different services to try to make their agency visible to the public. The mix of pathogens suggests this attack is backed by a highly sophisticated sponsor (could be State or well-resourced internal opposition). This sponsor sows additional confusion by using other assets to spread misinformation online
- Late 2022: Budget crisis due to lack of political direction and divided National Assembly
- Early 2022: Victorious political party is mired in corruption scandals left over from the election and does not command a majority. Money seems to be going missing or is not spent effectively. Public services (hospitals and public health in particular) are being shut down or reduced in services they offer due to lack of money
- Late 2021: Disputed and acrimonious election with many claims of corruption. Very little trust in politicians. Some indications of external interference in the elections
- Early 2021: Government falls following release of emails from parliamentary systems showing high levels of misbehaviour in public office (fiscal and other)
- Late 2020: Corrosive atmosphere to public; each political party seems to only represent narrow interests
- 2019: Increasing polarisation in the media. Increasing effect of "filter bubbles" in people's online life. Increasing intolerance of views different to one's own. Attempts to rewrite history by removal of public monuments and statues

Box 5a | Variants used to construct scenario described in Box 5b

(Inter-agency cooperation and emergency response plans) None

(Source of pathogen) Virus eradicated or uncommon in Europe

(Areas of water reached) City wide, medium impact, 30% of supply not chlorinated, reticulated

(Communications strategy & management of general public) Poor targeting leading to unnecessary public scare, total chaos

(Access to Health Care) Violence or nanic inside and around hospital

Box 5b | Attack scenario (5)

Ebola in Europe

- 2023: Attack on water supplies (Ebola virus injected into storage tanks) in large city in target European country. There is chaos with public panic, widespread misinformation about risks, impacts and safety measures that public can take.
- Late 2022: West African terrorists invite and help middle Eastern terrorists to collect and concentrate Ebola virus from faeces (suicide missions)
- Early 2022: Large Ebola outbreak starts in populous West African country
- 2018-2023: Debt crisis, cut in public services in target country. Leads to poor quality hospital services and few resources to maintain good quality public communications. Emergency response plans forgotten or abandoned or become very out of date. Capital investment in protecting and maintaining water supply network is especially badly affected, in large European city (or cities).

301 52

305

Table 4 | Results of feedback questionnaire

Criteria	Results
This workshop was relevant to my needs	81% agreed or strongly agreed
This workshop met my expectations	70% agreed or strongly agreed
This workshop helped me understand scenario planning better	97% agreed or strongly agreed
I enjoyed this workshop and I am glad I came to it	100% agreed or strongly agreed
I understood what to do in each phase of today's workshop, I didn't feel confused.	31% agreed or strongly agreed (46% unsure)
The content was well-organised	84% agreed or strongly agreed
The workshop has made me think about unusual risks in a helpful way	80% agreed or strongly agreed (20% unsure)
This workshop identified risk factors I never thought of before	50% agreed or strongly agreed (46% unsure)
The workshop made me feel better able to prepare for future unusual events	50% agreed or strongly agreed (42% unsure)
I could adapt these scenario methods to use in my own organization.	76% agreed or strongly agreed (20% unsure)
This workshop helped me see how to reduce possible impacts or better manage relevant risks at work	46% agreed or strongly agreed (42% unsure)

Discussion

We describe a set of procedures for running a one-day workshop to facilitate experts to envision a range of plausible scenarios that could threaten water supplies they are tasked to protect. Although the threats poised here were purely biological, the methods could easily be adapted to consider other hazards, such as chemical or radiological threats. Readers can judge for themselves if a similar workshop would be valuable to run within their organisation. The workshop steps described here are not definitively the best way to undertake such a workshop; we describe ways to improve the implementation below. Neither were our resulting scenarios definitive; a different set of experts probably would have identified somewhat different key factors and developed possibly quite different scenarios. The choices of key factors and how participants imagined scenarios was heavily influenced

by their own backgrounds (and therefore could be very country-specific). Nevertheless, we hope that our account and comments about implementation may be useful to others pondering whether to run similar planning exercises within their own organisations, in order to inform local risk assessment. Some of our observations described below seem likely to be very generalizable to many settings, with regard to protecting water supplies. A valuable outcome of scenario planning exercises like ours is going beyond worst and best case scenarios; looking for extremes is prone to strong human biases. There are an infinite number of scenarios for any one topic, and scenarios need to be adaptable to local conditions in order to be credible and not just very unlikely extremes. The intermediate steps in the methods we describe can also be a constructive exercise for those involved in water protection, such as identifying key factors (inherently important and uncertain), and the use of silent decision making (the silent sorting methods which encourage the widest range of views to emerge).

Real world implications

Participants found it difficult to plausibly imagine these scenarios except in an environment where protection measures and communications strategies were significantly inadequate. This is encouraging as it suggests an inverse situation is true in the group opinion: the scenarios were very unlikely as long as regulatory standards remain high, and monitoring and accountability mechanisms are well-resourced. Some of the participants with microbiology backgrounds felt strongly that genomic analysis would ensure very rapid identification of pathogen characteristics, so much so that delay in identification or difficulties in planning a management strategy would be negligible. Developments in genomic methods may indeed facilitate quicker removal from water supplies. However, the converse is also true. In an environment with inadequate safety controls, insufficient laboratory methods, poor resources or monitoring, the participants easily imagined a high diversity of ways that an emerging disease or attack could be highly disruptive to provision of safe drinking water in Europe.

Among the many policy changes that could increase likelihood of an emerging pathogen or successful attack were: declines in capital investment, delays in repairing infrastructure, prioritising other public services or promoting public preferences (such as having less chlorine taste in drinking water), as well as reduced intra-agency cooperation. The participants thought such policy changes were unlikely but not impossible, especially given the ever-present pressure on public services to be cost-effective. Policies that would reduce negative scenario impacts were to maintain effective communications with

the public and between relevant agencies, as well as well-defined strategies to make water quality testing easier, quicker, more specific and yet still cost-effective.

Event Implementation

The facilitators see many ways that the methods could have been concisely and clearly better explained. We were not prepared for how systematically and thoroughly the participants wanted to approach the task of choosing variants: we should have been firmer about insisting that they think of variants in concise terms. Many participants tried to devise elaborate scenarios as backstories for a possible variant, before they could decide on each variant option. It is certainly possible to dedicate an entire day to decide on factor variants, but for the purposes of scenario building in a one-day workshop, the variants only needed to be possible, rather than especially plausible. The time spent developing and identifying possible backstory details for the variants was still productive, ultimately, because these details were useful when fleshing out details during the scenario building stage.

Operational perspectives

In plenary discussion afterwards, attendees raised further observations about testing regimes. They commented that current methods for water quality testing are well suited for the required regulatory monitoring. However, deference to regulatory standards somewhat acts as a deterrent to trying new or more sensitive testing methods; since the new methods wouldn't be required and may represent an unnecessary and unreimbursed cost. There are also problems with accepting results from new techniques because of their lack of correspondence with data accumulated from conventional monitoring. Some methods are associated with different error patterns (such as higher numbers of false positives), and thus require sensitive decision making with regard to when an alert truly needed to be raised.

The regulatory sector should be receptive to new methods, but getting new methods into practice is often a slow process because of conservationism in the sector. Additional limitations are the usually high cost of newly developed methods. Therefore, the participants agreed that a distinction should be made between regulatory compliance and operational monitoring. It was suggested that a first step in operational implementation is event monitoring. This has been trialled since late 2017 in Spain (Brainard et al. 2018).

402

Relatively new technologies such as whole genome sequencing for typing and source tracking are relevant. These techniques offer value in outbreak investigation for tracing sources of contamination but are unlikely to be unsuitable for routine application in environmental sampling, as the large amounts of data generated could not necessarily be linked to health risks in humans. Microbial source tracking and whole genome sequencing techniques have been developed (Hjelmsø et al. 2017) which may be very valuable in bioterrorist attack investigations. Water quality testing strategies can be most protective when implemented in a tiered approach, such that test results are enhanced by complementary information from multiple sources (Rickert et al. 2014, Ryzinska-Paier et al. 2014).

Limitations

As observed previously, the key factors and scenarios identified by our participants cannot be definitive and may not all be generalizable, instead they are the products of the procedures as described. With regard to our procedures, we have tried to explain them clearly and be candid about what could be improved. We did not subject the workshop or the outputs to a rigorous evaluation scheme. Checklists for evaluating quality of participatory scenario planning exercises are not well-developed, but many resources exist from which such a checklist could be derived, some of which are contextualised with regard to water safety. Acosta Tellez (2014) describes different scenario planning approaches and their strengths and weaknesses when assessed against many evaluation criteria. Scott *et al.* (2012) described how scenario planning must identifying important but uncertain factors, their impacts and ultimately key adaptations to reduce risks of harm. Both Mott Lacroix *et al.* (2015) and Van der Merwe (2008) advocated that scenario planning exercises should produce outputs that are challenging, relevant and plausible; we believe that our scenario planning had many if not all of these attributes.

Conclusions

The scenario building workshop fostered increased awareness of possible risks to water supplies. The methods comprise a process that has credibility for participants by creating a multiplicity of considered perspectives in the process of identifying what are the key uncertain and important factors, and their possible variant states, that could have a high impact on the proposed problem situation. This meeting encouraged health and industry experts to identify vulnerabilities and novel pathways for an emerging disease or attack to threaten water supplies. Very positive feedback about the experience was received, even among those participants who found their scenario was very unlikely. Scenario building could be used to inform Water Safety Plans (Bridle *et al.* 2014, Ryzinska-

Paier et al. 2014, Gunnarsdottir et al. 2016, Gunnarsdottir et al. 2017) for better risk assessment and to improve protection of public water supplies.

Acknowledgements

Funding and support was provided by the EU-FP7 Aquavalens project (Grant agreement number: 311846). Mike van der Es, Claudia Serra Puigdomenech and Robert Pitchers gave helpful comments on this manuscript as it developed. Authors JB, CCH and PRH are affiliated to the UK National Institute for Health Research (NIHR) Health Protection Research Unit in Emergency Preparedness and Response in partnership with Public Health England (PHE). The views expressed are those of the authors and not necessarily those of the European Union, UK National Health Services, the NIHR, PHE or workshop participants. The funders had no role in event design, workshop delivery, collection of feedback, analysis, decision to publish, or preparation of the manuscript. Thank you to all of our participants for their intellectual input and expertise, creative and collaborative contributions, as well as Aquavalens partners cetAqua and Aigües de Barcelona who co-hosted the meeting.

References

Acosta Tellez, C. (2014). State of the art of scenario planning: 40.

Baldursson, S. and P. Karanis (2011). "Waterborne transmission of protozoan parasites: review of worldwide outbreaks-an update 2004-2010." Water Research 45(20): 6603-6614.

Brainard, J., C. Hammer, M. Bouzid, R. Pitcher and P. Hunter (2018). Protecting the health of Europeans by improving methods for the detection of pathogens in drinking water and water used in food preparation: 32.

46 433

Brainard, J. and P. R. Hunter (2016). "Contextual Factors Among Indiscriminate or Large Attacks on Food or Water Supplies, 1946-2015." Health Security 14(1): 19-28.

Bridle, H., B. Miller and M. P. Desmulliez (2014). "Application of microfluidics in waterborne pathogen monitoring: A review." Water Research 55: 256-271.

55 439

Clark, R. M. and R. A. Deininger (2000). "Protecting the nation's critical infrastructure: The vulnerability of US water supply systems." Journal of Contingencies and Crisis Management 8(2): 73-80.

443 1 444 Gleick, P. and M. Heberger (2014). Water and Conflict: Events, Trends and Analysis (2011-2012). The 445 World's Water. A. N. (ed). Washington DC, Island Press: 159-171. 3 4 446 5 447 Gunnarsdottir, M. J., S. M. Gardarsson, G. S. Jonsson and J. Bartram (2016). "Chemical quality and 6 regulatory compliance of drinking water in Iceland." International Journal of Hygiene and 448 7 8 449 Environmental Health 219(8): 724-733. 9 10 450 ¹¹ 451 Gunnarsdottir, M. J., K. M. Persson, H. O. Andradottir and S. M. Gardarsson (2017). "Status of small 12 452 water supplies in the Nordic countries: Characteristics, water quality and challenges." International 13 453 <u>Journal of Hygiene and Environmental Health</u> **220**(8): 1309-1317. 14 15 454 16 ₁₇ 455 Haimes, Y. Y., N. C. Matalas, J. H. Lambert, B. A. Jackson and J. F. Fellows (1998). "Reducing vulnerability of water supply systems to attack." Journal of Infrastructure Systems 4(4): 164-177. 18 **456** 19 20 457 21 458 Hjelmsø, M. H., M. Hellmér, X. Fernandez-Cassi, N. Timoneda, O. Lukjancenko, M. Seidel, D. Elsässer, 22 459 F. M. Aarestrup, C. Löfström and S. Bofill-Mas (2017). "Evaluation of Methods for the Concentration 23 460 and Extraction of Viruses from Sewage in the Context of Metagenomic Sequencing." PloS One 12(1): 24 461 e0170199. 25 26 27 **462** 28 **463** Kahn, H. and A. J. Wiener (1967). "The next thirty-three years: a framework for speculation." ²⁹ 464 Daedalus: 705-732. 30 ³¹ 465 32 466 Kwakkel, J. H., W. E. Walker and M. Haasnoot (2016). "Coping with the wickedness of public policy 33 467 problems: approaches for decision making under deep uncertainty." Journal of Water Resources 34 35 468 Planning and Management 142(3). 36 37 469 38 470 Meissner, P. and T. Wulf (2013). "Cognitive benefits of scenario planning: Its impact on biases and 39 471 decision quality." Technological Forecasting and Social Change 80(4): 801-814. 40 ⁴¹ 472 42 473 Mott Lacroix, K., A. Hullinger, M. Apel, W. Brandau and S. B. Megdal (2015). Using Scenario Planning 43 44 474 to Prepare for Uncertainty in Rural Watersheds, College of Agriculture, University of Arizona 45 475 (Tucson, AZ). 46 47 476 48 477 Mutchek, M. and E. Williams (2014). "Moving towards sustainable and resilient smart water grids." 49 478 <u>Challenges</u> **5**(1): 123-137. 50 51 479 52 480 Qiao, J., D. Jeong, M. Lawley, J.-P. P. Richard, D. M. Abraham and Y. Yih (2007). "Allocating security 53 ₅₄ 481 resources to a water supply network." <u>IIE Transactions</u> **39**(1): 95-109. 55 56 482 57 **483** Rickert, B., O. Schmoll, A. Rinehold and E. Barrenberg (2014). Water safety plan: A field guide to ⁵⁸ 484 improving drinking-water safety in small communities: 91. 59 ⁶⁰ 485

	486	Ryzinska-Paier, G., T. Lendenfeld, K. Correa, P. Stadler, A. Blaschke, R. Mach, H. Stadler, A. Kirschner
	487	and A. Farnleitner (2014). "A sensitive and robust method for automated on-line monitoring of
2	488	enzymatic activities in water and water resources." <u>Water Science and Technology</u> 69 (6): 1349-1358.
4	400	
5	489	Schwartz, D. (1006). The Art of the Long View, Planning for the Euture in an Uncertain World
6	490	Schwartz, P. (1996). The Art of the Long View: Planning for the Future in an Uncertain World,
7	491	<u>Currency</u> . New York, Double Day.
8 9	492	
10	493	Scott, C. A., C. J. Bailey, R. P. Marra, G. J. Woods, K. J. Ormerod and K. Lansey (2012). "Scenario
	494	planning to address critical uncertainties for robust and resilient water—wastewater infrastructures
12	495	under conditions of water scarcity and rapid development." Water 4(4): 848-868.
13		· · · · · · · · · · · · · · · · · · ·
14 15	496	
16	497	Sinclair, R., E. Jones and C. Gerba (2009). "Viruses in recreational water-borne disease outbreaks: A
17	498	review." <u>Journal of Applied Microbiology</u> 107 (6): 1769-1780.
18 19	499	
20	500	Steinmüller, K. (1997). "Grundlagen und Methoden der Zukunftsforschung: Szenarien." Delphi,
21	501	Technikvorausschau, Gelsenkirchen: Sekretariat für Zukunftsforschung (Werkstatt Bericht 21).
22	301	recimitivorausseriau, deiserikirerien. Sekretariat far Zakariresforseriang (Werkstatt Berfent 21).
23	502	
24 25	503	Van der Merwe, L. (2008). "Scenario-based strategy in practice: a framework." Advances in
26	504	<u>Developing Human Resources</u> 10 (2): 216-239.
27	F0F	
28	505 506	Wilson, I. (1978). Scenarios. <u>Handbook of Futures Research</u> . J. Fowles. Westport and London,
29 30	507	Greenwood Press.
31	307	Greenwood Fress.
32	508	
33		
34 35	505	
36	510	
37		
38		
39 40		
41		
42		
43		
44 45		
46		
47		
48		
49		
50 51		
52		
53		
54		
55 56		
57		
58		
59		
60 61		
62		
63		
64		

Please complete the evaluation for today's workshop – your feedback is valuable.							
Topic = \square Emerging Infectious Disease	2	Delibe	rate Attack				
You do not need to write your name or	this form,	we want	your honest opin	nions. Pleas	e tick how		
much you agree or not.	·						
Criteria	Strongly Agree	Agree	Neither agree nor disagree	Disagree	Strongly Disagree		
This Workshop was relevant to my							
needs							
This workshop met my expectations							
This workshop helped me understand scenario planning better							
I enjoyed this workshop and I am							
glad I came to it							
I understood what to do in each							
phase of today's workshop, I didn't							
feel confused.							
The content was well-organised							
The workshop has made me think							
about unusual risks in a helpful way							
This workshop identified risk factors							
I never thought of before							
The workshop made me feel better							
able to prepare for future unusual events							
I could adapt these scenario							
methods to use in my own							
organization.							
This workshop helped me see how							
to reduce possible impacts or better							
manage relevant risks at work							
Do you have any other comments, sucl today?	h as what c	ould we l	nave improved, o	r what did y	ou especia		
e will NOT share any identifying informations with the ware identified in today's we		•	•	•			
your anonymous replies on this form a would-be attackers, to be shared more	and other a	nonymo	us contributions t	o this works	shop that a		
lete any category that you do not want	•			•			
Yes		No					