

# A Novel CMAQ-CNN Hybrid Model to Forecast Hourly Surface-Ozone Concentrations Fourteen Days in Advance

---

Alqamah Sayeed<sup>1</sup>, Yunsoo Choi<sup>1\*</sup>, Ebrahim Eslami<sup>1</sup>, Jia Jung<sup>1</sup>, Yannic Lops<sup>1</sup>,  
Ahmed Khan Salman<sup>1</sup>, Min-Hyeok Choi<sup>2</sup>, Hyun-Ju Park<sup>2</sup>, and Jae-Bum Lee<sup>2</sup>


Presenter: Dr. Yunsoo Choi (ychoi6@uh.edu)

<sup>1</sup> Department of Earth and Atmospheric Sciences, University of Houston

<sup>2</sup> National Institute of Environmental Research, Air Qualify Forecasting Center

CMAS Conference, October 2020

# Neural Network Model



# Our deep learning application on short-term forecasting

---

- ❑ Prediction of Air pollutants

  - ❑ Ozone

  - ❑ Pollen

  - ❑ PM

- ❑ Prediction of various weather phenomenon

  - ❑ Wind Speed

  - ❑ Temperature

  - ❑ Relative Humidity

  - ❑ Precipitation

- ❑ Bias correction of numerical modeling outputs (e.g., from CMAQ and WRF)


# Motivation of this study

---


- ❑ Is there a hybrid model to combine numerical and deep learning modeling?
- ❑ Can this hybrid model be used to forecast long-term (e.g., one week or two weeks) concentrations of air pollutants?

# Data Set-up


---



# Data Set-up



# Model Architecture



# Loss Function

---


- ❑ Method to Evaluate the Performance of Model on a Sample Dataset.
- ❑ Common Loss Functions:
  - ❑ Mean Squared Error (Method 1)
  - ❑ Mean Absolute Error
  - ❑ Mean Absolute Percentage Error etc.
- ❑ Loss Function
  - ❑ Index of Agreement (Method 2)

$$IOA = 1 - \frac{\sum (O_i - P_i)^2}{\sum (\text{abs}(O_i - \bar{O}) + \text{abs}(P_i - \bar{O}))^2}$$




# Study Area and Data

- 255 Air Quality Monitoring Station of South Korea
- WRF Surface Meteorology Processed by MCIP (Meteorology-Chemistry Interface Processor).
- CMAQ predicted Surface Chemistry (Air Pollutant)
- Training: 2015-2017 Forecasting:2018




# Results : Based on Loss Function




MSE – Method 1  
IOA – Method 2


# Results : Different Method Comparison



# Results : Performance Comparison (Days)



# Results




# Factors Affecting Model's Performance


---

- ❑ Distance from Coast
- ❑ Base Model Performance (CMAQ)
- ❑ Ozone diurnal variation
- ❑ Urbanization

# Distance from Coast









# Dependence on CMAQ







# Ozone diurnal variation



# Urbanization



Source: Chan, C. H., Caine, E. D., You, S. & Yip, P. S. F. Changes in South Korean urbanicity and suicide rates, 1992 to 2012. *BMJ Open* 5, e009451 (2015).

# Summary & Funding

---

- ❑ Developed a Hybrid CMAQ-CNN model to predict 2-week Hourly Ozone Concentration
- ❑ Model significantly improved performance from the CMAQ Day 1.
- ❑ Average IOA for 1<sup>st</sup> Day was 0.91 that gradually reduces to 0.78 for 14<sup>th</sup> day.
- ❑ Computational time was significantly reduced.
- ❑ This approach can be used for other pollutants and weather variables.
  
- ❑ Funding: The National Institute of Environment Research (NIER), NIER-2018-04-02-056