

9. A teenager who works in a dog kennel after school has had a skin rash, eosinophilia and an enlarged liver and spleen for 2 years. Which of the following is the most likely cause of this infection?

- a. Giardiasis
- b. Toxoplasmosis
- c. Trichinosis
- d. Cysticercosis
- e. Visceral larva migrans

16. Cardiomyopathy is a complication of:

- a. Chagas disease
- b. Black water fever
- c. Kala azar
- d. Extra intestinal amoebiasis
- e. Dermal leishmanoid

22. When activated by free or heme-bound iron, it forms a carbon-centered free-radical compound that has the ability to alkylate many proteins such as PfATP6 as well as heme.

- A. chloroquine B. quinine
- C. sulfadiazine D. trimethoprim
- E. Artemisinin

27. Which of the following infections resemble malignancy

- a. Unilocular hydatid cyst disease
- b. Multilocular hydatid cyst disease
- c. Schistosome granuloma
- d. Sparganosis in human
- e. Trichinellosis

33. Ascariasis causes:

- a. Appendicitis
- b. Intestinal obstruction

- c. Haemorrhagic pancreatitis
- d. Intestinal intussusception
- e. All of the above

39. Napier's Aldehyde test is the serological test for diagnosis of

- a. Kala azar
- b. Oriental sore
- c. Chagas disease
- d. Dermal leishmanoid
- e. African sleeping sickness

40. It is contraindicated for the treatment of onchocerciasis because it causes severe reactions related to microfilarial destruction.

- A. piperazine
- B. praziquantel
- C. diethylcarbamazine (DEC)
- D. ivermectin
- E. antimony potassium tartrate

43. All are intracellular parasites except ()

- a. Leishmania donovani
- b. Toxoplasma gondii
- c. Plasmodium vivax
- d. Cryptosporidium parvum
- e. None of the above

55. Formed during chronic infections, slowly multiplying parasites within the cyst is known as –

- a. Tachyzoites
- b. Bradyzoites
- c. Phanerozoites
- d. Cryptozoites
- e. Merozoites

71.人兽共患寄生虫病中的家畜或野生动物，在流行病学上是该种人体寄生虫的：
()

- A.definitive host
- B.intermediate host
- C.reservoir host
- D.paratenic host
- E.以上都不是

79.下列虫种，哪个进入体后，先在肺部停留并发育，然后才到正常寄生部位寄生：

- A.Taenia solium
- B.Clonorchis sinensis
- C.Ascaris lumbricoides
- D.Enterobius vermicularis
- E.Hymenolepis nana

86.Concomitant Immunity' is shown by

- a. Enterobius vermicularis
- b. Wuchereria bancrofti
- c. Brugia malayi
- d. Ancylostoma duodenale
- e. Schistosoma japonicum

116.The antimalarial activity is probably attributable to quinone, a metabolite of the following drug that interferes with the function of ubiquinone as an electron carrier in the respiratory chain.

- A、chloroquine
- B、artemisinin
- C、quinine
- D、primaquine
- E、pyrimethamine

128.Morula cells with IgM deposits are seen in –

- a. Kala azar
- b. Sleeping sickness
- c. Classical filariasis
- d. Toxoplasmosis
- e. Neurocysticercosis

139.Select the false statement regarding Guinea worm

- a. It is mentioned as “Fiery serpent” in the Bible
- b. The female worm protrudes through blister to lay eggs
- c. The next parasitic disease mentioned in WHO’s eradication list
- d. The natural habitat of the worm is subcutaneous tissue in human
- e. None of the above

161.Sputum examination is useful in diagnosis caused by following except –

- a. Trichuris trichura
- b. Ancylostoma duodenale
- c. Paragonimus westermani
- d. Strongyloides stercoralis
- e. None of the above

170.Calabar swellings are the characteristic clinical feature of – infections

- a. Ground itch
- b. Loa loa
- c. Angiostrongylus cantonensis
- d. Schistosome granuloma
- e. Blinding filarial

191.Central body of Manson’ in microfilaria extends from

- a. Posterior V spot to G1 cell
- b. Anterior V spot to G1 cell
- c. Excretory pore to G4 cell
- d. Nerve ring to G4 cell

e. None of the above

205.下列蠕虫中不属于线虫的是()

A.Trichinella spiralis

B.Trichuris trichiura

C.Enterobius vermicularis

D.Ascais lumbricoides

E.Pagumogonimus skrjabini

232.Parasite commonly causing pulmonary eosinophilia syndrome –

a. Pin worm

b. Trichinella spiralis

c. Hook worm

d. Whip worm

e. Filaria bancrofti

233.Cutaneous larva migrans is due to the larvae of

a. Ancylostoma duodenale

b. Ancylostoma braziliensis

c. Microfilariae bancrofti

d. Necator americanus

e. Toxocara canis

239.Creeping eruption is caused by - of hook worm

a. Rhabditiform larva

b. Filariform larva

c. Egg

d. Adult worm

e. Intermediate host with larvae

257.Select the false statement regarding Guinea worm

a. It is mentioned as “Fiery serpent” in the Bible

b. The female worm protrudes through blister to parturate

c. The next parasitic disease mentioned in WHO's eradication list

d. The natural habitat of the worm is subcutaneous tissue in human

e. None of the above

275."Occult filariasis" is a synonym for –

a. Larva migrans

b. River blindness

c. Tropical pulmonary eosinophilia

d. Classical filariasis

e. Cutaneous allergy to ascariasis

286.下列蠕虫不属于线虫纲的是

A.Enterobius vermicularis

B.Wuchereria bancrofti

C.Trichinella spiralis

D.Ascais lumbricoides

E.Pagumogonimus skrjabini

289.Wintern bottom sign in sleeping sickness refers to –

a. Posterior cervical lymphadenopathy

b. Narcolepsy

c. Unilateral conjunctivitis

d. Transient erythema

e. Canker sore

292.The best method for the detection of Strongyloides larvae is which of the following?

a. Baermann technique

b. Enzyme immuno assay

c. NIH swab specimen

d. Dilution followed by egg count

e. Sigmoidoscopy and mucosal lesion biopsy

299. Most common filarial infections in South India is

- a. *Wuchereria bancrofti*
- b. *Brugia malayi*
- c. *Loa loa*
- d. *Onchocerca volvulus*
- e. *Mansonella ozzardi*

308. Intestinal intussusception is a characteristic clinical feature of which parasitic infection?

- a. Ancylostomiasis
- b. Beef tapeworm infections
- c. Trichuriasis
- d. Enterobiasis
- e. Ascariasis

328. All are transmitted by faeco-oral route except –

- a. Ascariasis
- b. Giardiasis
- c. Balantidiasis
- d. Strongyloidiasis
- e. Enterobiasis

362. Intermediate host in the life cycle of *Dracunculus medinensis* is –

- a. Man
- b. Cray fish
- c. Cyclops
- d. Rat flea
- e. Female *Anopheles* mosquito

366. Chronic dysentery, abdominal pain and rectal prolapse in children is caused by

–

- a. Pin worm
- b. Trichinella worm
- c. Whip worm
- d. Round worm
- e. Guinea worm

385. Calabar swellings are the characteristic clinical feature of – infections

- a. Ground itch
- b. *Loa loa*
- c. *Angiostrongylus cantonensis*
- d. Schistosome granuloma
- e. Blinding filaria

392. Which among the following is viviparous :

- a. *Strongyloides stercoralis*
- b. *Trichinella spiralis*
- c. *Enterobius vermicularis*
- d. *Sparganum mansoni*
- e. None of the above

393. Unarmed scolex is the characteristic feature of –

- a. *Echinococcus granulosus*
- b. *Taenia solium*
- c. *Hymenolepis diminuta*
- d. *Taenia saginata*
- e. *Hymenolepis nana*

426. The following regarding cysticercosis are true except –

- a. The commonest sites are meninges and cerebral ventricle
- b. Calcification is common
- c. Causes focal neurological complications
- d. Found in subcutaneous tissues

e. Seen in both human and intermediate hosts

433.The following infections resemble malignancy –

- a. Unilocular hydatid cyst disease
- b. Multilocular hydatid cyst disease
- c. Schistosoma granuloma
- d. Sparganosis in human
- e. Trichinellosis

438.Casoni's test is done for the diagnosis of –

- a. Hook worm infections
- b. Neurocysticercosis
- c. Filarial Infections
- d. Bilharziasis
- e. Hydatid Cyst Disease

451.Which parasite produces an operculated egg?

- a. Taenia saginata
- b. Hymenolepis nana
- c. Strongyloides stercoralis
- d. Trichuris trichura
- e. Spirometra mansoni

454.Which parasite egg does not float in saturated salt solution –

- a. Hook worm
- b. Whip worm
- c. Pork tape worm
- d. Round worm – fertilized egg
- e. Pin worm

459.Which parasite's egg consist of polar filaments arising from either end of the

embryo –

- a. Taenia solium
- b. Diphylobothrium latum
- c. Echinococcus granulosus
- d. Hymenolepis nana
- e. Schistosoma japonicum

502.Dwarf tapeworm refers to –

- a. Echinococcus multilocularis
- b. Taenia solium
- c. Hymenolepis diminuta
- d. Hymenolepis nana
- e. Spirometra mansoni

533.下列蠕虫中属于绦虫的是()

- A.Echinococcus granulosus
- B.Entamoeba histolytica
- C.Schistosoma japonicum
- D.Wuchereria bancrofti
- E.Paragonimus westermani

577.The most commonly affected tissue in cysticercosis is –

- a. Brain
- b. Eye
- c. Muscle
- d. Liver
- e. Intestine

623.Sparganum is the name of which larvae –

- a. Oncosphere
- b. Proceroid
- c. Coracidium

- d. Plerocercoid
- e. Free swimming miracidium

705. Cholangiocarcinoma is caused by –

- a. *Paragonimus westermani*
- b. *Schistosoma japonicum*
- c. *Clonorchis sinensis*
- d. *Fasciolopsis buskii*
- e. *Fasciola hepatica*

759. Which parasitic egg does not undergo development in soil

- a. *Ancylostoma duodenale*
- b. *Taenia solium*
- c. *Ascaris lumbricoides*
- d. *Trichuris trichura*
- e. *Necator americanus*

784. *Schistosoma japonicum* resides in

- a. Vesical venous plexes
- b. Systemic circulation
- c. Gall bladder
- d. Splenic vein
- e. Liver

792. Select the true statement :

- a. *Cysticercus bovis* can be seen in human host.
- b. Schistosomulae of *Schistosoma japonicum* mature in liver.
- c. Miracidium of *Clonorchis sinensis* will have a free swimming life.
- d. The life span of microfilariae is about 7 days.
- e. *Schistosoma japonicum* has two generations of rediae

819. The snail host acting as intermediate host for *Schistosoma haematobium* in India is –

- a. *Bulinus truncatus*
- b. *Biomphalaria alexandrina*
- c. *Oncomelania*
- d. *Ferrisia tenuis*
- e. *Planorbis*

838. Gynaecophoric canal is seen in the –

- a. Female *Schistosoma*
- b. Male *Schistosoma*
- c. *Schistosoma cercaria*
- d. *Microfilaria* of *W. bancrofti*
- e. None of the above

849. Cercariae of *Schistosoma* has a maximum free swimming life of

- a. 70 days
- b. 70 hours
- c. 30 days
- d. 3 hours
- e. None of the above

866. 下列蠕虫中属于吸虫的是()

- A. *Ancylostoma duodenale*
- B. *Fasciolopsis buskii*
- C. *Taenia solium*
- D. *Hymenolepis nana*
- E. *Necator americanus*

894. Fairley's test is done in –

- a. Bancroftian filariasis
- b. Cysticercosis
- c. Dermal leishmanoid
- d. Kala azar
- e. Vesical Schistosomiasis

897.Which of the following may be injected with raw fish, affects the liver, and has an operculated egg ?

- a. Clonorchis sinensis
- b. Paragonimus westermani
- c. Schistosoma japonicum
- d. Schistosoma mansoni
- e. Hymenolepis nana

907.Clay pipe-stem cirrhosis' is the pathogenic feature of

- a. Katayama disease
- b. Hepatic amoebiasis
- c. Clonorchiasis
- d. Bilharziasis
- e. Hepatic hydatid cyst disease

975.Which among these is also known as 'Assasin bugs'

- a. Bed bug
- b. Reduvid bug
- c. Pediculus humanus
- d. Phthirus pubis
- e. Xenopsylla cheopis

980.Laboratory diagnosis of scabies is best done by:

- a. Fecal examination
- b. Serology
- c. Blood examination
- d. Skin scraping
- e. Rectal biopsy

984.Which mosquitos are known as Tiger mosquitoes

- a. Anopheles
- b. Culex
- c. Aedes
- d. Mansonia
- e. None of the above

1037.Human flea among these are

- a. Xenopsylla cheopis
- b. Pediculus humanus
- c. Phthirus pubis
- d. Sarcoptes scabies
- e. Pulex irritans

1043.The scientist who discovered the transmission of malaria by Anopheline mosquitoes –

- a. Laveran
- b. Paul Muller
- c. Ronald Ross
- d. Golgi
- e. Romanowsky

1049.Simulium acts as vectors for spreading –

- a. Trypanosoma cruzi
- b. Malayan filaria
- c. African eye worm
- d. Bancroft's filarial
- e. Blinding filaria

1064.Tse tse fly is not the vector for

- a. Trypanosoma rhodesiense
- b. Trypanosoma gambiense
- c. Trypanosoma brucei

- d. *Trypanosoma cruzi*
- e. None of the above

1102. Wingless insects among the following

- a. *Pediculus humanus*
- b. *Glossina palpalis*
- c. Reduviid bug
- d. *Phlebotomus argentipus*
- e. Black fly

1131. Which disease is transmitted by mites

- a. Babesiosis
- b. Pediculosis
- c. Myiasis
- d. Chagas disease
- e. Scabies

1155. 对人致病力较强的两种阿米巴原虫是()

- A. *Entamoeba histolytica* & *E. dispar*
- B. *Entamoeba histolytica* & *E. coli*
- C. *Entamoeba histolytica* & *Naegleria fowleri*
- D. *Entamoeba histolytica* & *Endolimax nana*
- Entamoeba histolytica* & *Dientamoeba fragilis*

1159. A butcher fond of eating raw hamburger develops chorioretinitis. A Sabin-Feldman dye test is done which came positive. The most likely cause of infection is –

- a. Giardiasis
- b. Trichinosis
- c. Schistosomiasis
- d. Toxoplasmosis

- e. Kala azar

1160. "Anchovy sauce pus" is formed in

- a. Intestinal Amoebic abscess
- b. Cerebral malaria
- c. Amoebic Liver abscess
- d. Algid malaria
- e. Neural cysticercosis

1161. The parasite that results in symptoms of duodenitis is:

- a. *Entamoeba coli*
- b. *Balantidium coli*
- c. *Entamoeba histolytica*
- d. *Giardia lamblia*
- e. *Isospora belli*

1162. Merozoites liberated from Exo-erythrocytic schizogony are called as

- a. Phanerozoites
- b. Rhoptries
- c. Bradyzoites
- d. Tachyzoites
- e. Cryptozoites

1167. 引起原发性阿米巴脑膜脑炎的病原体

- A. *Naegleria fowleri*
- B. *Entamoeba histolytica*
- C. *Entamoeba gingivalis*
- D. *Entamoeba hartmani*
- Endolimax nana*

1168. Malaria attacks (paroxysms) are caused by:

- a. release of toxins from RBC
- b. invasion of RBC by merozoites

- c. severe hemolytic anemia
- d. all of the above
- e. a & c

1170. "Accoli" form of Trophozoites are seen in

- a. Plasmodium falciparum
- b. Plasmodium ovale
- c. Plasmodium malariae
- d. Plasmodium vivax
- e. None of the above

1173. 下列原虫中属于孢子虫的有()

- canthamoeba culbertsoni
- B. Giardia lamblia
- C. Toxoplasma gondii
- D. Clonorchis sinensis
- ntamoeba histolytica

1176. Which species shows less days of pre erythrocytic schizogony among the following –

- a. Plasmodium vivax
- b. Plasmodium falciparum
- c. Plasmodium malariae
- d. Plasmodium ovale
- e. Plasmodium berghei

1177. 十二指肠引流检查法用于诊断

- A. Trichomonas vaginalis
- B. Entamoeba histolytica
- C. Giardia lamblia
- D. Entamoeba coli
- E. Trichomonas hominis

1181. Presence of Ziemann's dots in RBC's are characteristic feature of – infections

- a. Plasmodium ovale
- b. Plasmodium falciparum
- c. Plasmodium malariae
- d. Plasmodium vivax
- e. All of the above

1188. 能引起阿米巴痢疾样症状的寄生虫是()

- scaris lumbricoides
- B. Enterobius vermicularis
- C. Balantidium coli
- D. Paragonimus
- E. Plasmodium

1189. Pick up the wrong choice :

- a. Toxoplasma gondii : Sporozoa
- b. Naegleria fowlveri : Rhizopoda
- c. Cryptosporidium : Coccidia
- d. Trichomonas hominis : Telosporia
- e. Plasmodium : Coccidia

1196. Cryptosporidium multiplies in:

- a. the lumen of small intestine
- b. the brush border of epithelial lining of small intestine
- c. the lumen of large intestine
- d. the reticuloendothelial cells
- e. the respiratory mucosal cells

1199. "Anchovy sauce pus" is formed in

- a. Intestinal Amoebic abscess
- b. Cerebral malaria

- c. Amoebic Liver abscess
- d. Algid malaria
- e. Neural cysticercosis

1212.All the structures are present in trophozoite of *Entamoeba histolytica* except –

- a. Glycogen mass
- b. Chromatoid bars
- c. Eccentric nucleus
- d. Karyosome
- e. Cytostome

1227.Pickup the wrong choice :

- a. *Toxoplasma gondii* : Sporozoa
- b. *Naegleria fowleri* : Rhizopoda
- c. *Cryptosporidium* : Coccidia
- d. *Trichomonas hominis* : Telosporia
- e. *Plasmodium* : Coccidia

1253.Which is not a vector born disease

- a. Toxoplasmosis
- b. Kala azar
- c. Filariasis
- d. Chagas disease
- e. African sleeping sickness

1262.Butterfly erythema appears in

- a. African sleeping sickness
- b. Oriental sore
- c. Dermal Leishmanoid
- d. Chagas disease
- e. Katayama disease

1271.Enteric and Exoenteric life cycle is the characteristic feature of

- a. *Leishmania donovani*
- b. *Toxoplasma gondii*
- c. *Cryptosporidium*
- d. *Pneumocystis carinii*
- e. *Balantidium coli*

1289.A stained blood smear reveals giant platelets and many reticulocytes containing Schuffner's dots, large ring forms, and schizonts with 12 to 24 merozoites. You would name this as

- a. *Plasmodium vivax*
- b. *Plasmodium malariae*
- c. *Plasmodium falciparum*
- d. *Plasmodium ovale*
- e. *Babesia* spp

1293.Band shaped trophozoites are seen in

- a. *Plasmodium ovale*
- b. *Plasmodium malariae*
- c. *Plasmodium vivax*
- d. *Plasmodium falciparum*
- e. *Plasmodium knowlesi*

1296.True regarding *Trichomonas vaginalis* infection:

- a. is caused by a protozoan flagellate
- b. the infective stage is the trophozoite
- c. causes vaginal discharge in females
- d. may infect males
- e. all of the above

1312.The metacyclic forms of *Trypanosoma cruzi* are found in:

- a. Reticuloendothelial cells

- b. Heart muscles
- c. Mouth part of insect vector
- d. Hind gut of infected insect
- e. None of the above

1324. Select the true statement regarding Leishmaniasis :

- a. Amastigotes develops only inside RBC's
- b. Promastigote form in bone marrow smear is diagnostic of Kala-azar
- c. *L. tropica* and *L. donovani* can be distinguished morphologically
- d. Positive aldehyde test is due to high albumin in serum
- e. Kala-azar is present in eastern parts of India

1329. Abdominal pain, fat malabsorption and frothy stools suggests –

- a. Intestinal amoebiasis
- b. Extra intestinal amoebiasis
- c. Bacillary dysentery
- d. Acute Giardiasis
- e. Katayama disease

1335. Amoebic liver abscess can be diagnosed by demonstrating –

- a. Cysts in sterile pus
- b. Trophozoites in pus
- c. Cysts in the intestine
- d. Trophozoites in the faeces
- e. All of the above

1345. 经治疗后宿主可获得终身免疫的寄生虫感染是

- A. Malaria
- B. Amoebic liver abscess
- C. Visceral leishmaniasis
- D. Giardiasis
- E. Trichomoniasis

1347. 1. Cyst phase does not exist in –

- a. *Entamoeba histolytica*
- b. *Entamoeba coli*
- c. *Giardia intestinalis*
- d. *Trichomonas vaginalis*
- e. *Toxoplasma gondii*

1356. Contamination of contact lenses could lead to eye infection with :

- a. *Acanthamoeba*
- b. *Entamoeba coli*
- c. *Onchocerca volvulus*
- d. *Toxocara canis*
- e. Loa Loa

1365. Nephrotic syndrome in *Plasmodium malariae* is due to :

- a. deposition of immune-complexes in glomerular capillaries
- b. the severe anemia
- c. hypoproteinemia
- d. release of malarial pigments
- e. All of the above

1369. *Entamoeba dispar* Brumpt 是: ()

- A. 一种寄生人体的条件致病内阿米巴
- B. 一种非致病的人体内阿米巴
- C. 与溶组织内阿米巴有同等致病力的内阿米巴种
- D. 一种致病的尚未确定的内阿米巴种
- E. 一种目前尚无法认定的阿米巴种

1372. Mucocutaneous Leishmaniasis is caused by –

- a. *Leishmania braziliensis*
- b. *Leishmania tropica*
- c. *Leishmania donovani*
- d. a & b

e. None of the above

1374. 10 year old boy develops chronic abdominal pain, low grade fever, intermittent diarrhoea, malabsorption and oocysts demonstrated in stool repeatedly during diarrhoeal symptoms for more than two years. Most likely cause of his diarrhea is –

- a. *Entamoeba histolytica*
- b. *Balantidium coli*
- c. *Giardia lamblia*
- d. *Pneumocystis carinii*
- e. *Cryptosporidium parvum*

1394. All of the following amoebae live in the large intestine except –

- a. *Entamoeba coli*
- b. *Endolimax nana*
- c. *Entamoeba gingivalis*
- d. *Iodamoeba buschii*
- e. *Entamoeba histolytica*

1404. Select the true statement regarding toxoplasmosis in humans:

- a. Becomes active in immuno-compromised patients
- b. Acute infection could be transmitted from mother to fetus
- c. Is mainly diagnosed by serological tests
- d. Is transmitted by ingestion of cysts in undercooked meat
- e. All of the above

1428. The metacyclic forms of *Trypanosoma cruzi* are found in:

- a. Reticuloendothelial cells
- b. Heart muscles
- c. Mouth part of insect vector
- d. Hind gut of infected insect
- e. None of the above

1431. The Amoebae which are morphologically more related to *Entamoeba histolytica* is

- a. *Entamoeba dispar*
- b. *Entamoeba coli*
- c. *Dientamoeba fragilis*
- d. *Entamoeba gingivalis*
- e. *Iodamoeba buskii*

1434. In Kala-azar there's no -----

- a. Hydrocephalus
- b. thrombocytopenia
- c. hemolysis of RBC's
- d. Leucopenia
- e. Anemia

1459. 可引起肝脓肿的原虫

- A. *Plasmodium vivax*
- B. *Leishmania donovani*
- C. *Toxoplasma gondii*
- D. *Giardia lamblia*
- Entamoeba histolytica*

1462. A 24 year old patient presented with features of acute meningoencephalitis in causality, his CSF on wet mount showed motile unicellular microorganism. The most likely cause is –

- a. *Acanthamoeba castellanii*
- b. *Entamoeba histolytica*
- c. *Naegleria fowleri*
- d. *Trypanosoma cruzi*
- e. None of the above

1463. Amoebic liver abscess can be diagnosed by –

- a. Cysts in the sterile pus
- b. Trophozoites in the pus
- c. Cysts in the intestine
- d. Trophozoites in the faeces
- e. All of the above

1472.虫体有两个核和一个吸器的原虫是

- A.Dientamoeba fragilis
- B.Iodamoeba butschlii
- C.Giardia lamblia
- D.Balantidium coli
- E.Nidolimax nana

1474.Which among these are not considered as opportunistic parasite in AIDS

- a. Pneumocystis carinii
- b. Toxoplasma gondii
- c. Trichomonas vaginalis
- d. Cryptosporidium parvum
- e. None of the above

1475.'Ring haemorrhages' are characteristic feature of –

- a. Cerebral malaria
- b. Algid malaria
- c. Septicaemic malaria
- d. Neurocysticercosis
- e. Hepatic amoebiasis

1489.Which staining method is not used for malarial parasites in blood smear

- a. Leishman's staining
- b. J.S.B's staining
- c. Field's staining
- d. Giemsa's staining
- e. Albert's staining

1496.下列哪种原虫不是寄生在细胞内： ()

- A.Plasmodium malariae
- B.Pneumocystis carinii
- C.Leishmania donovani
- D.Trichomonas vaginalis
- E.Toxoplasma gondii

1506.Toxoplasma gondii is transmitted to human by:

- a. ingestion of soil contaminated by oocyst from cats
- b. congenital transmission from mother with acute toxoplasmosis
- c. organ transplant and blood transfusion (infected donor)
- d. ingestion of cysts in undercooked meat
- e. all of the above

1508.Malaria attacks (paroxysms) are caused by:

- a. release of toxins from RBC
- b. invasion of RBC by merozoites
- c. severe hemolytic anemia
- d. destruction of liver parenchyma
- e. all of the above

1514.Which protozoan infection is not transmitted through contaminated water

- a. Cryptosporidiosis
- b. Amoebiasis
- c. PCP
- d. PAM
- e. Balantidiasis

1518.Which among these are not considered as opportunistic parasite in AIDS

- a. Pneumocystis

- b. *Toxoplasma gondii*
- c. *Trichomonas vaginalis*
- d. *Cryptosporidium parvum*
- e. None of the above

1527. One of the following is a feature of *Falciparum malaria* –

- a. All stages of schizogony seen in blood
- b. Multiple infection of erythrocytes
- c. Size of parasitised RBC increases
- d. Relapses seen in infection
- e. Erythrocytic cycle complete in 72 hours

1532. *Acanthamoeba keratitis* is caused by

- a. *A. culbertsoni*
- b. *A. castellanii*
- c. *A. polyphaga*
- d. *A. hatchetti*
- e. *A. rhysodes*

1540. 机会致病原虫

- A. *Plasmodium*
- B. *Leishmania*
- C. *Naegleria*
- D. *Acanthamoeba*
- E. *Toxoplasma*

1543. Mature trophozoites and young schizonts in blood smear show a distinct tendency toward band formation and the red blood cells are not enlarged.

- a. *Plasmodium vivax*
- b. *Plasmodium falciparum*
- c. *Plasmodium malariae*

- d. *Plasmodium ovale*
- e. *Plasmodium knowlesi*

1555. NNN Media is used for the culture of

- a. *Entamoeba histolytica*
- b. *Leishmania donovani*
- c. *Toxoplasma gondii*
- d. *Cryptosporidium parvum*
- e. *Acanthamoeba culbertsoni*

1556. 寄生宿主和寄生部位广泛的原虫是

- A. *Entamoeba histolytica*
- B. *Plasmodium vivax*
- C. *Leishmania donovani*
- D. *Trichomonas vaginalis*
- E. *Toxoplasma gondii*

1561. True about *Cryptosporidium parvum* –

- a. Affect only immunocompromised patient
- b. One among Common opportunistic infection in AIDS
- c. Cysts are passed in sputum and bronchial secretions
- d. Treatment is metronidazole
- e. All of the above

1563. Frenkel's skin test is positive in –

- a. *Neurocysticercosis*
- b. *Toxoplasmosis*
- c. Hydatid cyst disease
- d. Occult filariasis
- e. *Trichomoniasis*

1568. The most distinctive feature of pathogenic *Entamoeba histolytica* on fresh stool examination is –

- a. Presence of active pseudopodia
- b. Erythrophagocytosis
- c. Presence of intracytoplasmic vacuoles
- d. Presence of two karyosomes
- e. All of the above

1572. Presence of two nuclei are the characteristic feature of

- a. *Pneumocystis carinii*
- b. *Cryptosporidium parvum*
- c. Amastigote of *Leishmania*
- d. *Balantidium coli*
- e. Trophozoites of *Trichomonas vaginalis*

1574. Regarding cutaneous amoebiasis which is not true –

- a. Is a spreading necrotizing inflammation of skin and subcutaneous tissue
- b. Can occur in the perianal region
- c. Improvement with anti-amoebic treatment occurs
- d. Infection acquires through haematogenous route
- e. None of the above

1575. *Plasmodium falciparum* causes most serious form of infection due to:

- a. absence of true relapse
- b. irregular paroxysms
- c. adherence and blocking of capillaries of internal organs
- d. multiple infections inside the RBC's
- e. all of the above

1586. A protozoan with characteristic jerky motility is most commonly observed in which of the following

- a. Biopsied muscle
- b. Blood
- c. Duodenal aspirate
- d. Sputum

e. Vaginal secretions

1591. 下列寄生虫中属于鞭毛虫的是()

- A. *Taenia saginata*
- B. *Trichomonas vaginalis*
- C. *Cryptosporidium*
- D. *Naegleria fowleri*
- E. *Trichuris trichiura*