


1
point

1.

Suppose you have trained an SVM classifier with a Gaussian kernel, and it learned the following decision boundary on the training set:



The figure shows a decision boundary that is underfit to the training set, so we'd like to lower the bias / increase the variance of the SVM. We can do so by either increasing the parameter C or decreasing σ^2 .

high bias

You suspect that the SVM is underfitting your dataset. Should you try increasing or decreasing C ? Increasing or decreasing σ^2 ?

- ☐ It would be reasonable to try **decreasing** C . It would also be reasonable to try **increasing** σ^2 .
- ☒ It would be reasonable to try **increasing** C . It would also be reasonable to try **decreasing** σ^2 .
- ☐ It would be reasonable to try **decreasing** C . It would also be reasonable to try **decreasing** σ^2 .



It would be reasonable to try **increasing** C . It would also be reasonable to try **increasing** σ^2 .


1
point

2.

The formula for the Gaussian kernel is given by

$$\text{similarity}(x, l^{(1)}) = \exp\left(-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}\right).$$

The figure below shows a plot of $f_1 = \text{similarity}(x, l^{(1)})$ when $\sigma^2 = 1$.



Which of the following is a plot of f_1 when $\sigma^2 = 0.25$?






Figure 3.

○




Figure 2.

○





Figure 4.

This figure shows a "narrower" Gaussian kernel centered at the same location which is the effect of decreasing σ^2 .



1
point

3.


The SVM solves

$$\min_{\theta} C \sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x^{(i)}) + (1 - y^{(i)}) \text{cost}_0(\theta^T x^{(i)}) + \sum_{j=1}^n \theta_j^2$$

where the functions $\text{cost}_0(z)$ and $\text{cost}_1(z)$ look like this:

Support Vector Machines

Quiz, 5 questions



The first term in the objective is:

$$C \sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x^{(i)}) + (1 - y^{(i)}) \text{cost}_0(\theta^T x^{(i)}).$$

This first term will be zero if two of the following four conditions hold true. Which are the two conditions that would guarantee that this term equals zero?

- ☐ For every example with $y^{(i)} = 0$, we have that $\theta^T x^{(i)} \leq -1$.
- ☐ For every example with $y^{(i)} = 1$, we have that $\theta^T x^{(i)} \geq 1$.
- ☐ For every example with $y^{(i)} = 1$, we have that $\theta^T x^{(i)} \geq 0$.
- ☐ For every example with $y^{(i)} = 0$, we have that $\theta^T x^{(i)} \leq 0$.

1
point

4.

Suppose you have a dataset with $n = 10$ features and $m = 5000$ examples.

After training your logistic regression classifier with gradient descent, you find that it has underfit the training set and does not achieve the desired performance on the training or cross validation sets.

Which of the following might be promising steps to take? Check all that apply.

- ☐ Increase the regularization parameter λ .





Use an SVM with a linear kernel, without introducing new features.



Create / add new polynomial features.



Use an SVM with a Gaussian Kernel.

A neural network with many hidden units is a more complex (higher variance) model than logistic regression, so it is less likely to

or: Try using a neural network with a large number of hidden units.

greater complexity and can avoid underfitting the data.

Support Vector Machines

Quiz, 5 questions

1 point

5.

Which of the following statements are true? Check all that apply.



If the data are linearly separable, an SVM using a linear kernel will return the same parameters θ regardless of the chosen value of C (i.e., the resulting value of θ does not depend on C).

A linearly separable dataset can usually be separated by many different lines. Varying the parameter C will cause the SVM's decision boundary to vary among these possibilities. For example, for a very large value of C , it might learn larger values of θ in order to increase the margin on certain examples.



Suppose you are using SVMs to do multi-class classification and would like to use the one-vs-all approach. If you have K different classes, you will train $K - 1$ different SVMs.



The maximum value of the Gaussian kernel (i.e., $\text{sim}(x, l^{(1)})$) is 1.



It is important to perform feature normalization before using the Gaussian kernel.

The similarity measure used by the Gaussian kernel expects that the data lie in approximately the same range.



I, **Jun-Chieh Wang**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account. Learn more about Coursera's Honor Code

Submit Quiz

