


1
point



1.



Consider the following 2D dataset:



Which of the following figures correspond to possible values that PCA may return for $u^{(1)}$ (the first eigenvector / first principal component)? Check all that apply (you may have to check more than one figure).







1
point

2.

Which of the following is a reasonable way to select the number of principal components k ?

(Recall that n is the dimensionality of the input data and m is the number of input examples.)

- ☐ Use the elbow method.
- ☐ Choose k to be the largest value so that at least 99% of the variance is retained
- ☒ Choose k to be the smallest value so that at least 99% of the variance is retained.
- ☐ Choose k to be 99% of m (i.e., $k = 0.99 * m$, rounded to the nearest integer).

Principal Component Analysis

Quiz, 5 questions

1
point

3.

Suppose someone tells you that they ran PCA in such a way that "95% of the variance was retained." What is an equivalent statement to this?

- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \geq 0.05$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2} \geq 0.95$
- ☐ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \geq 0.95$
- ☒ $\frac{\frac{1}{m} \sum_{i=1}^m \|x^{(i)} - x_{\text{approx}}\|^2}{\frac{1}{m} \sum_{i=1}^m \|x^{(i)}\|^2} \leq 0.05$

1
point

4.

Which of the following statements are true? Check all that apply.



Feature scaling is not useful for PCA, since the eigenvector calculation (such as using Octave's `svd(Sigma)` routine) takes care of this automatically.

- ☐ PCA can be used only to reduce the dimensionality of data by 1 (such as 3D to 2D, or 2D to 1D).
- ☐ Given an input $x \in \mathbb{R}^n$, PCA compresses it to a lower-dimensional vector $z \in \mathbb{R}^k$.
- ☐ If the input features are on very different scales, it is a good idea to perform feature scaling before applying PCA.

Principal Component Analysis

Quiz, 5 questions

point

5.

Which of the following are recommended applications of PCA? Select all that apply.

- ☐ Data compression: Reduce the dimension of your data, so that it takes up less memory / disk space.
- ☐ To get more features to feed into a learning algorithm.
- ☐ Data visualization: Reduce data to 2D (or 3D) so that it can be plotted.
- ☐ Preventing overfitting: Reduce the number of features (in a supervised learning problem), so that there are fewer parameters to learn.



I, **Jun-Chieh Wang**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account. Learn more about Coursera's Honor Code

Submit Quiz

