

To further validate generalization performance, we have additionally collected two other publicly available SAR datasets (Ottawa and Yellow River II) and conducted supplementary experiments. The results show that on the Ottawa dataset, our CAFNet achieves competitive performance in terms of F1-score, IoU, and KC, comparable to the best-performing M2HG method. On the Yellow River II dataset, CAFNet attains the highest F1-score, IoU, and KC among all compared methods. These supplementary results further confirm that our method maintains robust performance across different imaging conditions and change patterns.




Fig. 1. Experimental results from Ottawa and Yellow River II. (a) Image I_1 . (b) Image I_2 . (c) Ground truth. (d) PGAG. (e) M2HG. (f) CAMixer. (g) TSPLR. (h) FFITN. (i) NDSNet. (j) WBANet. (k) CAFNet. White:TP, black: TN, green: FN, and red: FP.

TABLE I
CHANGE DETECTION RESULTS OF OTTAWA AND YELLOW RIVER II

Method	Result on the Ottawa dataset				
	Precision	Recall	F1	IoU	KC
PGAG	97.47	86.25	91.52	84.37	90.04
M2HG	98.56	92.57	95.47	91.34	94.65
CAMixer	95.00	93.05	94.02	88.71	92.91
TSPLR	95.97	92.54	94.22	89.07	93.16
FFITN	92.47	96.14	94.27	89.16	93.17
NDSNet	94.28	93.40	93.83	88.38	92.68
WBANet	95.32	93.73	94.52	89.61	93.50
CAFNet	96.19	94.73	95.46	91.31	94.61

Method	Result on the Yellow River II dataset				
	Precision	Recall	F1	IoU	KC
PGAG	76.92	80.78	78.80	65.02	78.06
M2HG	77.09	75.68	76.38	61.78	75.58
CAMixer	72.66	85.45	78.54	64.66	77.75
TSPLR	67.45	90.39	77.25	62.94	76.36
FFITN	73.27	80.99	76.94	62.52	76.11
NDSNet	69.10	84.91	76.19	61.54	75.29
WBANet	68.06	88.70	77.02	62.63	76.13
CAFNet	75.55	87.00	80.87	67.89	80.17