TSHR and IGF1R tissue extension

A Instrument availability

GTEx v8 tissue	*TSHR*	*IGF1R*
Thyroid	X	**1**
Cultured fibroblasts	−	**1**
Subcutaneous adipose	**1**	X
Visceral adipose	**1**	X
Whole blood	X	X

1 One qualifying instrument

x No eQTL meeting P < 5e−8

− Gene absent from the tested−gene file

B Primary comparison within the tissue extension

Disease dataset: FinnGen Graves disease

Expression instrument	OR (95% CI)	P value	Adjusted P
TSHR Subcutaneous adipose	0.36 (0.31−0.43)	**1.28e−34**	**1.28e−33***
TSHR Visceral adipose	0.50 (0.43−0.58)	**4.09e−19**	**4.09e−18***
IGF1R Thyroid	0.76 (0.56−1.02)	0.0692	0.692
IGF1R Cultured fibroblasts	1.11 (0.89−1.38)	0.359	1.000



* Corrected support: raw P < 0.005; adjusted P = min(1, 10 x P). Filled squares mark corrected support.

Ten primary combinations were planned; six were unestimable. All 40 secondary comparisons are reported separately.

GTEx blood differs from eQTLGen. ORs are per normalized expression unit. Non−orbital tissues; no tissue−specific colocalization.