Tissue−specific expression associations across five disease datasets

A TSHR

Disease dataset	OR (95% CI)	P value
Subcutaneous adipose		
FinnGen GD	0.36 (0.31−0.43)	**1.28e−34***
BBJ GD	0.46 (0.38−0.55)	**1.92e−15***
UKB hyperthyroidism	0.31 (0.27−0.37)	**2.86e−42***
FinnGen ophthalmopathy	0.45 (0.32−0.63)	**5.76e−06***
FinnGen strict ophthalmopathy	0.51 (0.35−0.74)	**3.67e−04***
Visceral adipose		
FinnGen GD	0.50 (0.43−0.58)	**4.09e−19***
BBJ GD	0.56 (0.46−0.67)	**2.64e−10***
UKB hyperthyroidism	0.39 (0.34−0.45)	**9.19e−38***
FinnGen ophthalmopathy	0.69 (0.49−0.98)	0.0368
FinnGen strict ophthalmopathy	0.78 (0.54−1.13)	0.1863



B IGF1R

Disease dataset	OR (95% CI)	P value
Thyroid		
FinnGen GD	0.76 (0.56−1.02)	0.0692
BBJ GD	1.23 (0.90−1.68)	0.1996
UKB hyperthyroidism	1.33 (0.99−1.80)	0.0594
FinnGen ophthalmopathy	1.02 (0.54−1.94)	0.9449
FinnGen strict ophthalmopathy	1.23 (0.62−2.44)	0.5493
Cultured fibroblasts		
FinnGen GD	1.11 (0.89−1.38)	0.3591
BBJ GD	0.91 (0.70−1.20)	0.5111
UKB hyperthyroidism	1.03 (0.82−1.29)	0.8147
FinnGen ophthalmopathy	0.78 (0.49−1.25)	0.3005
FinnGen strict ophthalmopathy	0.78 (0.47−1.30)	0.3413



Each estimate uses one cis−eQTL. ORs are per unit of normalized, inverse−normal−transformed expression.

* Family−corrected: FinnGen GD P < 0.005; other datasets P < 0.00125. Unmarked estimates do not pass these thresholds.

FinnGen endpoints overlap. Adipose and cultured fibroblasts are not orbital tissue. Thirty unestimable planned comparisons are tabulated separately.