Table 1. Data sources and their roles

Dataset / endpoint	Cases	Controls	Role
eQTLGen blood cis-eQTL	—	—	Exposure; up to 31,684 participants (median 28,092 per variant); predominantly European
BBJ GD (GCST90018627)	2,809	172,656	Discovery; East Asian
FinnGen GD (E4_GRAVES_STRICT)	3,962	496,386	Post hoc same-disease comparison; Finnish
UKB hyperthyroidism (GCST90038636)	3,731	480,867	Broader-phenotype comparison; self-reported hyperthyroidism or thyrotoxicosis; ancestry not established as exclusively European
FinnGen GO (GRAVES_OPHT)	858	499,490	Original ophthalmopathy (TED-enriched) comparison; Finnish
FinnGen GO strict (E4_GRAVES_OPHT_STRICT)	753	499,595	Post hoc ophthalmopathy definition sensitivity; Finnish
GTEx v8 tissue eQTL	—	—	Post hoc tissue extension (thyroid, cultured fibroblasts, subcutaneous and visceral adipose, whole blood; n = 469–670; Table S4)
1000 Genomes Phase 3 European reference (n = 503)	—	—	LD clumping of instruments and reference allele frequencies

The three FinnGen R12 endpoints share participants and are not independent replications. FinnGen ophthalmopathy controls are all other participants, including people with GD but no recorded eye disease, so GO estimates do not estimate TED risk among patients with GD. UKB hyperthyroidism is broader than GD. ‘Strict’ is the FinnGen endpoint label, not proof of perfect specificity. The descriptive orbital RNA-sequencing series (four TED patients, one control) is separate from these datasets. eQTL, expression quantitative trait locus; GTEx, Genotype-Tissue Expression; LD, linkage disequilibrium; TED, thyroid eye disease; BBJ, Biobank Japan; FinnGen GD, E4_GRAVES_STRICT; GD, Graves disease; GO, Graves ophthalmopathy; UKB, UK Biobank.