Table S3. *TSHR* regional colocalization and prior sensitivity

Dataset	Matched variants	Disease lead (*P*)	rs179252 *P*	rs179252-G OR (95% CI)	H4 top variant	PP.H4, 10^{-5} / 10^{-6}	PP.H4 range across windows
BBJ GD	4,336	rs28414437 (1.83×10^{-15})	1.09×10^{-14}	0.80 (0.76–0.85)	rs179252	**0.951** / 0.661	0.951–0.951
FinnGen GD	6,579	rs1023586 (1.58×10^{-36})	3.19×10^{-32}	0.77 (0.73–0.80)	rs1023586	0.264 / 0.035	0.264–0.264
UKB hyperthyroidism	5,667	rs4903961 (1.17×10^{-43})	8.77×10^{-28}	0.77 (0.74–0.81)	rs1023586	0.226 / 0.028	0.226–0.226
FinnGen GO	6,579	rs3783947 (1.23×10^{-7})	2.82×10^{-7}	0.78 (0.71–0.86)	rs179252	**0.986** / **0.875**	0.986–0.987
FinnGen GO strict	6,579	rs2110695 (1.34×10^{-5})	9.02×10^{-5}	0.82 (0.74–0.90)	rs179252	**0.891** / 0.451	0.891–0.905

Disease lead, the strongest disease-associated variant among the matched cis variants, with its *P* value; rs179252 *P*, z-test *P* of the aligned G-allele disease association (the UKB source BOLT-LMM *P* was 1.3×10^{-27}). H4 top variant, the variant with the highest posterior under the shared-variant hypothesis; it is not necessarily the disease lead. PP.H4 under p12 = 10^{-5} and 10^{-6} (p1 = p2 = 10^{-4}); bold, ≥0.80. Window range covers the full source cis-region, ±500 kb, ±250 kb and the five-outcome intersection at p12 = 10^{-5}; it is not a confidence interval. All 240 window settings are in TSHR_windows. BBJ pairs European eQTL with East Asian disease statistics. CI, confidence interval; OR, odds ratio; PP.H4, shared-variant posterior probability; BBJ, Biobank Japan; FinnGen GD, E4_GRAVES_STRICT; GD, Graves disease; GO, Graves ophthalmopathy; UKB, UK Biobank.