Table S3. Tissue-specific expression and FinnGen GD

Gene / tissue	N	IVs	OR (95% CI)	*P* value	Adjusted *P*
TSHR / Thyroid	574	0	No qualifying IV	—	—
IGF1R / Thyroid	574	1	0.76 (0.56–1.02)	0.0692	0.692
TSHR / Cultured fibroblasts	483	0	Not tested	—	—
IGF1R / Cultured fibroblasts	483	1	1.11 (0.89–1.38)	0.359	1.000
TSHR / Subcutaneous adipose	581	1	0.36 (0.31–0.43)	1.28×10^{-34}	1.28×10^{-33}
IGF1R / Subcutaneous adipose	581	0	No qualifying IV	—	—
TSHR / Visceral adipose	469	1	0.50 (0.43–0.58)	4.09×10^{-19}	4.09×10^{-18}
IGF1R / Visceral adipose	469	0	No qualifying IV	—	—
TSHR / Whole blood	670	0	No qualifying IV	—	—
IGF1R / Whole blood	670	0	No qualifying IV	—	—

Fixed-panel tissue extension with FinnGen GD as the primary disease dataset. IVs are retained instrumental variants. ORs are per unit of normalized, inverse-normal-transformed expression. Adjusted $P = \min(1, 10P)$; the ten-comparison denominator includes unestimable cells. Not tested means the gene was absent from the tissue eGenes file; no qualifying IV means the minimum eQTL *P* did not meet 5×10^{-8}. All four estimable combinations used one SNP (F 32.1–61.1), so each *P* value equals that variant's disease-association *P* value. The adipose *TSHR* instruments (rs12101255 and rs74064796, 50 bp apart in *TSHR* intron 1) are themselves blood *TSHR* eQTLs near rs179252 and therefore re-express the regional *TSHR* association rather than providing independent evidence. The 40 secondary comparisons use a separate denominator of 40 and are retained, with all source statistics, in Tissue_extension. Cultured fibroblasts and adipose tissues are not orbital tissue. Original detection benchmarks remain in Detection_original.