



UPPSALA  
UNIVERSITET



# Dynamic social networks in dairy cows

## Summary

From the positional data of cows in a free-stall barn in the Netherlands we obtain the adjacency matrices, social networks and communities of dairy cows. Then monadic, dyadic and polyadic features of social networks are analysed and combined to investigate the dynamics of the dairy cows' social networks.

## Methods

The positional data was converted into adjacency matrices, which were then converted to social networks. Python is suitable for analyzing social network, since there are a lot of pre-existing functions to compute metrics in it.

	cow1	cow2	...	...	cowN
cow1	0				
cow2		0			
...			0		
...				0	
cowN					0

**Figure 1: The structure of adjacency matrix, the adjacency matrix is symmetric, elements represent the relationship between two cows**


We looked at community detection, where we used the following algorithms: Girvan-Newman, Louvain, and Clique Percolation Method (CPM).

We also looked at some monadic metrics: centrality, edge distribution, and eigenvector centrality. One dyadic metrics: average shortest path, and for the polyadic metrics we looked at transitivity, eigenvalues, and laplacian eigenvalues. Features of graphs and communities are also considered as part of polyadic metrics.

We also compared our results with simulated data to see how statistically significant our results were.


## Results

The monadic metrics showed that there are some clear, significant distribution of how many interactions cows had. There are some cows that have a large number of interactions, and some have almost no interactions.



**Figure 2: The Centrality for all cows in the barn, higher centrality means more social contact**

The dyadic metric shows that the social network is well connected as two cows are on average less than two connections away from each other.



**Figure 3: The Normalized Mutual Information (NMI) for partitions of different methods over 14 days, low NMI means that the partitions of dairy cows' communities change tremendously**

The polyadic metrics show that cows do not follow the saying a friend of a friend is my friend. The chance that two of a cow's connecting cows are also connected is lower than in the randomly simulated networks.

Especially, the features of communities show that the communities of cows on different days are relatively independent. They are more stable than random communities but much less than we expected them to be.



Photo Credit: iStock

## Conclusion

- There is a consistency in how cows interact, but only when looking at monadic or dyadic interactions.
- The dairy cows' communities are more stable than the random communities. But in general, the communities of different days are still very independent. There is no preference for group forming, and cows do not have stable social communities like humans.
- The social structure is more dependent on the individuals (some very active cows) standing in the group than on any type of social group or community.

Authors:

Emil Helmryd Grosfilley  
emil.helmrydgrosfilley.163  
2@student.uu.se

Junjie Chu  
allen7chu@foxmail.com

Dap De Bruijkere  
dap.debruijkere.2546@stu  
dent.uu.se

Supervisors:

Lars Rönnegård (SLU)

Moudud Alam (SLU)