

Statistical Computing

Constrained Optimization & Duality

Spring 2026

Constrained Optimization

Equality Constrained Optimization


Inequality Constrained Optimization

Lagrangian Duality

Constrained Optimization

Constrained Maxima and Minima

- In many practical optimization problems, we must maximize or minimize a function in which the independent variables are subjected to certain further constraints.



Equality Constrained Optimization

Substitution Method


- Find the relative minimum of $f(x, y) = 2x^2 + y^2$ subject to $g(x, y) = x + y - 1 = 0$.

Solution.

Solving the constraint for y gives $y = -x + 1$. Substituting into f :

$$h(x) = 2x^2 + (1 - x)^2 = 3x^2 - 2x + 1.$$

Solve $h'(x) = 6x - 2 = 0 \Rightarrow x = \frac{1}{3}$. Since $h''(x) = 6 > 0$, this is a relative minimum. With $y = \frac{2}{3}$, the constrained minimum is at $(\frac{1}{3}, \frac{2}{3})$.



The Method of Lagrange Multipliers

- To find the relative extrema of $f(x, y)$ subject to $g(x, y) = 0$:

1. Form the **Lagrangian function** (where λ is the **Lagrange multiplier**):

$$F(x, y, \lambda) = f(x, y) + \lambda g(x, y).$$

2. Solve the system of equations

$$F_x = 0, \quad F_y = 0, \quad F_\lambda = 0$$

for all values of x , y , and λ .

3. The solutions found in step 2 are candidates for the extrema of f .

Example: Lagrange Multipliers

- Find the relative minimum of $f(x, y) = 2x^2 + y^2$ subject to $x + y = 1$ using Lagrange multipliers.

Solution.

- Write $g(x, y) = x + y - 1 = 0$. Form the Lagrangian:

$$F(x, y, \lambda) = (2x^2 + y^2) + \lambda(x + y - 1).$$


- Solve the system:

$$F_x = 4x + \lambda = 0, \quad F_y = 2y + \lambda = 0, \quad F_\lambda = x + y - 1 = 0.$$

- From the first two equations: $x = -\frac{\lambda}{4}$ and $y = -\frac{\lambda}{2}$.
- Substituting into the third: $\lambda = -\frac{4}{3}$, so $x = \frac{1}{3}$, $y = \frac{2}{3}$.

Geometric Interpretation

$$\begin{array}{ll}\text{Minimize} & 2x_1^2 + x_2^2 \\ \text{Subject to} & x_1 + x_2 = 1\end{array}$$



Geometric Interpretation

- Consider the gradients of f and h at the optimal point $\mathbf{x}^*$.
- They must be parallel (same direction, possibly different lengths):

$$\nabla f(\mathbf{x}^*) = \nu^* \nabla h(\mathbf{x}^*).$$

- This is equivalent to $\nabla L(\mathbf{x}^*, \nu^*) = 0$ along with feasibility of $\mathbf{x}^*$.
- $\nabla f(\mathbf{x})$ is perpendicular to a level curve of f ; similarly $\nabla h(\mathbf{x})$ is perpendicular to the level curve $h(\mathbf{x}) = 0$.
- From the example, at $x_1^* = \frac{1}{3}$, $x_2^* = \frac{2}{3}$, $\lambda_1^* = \frac{4}{3}$:

$$\nabla f(x_1^*, x_2^*) = \begin{bmatrix} \frac{4}{3} & \frac{4}{3} \end{bmatrix}, \quad \nabla h_1(x_1^*, x_2^*) = \begin{bmatrix} 1 & 1 \end{bmatrix}.$$

Equality Constrained Optimization

- Optimization problem with one equality constraint:

$$\begin{aligned} & \text{minimize} && f(\mathbf{x}) \\ & \text{subject to} && h(\mathbf{x}) = 0. \end{aligned}$$

- The **Lagrangian** is $L(\mathbf{x}, \nu) = f(\mathbf{x}) + \nu h(\mathbf{x})$.
- A solution must satisfy $\nabla_{\mathbf{x}} L(\mathbf{x}^*, \nu^*) = \nabla f(\mathbf{x}^*) + \nu^* \nabla h(\mathbf{x}^*) = \mathbf{0}$, i.e., there exists ν^* such that

$$\frac{\partial L(\mathbf{x}^*, \nu^*)}{\partial x_1} = \dots = \frac{\partial L(\mathbf{x}^*, \nu^*)}{\partial x_n} = 0, \quad \frac{\partial L(\mathbf{x}^*, \nu^*)}{\partial \nu} = h(\mathbf{x}^*) = 0.$$

Equality Constrained Optimization: Two Constraints

- Optimization problem with two equality constraints:

$$\begin{array}{ll}\text{minimize} & f(\mathbf{x}) \\ \text{subject to} & h_1(\mathbf{x}) = 0, \ h_2(\mathbf{x}) = 0.\end{array}$$

- The Lagrangian is $L(\mathbf{x}, \boldsymbol{\nu}) = f(\mathbf{x}) + \nu_1 h_1(\mathbf{x}) + \nu_2 h_2(\mathbf{x})$.
- Stationarity condition:

$$\nabla_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\nu}) = \nabla f(\mathbf{x}) + \nu_1 \nabla h_1(\mathbf{x}) + \nu_2 \nabla h_2(\mathbf{x}) = \mathbf{0}.$$

Equality Constrained Optimization: General Case

- General problem with m equality constraints:

$$\begin{array}{ll}\text{minimize} & f(\mathbf{x}) \\ \text{subject to} & h_j(\mathbf{x}) = 0, \quad j = 1, \dots, m.\end{array}$$

- Denote the constraint vector and multiplier vector:

$$\mathbf{h}(\mathbf{x}) = \begin{pmatrix} h_1(\mathbf{x}) \\ \vdots \\ h_m(\mathbf{x}) \end{pmatrix} \in \mathbb{R}^m, \quad \mathbf{v} = \begin{pmatrix} v_1 \\ \vdots \\ v_m \end{pmatrix} \in \mathbb{R}^m.$$

- The Lagrangian is

$$L(\mathbf{x}, \mathbf{v}) = f(\mathbf{x}) + \sum_{j=1}^m v_j h_j(\mathbf{x}) = f(\mathbf{x}) + \mathbf{v}^\top \mathbf{h}(\mathbf{x}).$$

Equality Constrained Optimization: General Case (cont.)

- **First-Order Necessary Conditions (equality constraints only):**
- Suppose $\mathbf{x}^*$ is a local minimizer and LICQ holds at $\mathbf{x}^*$.
 - **LICQ** (Linear Independence Constraint Qualification): the constraint gradients $\nabla h_1(\mathbf{x}^*), \dots, \nabla h_m(\mathbf{x}^*)$ are linearly independent.
- Then there exists $\boldsymbol{\nu}^*$ such that at $(\mathbf{x}^*, \boldsymbol{\nu}^*)$:

$$\nabla_{\mathbf{x}} L(\mathbf{x}^*, \boldsymbol{\nu}^*) = \nabla f(\mathbf{x}^*) + \sum_{j=1}^m \nu_j^* \nabla h_j(\mathbf{x}^*) = \mathbf{0},$$

$$h_j(\mathbf{x}^*) = 0, \quad j = 1, \dots, m.$$

LICQ: Why Linear Independence?

- **Example:**

$$\begin{aligned} &\text{Minimize} && x_1 + x_2 + x_3^2 \\ &\text{Subject to:} && x_1 = 1, \quad x_1^2 + x_2^2 = 1. \end{aligned}$$

The minimum is achieved at $x_1 = 1, x_2 = 0, x_3 = 0$.

The Lagrangian is:

$$L(x_1, x_2, x_3, \lambda_1, \lambda_2) = x_1 + x_2 + x_3^2 + \lambda_1(1 - x_1) + \lambda_2(1 - x_1^2 - x_2^2).$$

But $\frac{\partial L(1, 0, 0, \lambda_1, \lambda_2)}{\partial x_2} = 1$ for all λ_1, λ_2 : no multipliers can satisfy stationarity!

Observe: $\nabla h_1(1, 0, 0) = \begin{bmatrix} 1 & 0 & 0 \end{bmatrix}$ and $\nabla h_2(1, 0, 0) = \begin{bmatrix} 2 & 0 & 0 \end{bmatrix}$ are *linearly dependent*.

Inequality Constrained Optimization

- Optimization problem with one inequality constraint:

$$\begin{array}{ll}\text{minimize} & f(\mathbf{x}) \\ \text{subject to} & g(\mathbf{x}) \leq 0.\end{array}$$

- Two cases arise depending on where the constrained minimum $\mathbf{x}^*$ lies:
 - $g(\mathbf{x}^*) = 0$ (boundary): the constraint is active; the solution is pushed onto the constraint boundary.
 - $g(\mathbf{x}^*) < 0$ (interior): the constraint is inactive; the solution coincides with the unconstrained minimum and $\lambda^* = 0$.

Inequality Constrained Optimization: Active Constraint

- If the solution lies on the boundary ($g(\mathbf{x}^*) = 0$), the constraint is active and behaves like an equality constraint. The same Lagrange condition applies:

$$\nabla f(\mathbf{x}^*) + \lambda^* \nabla g(\mathbf{x}^*) = \mathbf{0}.$$




Figure 10.5. An active inequality constraint. The corresponding contour line is shown in red.




Figure 10.6. An inactive inequality constraint.

Example: Inequality Constraint

$$\begin{array}{ll}\text{Minimize} & 3x_1^2 + x_2^2 \\ \text{subject to} & x_1 + x_2 \geq 1\end{array}$$

The unconstrained minimum $(0, 0)$ violates $x_1 + x_2 \geq 1$, so the constraint is active at $\mathbf{x}^*$.

With $g(\mathbf{x}) = 1 - x_1 - x_2$, stationarity of $L = 3x_1^2 + x_2^2 + \lambda(1 - x_1 - x_2)$ gives $x_1 = \frac{\lambda}{6}$, $x_2 = \frac{\lambda}{2}$. Substituting into the constraint:

$$\frac{\lambda}{6} + \frac{\lambda}{2} = 1 \Rightarrow \lambda^* = \frac{3}{2}, \quad \mathbf{x}^* = \left(\frac{1}{4}, \frac{3}{4}\right).$$


Motivation for the KKT (Karush-Kuhn-Tucker) Conditions

- Imposing $g(\mathbf{x}) \leq 0$ is equivalent to minimizing a step-penalized objective:

$$\min_{\mathbf{x}} f(\mathbf{x}) \text{ s.t. } g(\mathbf{x}) \leq 0 \iff \min_{\mathbf{x}} f_{\infty}(\mathbf{x}),$$

where $f_{\infty}(\mathbf{x}) := f(\mathbf{x}) + \infty \cdot \mathbf{1}_{g(\mathbf{x}) > 0}(\mathbf{x})$ equals $f(\mathbf{x})$ when $g(\mathbf{x}) \leq 0$ and $+\infty$ otherwise.

- We use a linear penalty $\lambda g(\mathbf{x})$ ($\lambda \geq 0$) as a lower bound on $\infty \cdot \mathbf{1}_{g(\mathbf{x}) > 0}(\mathbf{x})$, penalizing infeasibility.



Motivation for the KKT Conditions

- Lagrangian for inequality constraints:


$$L(\mathbf{x}, \lambda) = f(\mathbf{x}) + \lambda g(\mathbf{x}).$$

- Recover f_∞ by maximizing over λ :

$$f_\infty(\mathbf{x}) = \max_{\lambda \geq 0} L(\mathbf{x}, \lambda).$$

- (Primal problem)

$$\min_{\mathbf{x}} \max_{\lambda \geq 0} L(\mathbf{x}, \lambda).$$



From $\min_{\mathbf{x}} \max_{\lambda \geq 0} L(\mathbf{x}, \lambda) := f(\mathbf{x}) + \lambda g(\mathbf{x})$, the following conditions must hold at $\mathbf{x}^*$:

- **Primal feasibility:** $g(\mathbf{x}^*) \leq 0$.
- **Dual feasibility:** $\lambda^* \geq 0$, required for $\lambda g(\mathbf{x})$ to serve as a lower bound on the step penalty.
- **Complementary slackness:** $\lambda^* g(\mathbf{x}^*) = 0$.
 - When $g(\mathbf{x}^*) < 0$, λ^* must be 0 to recover $f(\mathbf{x}^*)$ from $\max_{\lambda \geq 0} L(\mathbf{x}^*, \lambda)$.
 - When $g(\mathbf{x}^*) = 0$, λ^* may be positive.
- **Stationarity:** $\nabla_{\mathbf{x}} L(\mathbf{x}^*, \lambda^*) = \nabla f(\mathbf{x}^*) + \lambda^* \nabla g(\mathbf{x}^*) = \mathbf{0}$.
 - When the constraint is active, the level curves of f and g must be aligned.
 - When inactive, $\nabla f(\mathbf{x}^*) = \mathbf{0}$ and $\lambda^* = 0$.

KKT Conditions: General Case

- General problem with l inequality and m equality constraints:

$$\begin{aligned} & \text{minimize} && f(\mathbf{x}) \\ & \text{subject to} && g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, l, \quad h_j(\mathbf{x}) = 0, \quad j = 1, \dots, m. \end{aligned}$$

- Lagrangian:**

$$L(\mathbf{x}, \boldsymbol{\lambda}, \mathbf{v}) = f(\mathbf{x}) + \sum_i \lambda_i g_i(\mathbf{x}) + \sum_j v_j h_j(\mathbf{x}).$$

- Active set:** $\mathcal{A}(\mathbf{x}^*) = \{i : g_i(\mathbf{x}^*) = 0\} \cup \{j : h_j(\mathbf{x}^*) = 0\}$.
- LICQ** (Linear Independence Constraint Qualification): the gradients

$$\{\nabla g_i(\mathbf{x}^*) : g_i(\mathbf{x}^*) = 0\} \cup \{\nabla h_j(\mathbf{x}^*)\}$$

are linearly independent at $\mathbf{x}^*$.

- For almost every choice of differentiable objective and constraint functions, the LICQ holds at every local minimizer.

KKT Conditions: General Case (cont.)

- If $\mathbf{x}^*$ is a local minimizer and LICQ holds, there exist $(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ satisfying the **KKT conditions**:

- **Stationarity:**

$$\nabla_{\mathbf{x}} L(\mathbf{x}^*, \boldsymbol{\lambda}^*, \boldsymbol{\nu}^*) = \nabla f(\mathbf{x}^*) + \sum_i \lambda_i^* \nabla g_i(\mathbf{x}^*) + \sum_j \nu_j^* \nabla h_j(\mathbf{x}^*) = \mathbf{0}.$$

- **Primal feasibility:**

$$g_i(\mathbf{x}^*) \leq 0, \quad i = 1, \dots, l; \quad h_j(\mathbf{x}^*) = 0, \quad j = 1, \dots, m.$$

- **Dual feasibility:**

$$\lambda_i^* \geq 0, \quad i = 1, \dots, l.$$

- **Complementary slackness:**

$$\lambda_i^* g_i(\mathbf{x}^*) = 0, \quad i = 1, \dots, l.$$

Example: KKT with Inequality Constraints

$$\begin{array}{ll} \text{maximize} & -(x-2)^2 - 2(y-1)^2 \\ \text{subject to} & x+4y \leq 3, \quad x \geq y \end{array} \Rightarrow \begin{array}{ll} \text{minimize} & (x-2)^2 + 2(y-1)^2 \\ \text{subject to} & x+4y-3 \leq 0, \quad -x+y \leq 0. \end{array}$$

Lagrangian: $L = (x-2)^2 + 2(y-1)^2 + \mu_1(x+4y-3) + \mu_2(-x+y)$.

KKT stationarity:

$$\frac{\partial L}{\partial x} = 2(x-2) + \mu_1 - \mu_2 = 0, \quad \frac{\partial L}{\partial y} = 4(y-1) + 4\mu_1 + \mu_2 = 0.$$

Complementary slackness: $\mu_1(x+4y-3) = 0, \quad \mu_2(-x+y) = 0, \quad \mu_1, \mu_2 \geq 0.$

Example: KKT with Inequality Constraints (continued)

Two complementarity conditions $\Rightarrow$ check 4 cases:

- **Case 1** $\mu_1 = \mu_2 = 0$: stationarity $\Rightarrow x = 2, y = 1$. But $2 + 4(1) = 6 > 3$: *infeasible*.
- **Case 2** $\mu_1 = 0, x = y$: stationarity $\Rightarrow x = y = \frac{4}{3}$, but $\mu_2 = -\frac{4}{3} < 0$: *dual infeasible*.
- **Case 3** $x + 4y = 3, \mu_2 = 0$: stationarity $\Rightarrow x = \frac{5}{3}, y = \frac{1}{3}, \mu_1 = \frac{2}{3} > 0$.
Inactive constraint: $-x + y = -\frac{4}{3} \leq 0 \checkmark$. **Valid KKT point.**
- **Case 4** $x + 4y = 3, x = y$: stationarity $\Rightarrow x = y = \frac{3}{5}$, but $\mu_2 = -\frac{48}{25} < 0$: *dual infeasible*.

Optimal solution: $x^* = \frac{5}{3}, y^* = \frac{1}{3}, (x^* - 2)^2 + 2(y^* - 1)^2 = 1$.

Example: KKT Conditions

$$\begin{array}{ll}\text{minimize} & (x-3)^2 + (y-2)^2 \\ \text{subject to} & x^2 + y^2 \leq 5, \quad x + 2y \leq 4, \quad x, y \geq 0\end{array}$$


$$\nabla f = (2x-6, 2y-4), \quad \nabla g_1 = (2x, 2y), \quad \nabla g_2 = (1, 2), \quad \nabla g_3 = (-1, 0), \quad \nabla g_4 = (0, -1).$$

At $(x, y) = (2, 1)$: feasible with active set $\mathcal{I} = \{1, 2\}$. KKT stationarity:

$$\begin{bmatrix} 2 \\ 2 \end{bmatrix} - \mu_1 \begin{bmatrix} 4 \\ 2 \end{bmatrix} - \mu_2 \begin{bmatrix} 1 \\ 2 \end{bmatrix} = \mathbf{0} \quad \Rightarrow \quad \mu_1 = \frac{1}{3}, \quad \mu_2 = \frac{2}{3}.$$

Example: KKT with Multiple Constraints

$$\begin{aligned} &\text{minimize} && (x_1 - \tfrac{3}{2})^2 + (x_2 - \tfrac{1}{2})^2 \\ &\text{subject to} && c_1(\mathbf{x}) = x_1 + x_2 - 1 \leq 0, \\ & && c_2(\mathbf{x}) = x_1 - x_2 - 1 \leq 0, \\ & && c_3(\mathbf{x}) = -x_1 + x_2 - 1 \leq 0, \\ & && c_4(\mathbf{x}) = -x_1 - x_2 - 1 \leq 0. \end{aligned}$$



- Solution: $\mathbf{x}^* = (1, 0)^T$; active constraints: c_1, c_2 .

$$\nabla f(\mathbf{x}^*) = \begin{bmatrix} -1 \\ -\frac{1}{2} \end{bmatrix}, \quad \nabla c_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \quad \nabla c_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

- KKT multipliers: $\lambda^* = (\frac{3}{4}, \frac{1}{4}, 0, 0)^T$.

■ Class Exercise 1

$$\begin{aligned} &\text{minimize} && 2x_1^2 + 2x_1x_2 + x_2^2 - 10x_1 - 10x_2 \\ &\text{subject to} && x_1^2 + x_2^2 \leq 5, \quad 3x_1 + x_2 \leq 6. \end{aligned}$$

Solution: $x_1^* = 1$, $x_2^* = 2$, $\lambda^* = 1$.

■ Class Exercise 2

$$\begin{aligned} &\text{maximize} && \ln x + 1 + y \\ &\text{subject to} && 2x + y \leq 3, \quad x, y \geq 0. \end{aligned}$$

■ Class Exercise 3

$$\begin{aligned} &\text{minimize} && x^2 + y^2 \\ &\text{subject to} && x^2 + y^2 \leq 5, \quad x + 2y = 4, \quad x, y \geq 0. \end{aligned}$$

Lagrangian Duality

Motivation for Duality

- **Primal problem:**

$$\begin{aligned} & \text{minimize} && f(\mathbf{x}) \\ & \text{subject to} && \mathbf{g}(\mathbf{x}) \leq \mathbf{0}, \iff \min_{\mathbf{x}} \max_{\lambda \geq 0, \nu} L(\mathbf{x}, \lambda, \nu) = f(\mathbf{x}) + \lambda^\top \mathbf{g}(\mathbf{x}) + \nu^\top \mathbf{h}(\mathbf{x}). \\ & && \mathbf{h}(\mathbf{x}) = \mathbf{0} \end{aligned}$$

- **Dual problem:**

$$\max_{\lambda \geq 0, \nu} \min_{\mathbf{x}} L(\mathbf{x}, \lambda, \nu)$$

- For any (λ, ν) :

$$L(\mathbf{x}, \lambda, \nu) \leq \max_{\lambda' \geq 0, \nu'} L(\mathbf{x}, \lambda', \nu') \Rightarrow \min_{\mathbf{x}} L(\mathbf{x}, \lambda, \nu) \leq \min_{\mathbf{x}} \max_{\lambda' \geq 0, \nu'} L(\mathbf{x}, \lambda', \nu').$$

- Hence: $\max_{\lambda \geq 0, \nu} \min_{\mathbf{x}} L \leq \min_{\mathbf{x}} \max_{\lambda \geq 0, \nu} L$.
 - The solution to the dual problem provides a **lower bound** to the primal optimum.

The Lagrangian Duality

Primal problem (P)

$$\begin{aligned} & \text{minimize } f(\mathbf{x}) \\ & \text{subject to } g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, l, \\ & \quad \quad \quad h_j(\mathbf{x}) = 0, \quad j = 1, \dots, m. \end{aligned}$$

Lagrangian Dual Problem (D)

$$\begin{aligned} & \text{maximize } \mathcal{D}(\boldsymbol{\lambda}, \boldsymbol{\nu}) := \min_{\mathbf{x}} L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) \\ & \text{subject to } \lambda_i \geq 0, \quad i = 1, \dots, l, \end{aligned}$$

where $L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = f(\mathbf{x}) + \sum_i \lambda_i g_i(\mathbf{x}) + \sum_j \nu_j h_j(\mathbf{x})$.

Weak Duality Theorem

Weak Duality Theorem

Let $\mathbf{x}$ be feasible for (P) and $(\boldsymbol{\lambda}, \boldsymbol{\nu})$ be feasible for (D) (i.e., $\boldsymbol{\lambda} \geq \mathbf{0}$). Then

$$f(\mathbf{x}) \geq \mathcal{D}(\boldsymbol{\lambda}, \boldsymbol{\nu}).$$

Moreover, if $f(\mathbf{x}^*) = \mathcal{D}(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$, then $\mathbf{x}^*$ and $(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ are optimal for (P) and (D) respectively.

Proof. Since $\mathcal{D}(\boldsymbol{\lambda}, \boldsymbol{\nu}) = \inf_y L(y, \boldsymbol{\lambda}, \boldsymbol{\nu}) \leq L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu})$ and $\boldsymbol{\lambda} \geq \mathbf{0}$:

$$\mathcal{D}(\boldsymbol{\lambda}, \boldsymbol{\nu}) \leq f(\mathbf{x}) + \underbrace{\boldsymbol{\lambda}^\top \mathbf{g}(\mathbf{x})}_{\leq 0} + \underbrace{\boldsymbol{\nu}^\top \mathbf{h}(\mathbf{x})}_{=0} \leq f(\mathbf{x}). \quad \square$$

Duality Gap

If the optimal objective value of the primal problem is *strictly greater* than the optimal objective value of the dual problem, a **duality gap** is said to exist:

$$f^* - \mathcal{D}^* > 0.$$

Strong Duality Theorem

Strong Duality Theorem

Let f, g_i be convex ($i = 1, \dots, l$) and h_j be affine ($j = 1, \dots, m$). Suppose **Slater's condition** holds: $\exists \mathbf{x} \in \mathbb{R}^n$ such that $g_i(\mathbf{x}) < 0$ and $h_j(\mathbf{x}) = 0$. Then

$$\inf\{f(\mathbf{x}) : g_i(\mathbf{x}) \leq 0, h_j(\mathbf{x}) = 0\} = \sup\{\mathcal{D}(\boldsymbol{\lambda}, \boldsymbol{\nu}) : \boldsymbol{\lambda} \geq 0\}.$$

If the inf is finite, the sup is achieved at some $(\boldsymbol{\lambda}^*, \boldsymbol{\nu}^*)$ with $\boldsymbol{\lambda}^* \geq 0$. If the inf is achieved at $\mathbf{x}^*$, then $\lambda_i^* g_i(\mathbf{x}^*) = 0$ for all i .

- Under convexity and Slater's condition, the **duality gap is zero**.
- Affine: $h_j(\mathbf{x}) = \mathbf{a}_j^T \mathbf{x} + b_j$ (linear function plus constant; both convex and concave).
- Note that the Lagrangian $L(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu})$ is convex.

Example: Strong Duality


- Consider:

$$\begin{aligned} &\text{minimize } x_1^2 + x_2^2 \\ &\text{subject to } x_1 + x_2 \geq 4. \end{aligned}$$

Lagrangian: $L(\mathbf{x}, \lambda) = x_1^2 + x_2^2 + \lambda(4 - x_1 - x_2)$.

Dual function:

$$\begin{aligned} \mathcal{D}(\lambda) &= \min_{\mathbf{x}} L(\mathbf{x}, \lambda) \\ &= \min_{\mathbf{x}} \{x_1^2 + x_2^2 + \lambda(4 - x_1 - x_2)\} \\ &= \min_{\mathbf{x}} \{(x_1^2 - \lambda x_1) + (x_2^2 - \lambda x_2)\} + 4\lambda \\ &= 4\lambda + \min_{x_1} \{x_1^2 - \lambda x_1\} + \min_{x_2} \{x_2^2 - \lambda x_2\}. \end{aligned}$$



Example: Strong Duality (continued)

The minimum of $L(\mathbf{x}, \lambda)$ is attained at

$$x_1(\lambda) = \frac{\lambda}{2}, \quad x_2(\lambda) = \frac{\lambda}{2}.$$

Thus:

$$\mathcal{D}(\lambda) = L(\mathbf{x}(\lambda), \lambda) = 4\lambda - \frac{\lambda^2}{2} \quad \forall \lambda \geq 0.$$

The dual function is concave and differentiable. Maximize:

$$\frac{\partial \mathcal{D}}{\partial \lambda} = 4 - \lambda = 0 \Rightarrow \lambda^* = 4.$$

Therefore $\mathcal{D}(\lambda^*) = 8$ and $\mathbf{x}^* = (2, 2)$. No duality gap: $f(\mathbf{x}^*) = 8 = \mathcal{D}(\lambda^*)$.

✓

Exercise

- Solve the problem.
- Formulate and solve the dual problem.
- Check whether the primal and dual objective values are equal (strong duality).

$$\begin{array}{ll}\text{minimize} & x_1 \\ \text{subject to} & x_1^2 + x_2^2 = 1.\end{array}$$

- **Linear Programming (LP):**

$$\begin{aligned} & \text{minimize } c^T x \\ & \text{subject to } Ax = b, \quad x \geq 0. \end{aligned}$$

- Lagrangian: $L(x, \lambda, \nu) = c^T x - \lambda^T x + \nu^T (b - Ax)$.
- Stationarity for the dual: $c - \lambda - A^T \nu = 0, \lambda \geq 0$.
- **Dual LP:**

$$\begin{aligned} & \text{maximize } b^T \nu \\ & \text{subject to } A^T \nu \leq c. \end{aligned}$$

- **Quadratic Programming (QP):**

$$\begin{aligned} & \text{minimize } \frac{1}{2}x^T Hx + d^T x \\ & \text{subject to } Ax \leq b \end{aligned}$$

where H is symmetric and positive semidefinite.

- Lagrangian: $L(x, \lambda) = \frac{1}{2}x^T Hx + d^T x + \lambda^T (Ax - b)$.
- Stationarity: $Hx + A^T \lambda + d = 0 \Rightarrow x = -H^{-1}(d + A^T \lambda)$.
- **Dual QP:**

$$\begin{aligned} & \text{maximize } \frac{1}{2}\lambda^T D\lambda + \lambda^T c - \frac{1}{2}d^T H^{-1}d \\ & \text{subject to } \lambda \geq 0, \end{aligned}$$

where $D = -AH^{-1}A^T$ and $c = -b - AH^{-1}d$.