

This article was downloaded by: [Northwestern University]

On: 21 January 2015, At: 00:48

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Optimization: A Journal of Mathematical Programming and Operations Research

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gopt20>

Modeling of packing problems

G. Scheithauer & J. Terno

Published online: 20 Mar 2007.

To cite this article: G. Scheithauer & J. Terno (1993) Modeling of packing problems, Optimization: A Journal of Mathematical Programming and Operations Research, 28:1, 63-84, DOI: [10.1080/02331939308843904](https://doi.org/10.1080/02331939308843904)

To link to this article: <http://dx.doi.org/10.1080/02331939308843904>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

MODELING OF PACKING PROBLEMS

G. SCHEITHAUER and J. TERNO

*Technische Universität Dresden,
Inst. für Numerische Mathematik,
Mommstr. 13, O-8027 Dresden, Germany.*

(Received 2 June 1992; in final form 8 March 1993)

In this paper the modeling of packing problems (cutting and allocation problems) is investigated. Especially, linear mixed-integer models are proposed for two-dimensional packing problems of convex and non-convex polygons. The models are based on the so-called (outer) hodograph and the inner hodograph. Rotations of the polygons are not allowed. The principles of modeling in the two-dimensional case can be applied also for three- and higher dimensional packing problems.

KEY WORDS Cutting and packing problems, packing of polygons, modeling, linear mixed-integer modeling, 0/1-problems, hodograph.

Mathematics Subject Classification 1991:
Primary: 90C11; Secondary: 90C09.

1. INTRODUCTION AND PROBLEM FORMULATION

In this paper packing problems of the following type are investigated:

Let be given a closed and bounded region B with $B \subset R^n$, ($n = 1, 2, 3$) whose interior int B is connected, and m (closed and bounded) objects (pieces, figures) T_i with $T_i \subset R^n$, int T_i connected, $i = 1, \dots, m$. To each object T_i a non-negative valuation c_i is assigned. The objects may be translated but not rotated.

An allocation (or packing) of a subset of the objects within B is called feasible if the packed objects do not overlap each other.

Find a feasible allocation such that the total valuation is a maximum.

In the literature, problems of this kind are called packing problems but also allocation or cutting problems. A typology of such problems is given in [4]. Because of their practical relevance a lot of work is done with respect to modeling, exact and heuristic solutions. A wide field of such investigations is contained in [8]. Bibliographies are given in [5] and [7].

Packing problems belong in general to the NP-hard problems but in many cases problems of medium size can be solved exactly with acceptable computational amount, especially in the case of rectangular regions and objects.

In [2] a linear optimization model with 0/1-variable for the two-dimensional non-guillotine cutting problem of rectangular objects is proposed. The branch and bound algorithm given there uses a Lagrangean relaxation and a subgradient iteration to compute upper bounds.

A mixed-integer nonlinear programming model is used in [3] to describe the two-dimensional assortment problem with rectangular objects. The model contains a nonlinear objective function but only linear constraints.

The problem of deciding whether a polygon can be translated to fit inside another polygon is investigated in [1] from an algorithmical point of view.

In this paper we develop linear optimization models with continuous and 0/1-variables for packing problems with convex and non-convex polygons. The number of 0/1-variables is essentially reduced in comparison to [2] and [3]. This reduction is based on the use of very helpful tools – the so-called (outer) hodograph and the inner hodograph of two polygons. The (outer) hodograph can be used to handle the non-overlapping condition of two packed objects. The containment of a polygon in another polygon can be characterized by using the inner hodograph.

2. SET-THEORETICAL MODEL

Within the paper an orthogonal x, y, z -coordinate system is used for the description of the figures T_i and the region B . It is called reference system. Depending on the considered dimension the investigations are made with respect to the x -axis, the x, y -plane or the x, y, z -space. (The x -axis corresponds to the length, the y -axis to the width and the z -axis to the height.) The normalized position of an object T_i with respect to the reference system is characterized by the following conditions:

$$\min\{x : (x, y, z) \in T_i\} = 0, \quad \min\{y : (x, y, z) \in T_i\} = 0, \quad \min\{z : (x, y, z) \in T_i\} = 0.$$

The normalized position of the region B is defined analogously. The translation of T_i to $v^i \in R^n$ yields the set of points

$$T_i(v^i) := T_i + v^i := \{v \in R^n : v = v^i + w, w \in T_i\}.$$

The point v_i is called the allocation point of $T_i(v^i)$ or for short, the allocation point of T_i .

Now, the considered packing problem can be formulated as follows: Find a subset I^* of $I := \{1, \dots, m\}$ (a subset of the objects) and the corresponding allocation points $v^i, i \in I^*$, such that the packing conditions are fulfilled and the total value of the packed objects is a maximum. Hence:

$$\sum_{i \in I^*} c_i \rightarrow \max$$

subject to

$$\begin{aligned}
I^* &\subset I, \\
T_i(v^i) &\subset B \quad \text{for } i \in I^*, \\
\text{int}(T_i(v^i)) \cap \text{int}(T_j(v^j)) &= \emptyset \quad \text{for } i \neq j, i, j \in I^*.
\end{aligned} \tag{1}$$

This model is contained in [6]. More detailed theoretical and practical investigations require the knowledge of the structure of the region B and the objects T_i

3. MODELS OF ONE-DIMENSIONAL PACKING PROBLEMS

In the one-dimensional case the region B and the objects T_i can be considered as intervals $[0, L]$ and $[0, l_i]$, $i = 1, \dots, m$, respectively. The task is to find allocation points $v^i = x_i$ of the objects T_i .

For any feasible packing which contains the objects T_i and T_j with $x_i > x_j$ there exists a packing with the same valuation for which $x_i < x_j$ holds. For that reason, without loss of generality, it can be assumed that $x_i + l_i \leq x_j$ is valid for any two packed objects T_i and T_j with $i < j$.

To describe the set I^* of the packed objects T_i 0/1-variables a_i are defined as follows:

$$a_i = \begin{cases} 1 & , T_i \text{ is contained in } B, \\ 0 & , \text{otherwise.} \end{cases} \tag{2}$$

The set-theoretical model (1) has now the following form:

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max$$

subject to

$$x_1 \geq 0, \quad x_i + a_i l_i \leq x_{i+1}, \quad i = 1, \dots, m-1, \quad x_m + a_m l_m \leq L, \tag{3}$$

$$a_i \in \{0, 1\}, \quad i = 1, \dots, m.$$

Here, m continuous, m 0/1-variables and $m+1$ inequalities are used in model (3). Fixing of x_i according to

$$x_1 = 0, \quad x_i = \sum_{j=1}^{i-1} l_j a_j, \quad i = 2, \dots, m,$$

corresponds to a dense packing of the objects, or also to the consideration of so-called "normalized patterns" ([2], "left-justified"). Doing so, the well-known 0/1 knapsack problem arises from the model (3):

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max$$

subject to

$$\sum_{i=1}^m l_i a_i \leq L \quad a_i \in \{0, 1\}, i = 1, \dots, m.$$

Only m 0/1-variables and 1 inequality are contained in the knapsack problem.

4. MODELS FOR TWO-DIMENSIONAL PACKING PROBLEMS

4.1 Packing of Rectangles within a Rectangle

Within this section it is assumed that the region B is rectangular as well as all objects T_i which are edge-parallel to the reference system. Then B can be characterized by its length L and width W . T_i has length l_i and width w_i ($i = 1, \dots, m$). Only edge-parallel allocations are allowed.

The aim of the following investigations is to develop a linear mixed-integer optimization model. To describe the index set I^* 0/1-variables a_i are introduced in analogy to section 3 according to (2).

The modeling is based on the following idea:

Let be given a sufficiently large rectangle B' of length L' , $L' \geq L$, and width W' , $W' \geq W$, in which all objects can be packed. But only the total containment of the piece T_i within B gets a positive valuation. This can be realized using the inequalities (5) where x_i, y_i are the coordinates of the allocation point v^i of T_i ($i = 1, \dots, m$).

To characterize the mutual position of two objects $T_i(v^i)$ and $T_j(v^j)$ the notions “right-left” and “above-below” are used.

Thereby

$$\begin{aligned} \text{“} T_j \text{ right (left) to } T_i \text{” if } p = (x, y) \in T_j(v^j) \Rightarrow x \geq x_i + l_i \text{ (resp. } x \leq x_i), \\ \text{“} T_j \text{ above (below) to } T_i \text{” if } p = (x, y) \in T_j(v^j) \Rightarrow y \geq y_i + w_i \text{ (resp. } y \leq y_i). \end{aligned}$$

To describe the mutual position of T_i and T_j the following 0/1-variables are defined:

$$v_{ij}^r = \begin{cases} 1 & , \quad \text{if } T_j \text{ is right to } T_i, \\ 0 & , \quad \text{otherwise,} \end{cases} \quad u_{ij}^o = \begin{cases} 1 & , \quad \text{if } T_j \text{ is above to } T_i, \\ 0 & , \quad \text{otherwise.} \end{cases}$$

The condition $\text{int}(T_i(v^i)) \cap \text{int}(T_j(v^j)) = \emptyset$ is fulfilled if and only if T_j is right or left or above or below to T_i .

Hence, the condition $\text{int}(T_i(v^i)) \cap \text{int}(T_j(v^j)) = \emptyset$ can be described by the system of inequalities (6). Summarizing, the following linear mixed-integer model is obtained:

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max \quad (4)$$

subject to

$$\begin{aligned} 0 \leq x_i, \quad x_i &\leq (L - l_i)a_i + L'(1 - a_i), \quad i = 1, \dots, m, \\ 0 \leq y_i, \quad y_i &\leq (W - w_i)a_i + W'(1 - a_i), \quad i = 1, \dots, m, \\ a_i &\in \{0, 1\}, \quad i = 1, \dots, m; \end{aligned} \quad (5)$$

and

$$\begin{aligned} x_i + l_i &\leq x_j + L'(1 - u_{ij}^r), \quad x_j + l_j \leq x_i + L'(1 - u_{ji}^r), \\ y_i + w_i &\leq y_j + W'(1 - u_{ij}^o), \quad y_j + w_j \leq y_i + W'(1 - u_{ji}^o), \\ u_{ij}^r + u_{ji}^r + u_{ij}^o + u_{ji}^o &\geq 1, \\ u_{ij}^r, u_{ji}^r, u_{ij}^o, u_{ji}^o &\in \{0, 1\}, \quad i = 1, \dots, m-1, \quad j = i+1, \dots, m. \end{aligned} \quad (6)$$

The model (4) – (6) contains $2m$ continuous, $m + 2m(m-1)$ 0/1-variables and $4m + 5m(m-1)/2$ inequalities.

Instead of the 4 variables $u_{ij}^r, u_{ji}^r, u_{ij}^o, u_{ji}^o$ in (6) only 2 0/1-variables u_{ij}, v_{ij} are sufficient to describe the four possibly mutual positions of two objects. (6) is equivalent to the system of inequalities (7):

$$\begin{aligned} x_i + l_i &\leq x_j + L'(2 - u_{ij} - v_{ij}) \\ x_j + l_j &\leq x_i + L'(1 - u_{ij} + v_{ij}) \\ y_i + w_i &\leq y_j + W'(1 + u_{ij} - v_{ij}) \\ y_j + w_j &\leq y_i + W'(u_{ij} + v_{ij}) \\ u_{ij}, v_{ij} &\in \{0, 1\}, \quad i = 1, \dots, m-1, \quad j = i+1, \dots, m. \end{aligned} \quad (7)$$

Since $u_{ij}, v_{ij} \in \{0, 1\}$ exactly one of the four conditions is non-trivial. Hence, $2m$ continuous, but only $m + m(m-1)$ 0/1-variables and $4m + 2m(m-1)$ inequalities are used in model (4,5,7).

4.2. PACKING OF CONVEX POLYGONS

Within this section it is assumed that the region B as well as the objects $T_i (i = 1, \dots, m)$ are convex polygons.

4.2.1 First Model

The region $T_0 := B$ and the objects T_i can be described as the intersection of k_i closed half-spaces (half-planes). Without loss of generality, it is assumed that the k_i corner points $e_{ik}, k = 1, \dots, k_i$, of $T_i (i = 0, 1, \dots, m)$ are numbered clockwise and let e_{i1} be that corner point having x -coordinate 0 and smallest y -coordinate (the normalized position of T_i with respect to the reference system is assumed). Hence T_i can be described as follows:

$$T_i = \{(x, y) \in R^2 : g_{ik}(x, y) := a_{ik}x + b_{ik}y + c_{ik} \leq 0, k = 1, \dots, k_i\},$$

where the corner point $e_{ik} = (x_{ik}, y_{ik})$ is the intersection of g_{ik} and $g_{i,k+1}, k = 1, \dots, k_i, (k_i + 1 \equiv 1), i = 1, \dots, m$.

The translation of T_i from the normalized position along the vector $v^i = (x_i, y_i)$ yields

$$T_i(v^i) = T_i(x_i, y_i) = \{(x, y) \in R^2 : g_{ik}(x - x_i, y - y_i) \leq 0, k = 1, \dots, k_i\}.$$

(x_i, y_i) is the allocation point of T_i .

The translated objects $T_i(v^i)$ and $T_j(v^j)$ do not overlap each other if they can be separated by a hyperplain resp. $T_i(v^i) \cap T_j(v^j)$ is part of a hyperplain. In both cases there exists a bounding hyperplain $g_{ik}(x - x_i, y - y_i) \geq 0$ of $T_i(v^i)$ such that for $(x, y) \in T_j(v^j)$ $g_{ik}(x - x_i, y - y_i) \geq 0$ is valid ($k \in \{1, \dots, k_i\}$), or a corresponding hyperplain of $T_j(v^j)$.

To test whether two fixed objects T_i and T_j do not overlap each other, it is sufficient, because of the convexity of the objects, to consider the bounding hyperplains of $T_i(v^i)$ and the corner points of $T_j(v^j)$ resp. the hyperplains of $T_j(v^j)$ and the corner points of $T_i(v^i)$. It holds:

Statement 1: *If there exists an index $k, k \in \{1, \dots, k_i\}$, such that for all corner points $(x_{jp}, y_{jp}), p = 1, \dots, k_j$,*

$$g_{ik}(x_{jp} + x_j - x_i, y_{jp} + y_j - y_i) = a_{ik}x_{jp} + b_{ik}y_{jp} + g_{ik}(x_j - x_i, y_j - y_i) \geq 0,$$

then $T_i(v^i)$ and $T_j(v^j)$ do not overlap each other.

Let

$$d_{ik}^j := \min_{p=1, \dots, k_j} \{a_{ik}x_{jp} + b_{ik}y_{jp}\}, i = 1, \dots, m, k = 1, \dots, k_i, j = 1, \dots, m, j \neq i.$$

The statement $\text{int}(T_i(v^i)) \cap \text{int}(T_j(v^j)) = \emptyset$ is true if at least one of the following $k_i + k_j$ conditions is fulfilled:

$$-g_{ik}(x_j - x_i, y_j - y_i) \leq d_{ik}^j, \quad k = 1, \dots, k_i \quad (8)$$

(Separating of $T_j(v^j)$ using a hyperplain of $T_i(v^i)$),

$$-g_{jk}(x_i - x_j, y_i - y_j) \leq d_{jk}^i, \quad k = 1, \dots, k_j. \quad (9)$$

(Separating of $T_i(v^i)$ using a hyperplain of $T_j(v^j)$).

The containment of $T_i(v^i)$, $i = 1, \dots, m$, within $B(= T_0)$ is equivalent to

$$g_{0k}(x_{ip} + x_i, y_{ip} + y_i) \leq 0, \quad p = 1, \dots, k_i, \quad k = 1, \dots, k_0.$$

Let $d_{0k}^i := \max_{p=1, \dots, k_i} \{a_{0ik}x_{ip} + b_{0k}y_{ip}\}$. Then $T_i(v^i) \subset B$ if and only if the following inequalities are fulfilled:

$$g_{0k}(x_i, y_i) \leq -d_{0k}^i, \quad k = 1, \dots, k_0 \quad (10)$$

Again, 0/1-variables a_i , $i = 1, \dots, m$, are used according to (2) to characterize the containment of T_i within B . Additionally, 0/1-variables w_{ik}^j are defined as follows to describe the mutual position of two objects $T_i(v^i)$ and $T_j(v^j)$, $i \neq j$:

$$w_{ik}^j = \begin{cases} 1 & , \quad \text{if } T_j(v^j) \text{ lies in the half-plain } g_{ik}(x - x_i, y - y_i) \geq 0, \\ 0 & , \quad \text{otherwise,} \end{cases}$$

$$i = 1, \dots, m, \quad k = 1, \dots, k_i, \quad j = 1, \dots, m, \quad i \neq j.$$

By this a natural generalization is given of the 0/1-variables in section 4.1. Using the same linear objective function (4) and the restrictions (8)–(10) the following linear mixed-integer model is obtained where M is a sufficiently large number.

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max \quad (11)$$

subject to

$$g_{0k}(x_i, y_i) + d_{0k}^i \leq M(1 - a_i), \quad a_i \in \{0, 1\}, \quad i = 1, \dots, m, \quad k = 1, \dots, k_0 \quad (12)$$

and

$$-g_{ik}(x_j - x_i, y_j - y_i) - d_{ik}^j \leq M(1 - u_{ik}^j),$$

$$k = 1, \dots, k_i, i = 1, \dots, m, j = 1, \dots, m, i \neq j. \quad (13)$$

$$\sum_{k=1}^{k_i} u_{ik}^j + \sum_{k=1}^{k_j} u_{jk}^i \geq 1, \quad i = 1, \dots, m-1, j = i+1, \dots, m, \quad (14)$$

$$u_{ik}^j \in \{1, 0\}, \quad i = 1, \dots, m, k = 1, \dots, k_i, j = 1, \dots, m, i \neq j.$$

If $a_i = 1$ then (12) secures $T_i(v^i) \subset B$. (13,14) ensures that $T_i(v^i)$ and $T_j(v^j)$ do not overlap each other.

The model (11)–(14) uses $2m$ continuous, $m + (m-1) \sum_{i=1}^m k_i$ 0/1-variables and $mk_0 + m(m-1)/2 + (m-1) \sum_{i=1}^m k_i$ inequalities. With respect to the fact that for fixed $i, j, i \neq j$, only one of the inequalities

$$-g_{ik}(x_j - x_i, y_j - y_i) - d_{ik}^j \leq M(1 - u_{ik}^j), \quad k = 1, \dots, k_i$$

$$-g_{jk}(x_i - x_j, y_i - y_j) - d_{jk}^i \leq M(1 - u_{jk}^i), \quad k = 1, \dots, k_j$$

has to be fulfilled, an equivalent model with¹ $\lceil \log_2(k_i + k_j) \rceil$ 0/1-variables can be obtained in analogy to (7). Then the inequalities (14) are unnecessary.

Hence, the model has $2m$ continuous, $m + \sum_{i=1}^{m-1} \sum_{j=i+1}^m \lceil \log_2(k_i + k_j) \rceil$ 0/1-variables and $mk_0 + \sum_{i=1}^{m-1} \sum_{j=i+1}^m (k_i + k_j) = mk_0 + (m-1) \sum_{i=1}^m k_i$ inequalities.

In the case of rectangular objects ($k_i = 4, i = 1, \dots, m$) we have the same number of continuous and 0/1-variables but $4m + 4m(m-1)$ inequalities. In section 4.1. additionally the fact could be exploited that the rectangular objects have to be parallel to the edges of the region.

4.2.2 Example

Let be given a region B and two convex polygons T_1 and T_2 as defined below and shown in Fig. 1.


$$B = \{(x, y) : g_{01}(x, y) := -x - y + 1 \leq 0, g_{02}(x, y) := -x \leq 0, g_{03}(x, y) := y - 2 \leq 0, g_{04}(x, y) := x - 2 \leq 0, g_{05}(x, y) := -y \leq 0\},$$

$$T_1 = \{(x, y) : g_{11}(x, y) := -y \leq 0, g_{12}(x, y) := -2x + y \leq 0, g_{13}(x, y) := x - 1 \leq 0\},$$

$$T_2 = \{(x, y) : g_{21}(x, y) := -x - y + 1 \leq 0, g_{22}(x, y) := -x + y - 1 \leq 0, g_{23}(x, y) := y - 2 \leq 0, g_{24}(x, y) := 2x - y - 2 \leq 0\}.$$

With $d_{01}^1 = d_{02}^1 = d_{05}^1 = 0, d_{03}^1 = 2, d_{04}^1 = 1$ and $d_{02}^2 = d_{05}^2 = 0, d_{01}^2 = -1, d_{03}^2 = d_{04}^2 = 2$, then (12) is as follows:

¹ $\lceil x \rceil$ denotes the smallest integer not smaller than x .

Figure 1 Region B and given objects T_1 and T_2

$$\begin{aligned}
 -x_1 - y_1 + 1 &\leq M(1 - a_1), & -x_2 - y_2 &\leq M(1 - a_2), \\
 -x_1 &\leq M(1 - a_1), & -x_2 &\leq M(1 - a_2), \\
 y_1 &\leq M(1 - a_1), & y_2 &\leq M(1 - a_2), \\
 x_1 - 1 &\leq M(1 - a_1), & x_2 &\leq M(1 - a_2), \\
 -y_1 &\leq M(1 - a_1), & -y_2 &\leq M(1 - a_2); \\
 a_1, a_2 &\in \{0, 1\}.
 \end{aligned}$$

Condition (13) has the form (because of $d_{11}^2 = d_{12}^2 = -2$, $d_{13}^2 = 0$ and $d_{21}^1 = -3$, $d_{22}^1 = -1$, $d_{23}^1 = d_{24}^1 = 0$):

$$\begin{aligned}
 y_2 - y_1 + 2 &\leq M(1 - u_{11}^2), & (x_1 - x_2) + (y_1 - y_2) + 2 &\leq M(1 - u_{21}^1), \\
 2(x_2 - x_1) - (y_2 - y_1) + 2 &\leq M(1 - u_{12}^2), & (x_1 - x_2) + (y_1 - y_2) + 2 &\leq M(1 - u_{22}^1), \\
 -(x_2 - x_1) + 1 &\leq M(1 - u_{13}^2), & -(y_1 - y_2) + 2 &\leq M(1 - u_{23}^1), \\
 & & -2(x_1 - x_2) + (y_1 - y_2) + 2 &\leq M(1 - u_{24}^1),
 \end{aligned}$$

$$\begin{aligned}
 u_{11}^2 + u_{12}^2 + u_{13}^2 + u_{21}^1 + u_{22}^1 + u_{23}^1 + u_{24}^1 &\geq 1, \\
 u_{11}^2, u_{12}^2, u_{13}^2, u_{21}^1, u_{22}^1, u_{23}^1, u_{24}^1 &\in \{0, 1\}.
 \end{aligned}$$

The objective function

$$z = c_1 a_1 + c_2 a_2 \quad (c_1 > 0, c_2 > 0)$$

has the maximum value $c_1 + c_2$ and the optimal solution is as follows:

$$\begin{aligned}
 a_1 = a_2 &= 1, & u_{12}^2 = u_{24}^1 &= 1, \\
 u_{11}^2 = u_{13}^2 = u_{21}^1 = u_{22}^1 = u_{23}^1 &= 0, \\
 (x_1, y_1) &= (1, 0), & (x_2, y_2) &= (0, 0).
 \end{aligned}$$

Remark : Instead of the use of $7 \lceil \log_2(k_1 + k_2) \rceil$ 0/1-variables u_{ik}^i , it is sufficient to define $3 = \lceil \log_2(k_1 + k_2) \rceil$ 0/1-variables.

4.2.3 Modeling with Hodographs

The description of the non-overlapping of two objects $T_i(v^i)$ and $T_j(v^j)$ is an essential part when handling packing problems. In the previous section, therefore the position of corner points of $T_j(v^j)$ with respect to bounding hyperplanes of $T_i(v^i)$, and reversely, is used. A second possibility consists in characterizing the beginning of the overlapping of T_i by $T_j(v^j)$. With other words, which are the allocation points v_j of $T_j(v^j)$ such that $\text{int } T_i \cap \text{int } T_j(v^j) = \emptyset$ and $T_i \cap T_j(v^j) \neq \emptyset$ (T_i is considered to be in normalized position). The set of such allocation points is called (outer) hodograph of the dense allocation of T_i and T_j (see also in [6] and [8]). The **hodograph** of two convex polygons T_i and T_j is the boundary of a convex polygon H_{ij} defined according to

$$H_{ij} = \{(x, y) \in R^2 : ((x, y) + T_j) \cap T_i \neq \emptyset\}.$$

Hence, H_{ij} can also be described using closed half-planes:

$$H_{ij} := \{(x, y) \in R^2 : h_k^{ij}(x, y) \leq 0, \quad k = 1, \dots, k_{ij}\}. \quad (15)$$

On the other hand, H_{ij} can be characterized by hyperplanes given T_i and T_j .

Statement 2: Let T_i and T_j be convex polygons and let $g_{ik} = a_{ik}x + b_{ik}y + c_{ik}$, $k = 1, \dots, k_i$, and $g_{jk} = a_{jk}x + b_{jk}y + c_{jk}$, $k = 1, \dots, k_j$, be the bounding hyperplanes of T_i and T_j . Then it holds with $p = (p_x, p_y)$:

$$\begin{aligned} H_{ij} = \{(x, y) \in R^2 : & g_{ik}(x, y) + \min_{p \in T_j} g_{ik}(p_x, p_y) - c_{ik} \leq 0, \quad k = 1, \dots, k_i, \\ & -g_{jk}(x, y) + \min_{p \in T_i} g_{jk}(p_x, p_y) + c_{jk} \leq 0, \quad k = 1, \dots, k_j\} \end{aligned} \quad (16)$$

Proof: Let $v = (x, y) \in H_{ij}$. Then, per definition, $T_j(v) \cap T_i \neq \emptyset$. Hence, for any bounding hyperplane of $T_j(v)$ there exists at least one element $(\bar{x}, \bar{y}) \in T_i$ such that $g_{jk}(\bar{x} - x, \bar{y} - y) \leq 0$. Moreover:

$$-g_{jk}(x, y) + \min_{p \in T_i} g_{jk}(p_x, p_y) + c_{jk} \leq 0, \quad k = 1, \dots, k_j.$$

Since $(v + T_j) \cap T_i \neq \emptyset$ if and only if $T_j \cap (T_i - v) \neq \emptyset$, one has:

$$g_{ik}(x, y) + \min_{p \in T_j} g_{ik}(p_x, p_y) - c_{ik} \leq 0, \quad k = 1, \dots, k_i \quad \blacksquare$$

Hence, the functions h_k^{ij} for (15) are chosen from the functions used in the inequalities in (16). The non-overlapping condition for $T_i(v^i)$ and $T_j(v^j)$ requires, using (15), that at least one of the inequalities

$$h_k^{ij}(x_j - x_i, y_j - y_i) \geq 0, \quad k = 1, \dots, k_{ij}, \quad (17)$$

is fulfilled. Since the non-overlapping of T_j and T_i is ensured if T_i and T_j are non-overlapping. $\sum_{i=1}^{m-1} \sum_{j=i+1}^m k_{ij} =: K$ inequalities of the type (17) are to consider. Since the set H_{ij} has at most $k_i + k_j$ corner points (compare in [8], page 67), that is $k_{ij} \leq k_i + k_j$, it follows $K \leq (m-1) \sum_{i=1}^m k_i$. Hence, in the case $k_{ij} < k_i + k_j$, the consideration of K inequalities of the type (17) is more favourable than to consider $(m-1) \sum_{i=1}^m k_i$ inequalities of type (8) resp. (9).

To guarantee that at least one of the k_{ij} inequalities is fulfilled, $\lceil \log_2(k_{ij}) \rceil$ 0/1-variables u_p^{ij} , $p = 1, \dots, v$, with $v = v_{ij} := \lceil \log_2(k_{ij}) \rceil$ are necessary. The k -th inequality

$$h_k^{ij}(x_j - x_i, y_j - y_i) + M \left(\sum_{p=1}^v \alpha_p^k + \sum_{p=1}^v (-1)^{\alpha_p^k} u_p^{ij} \right) \geq 0, \quad (18)$$

(where $\alpha_1^k, \dots, \alpha_v^k$ are the coefficients of the binary representation of $2^v - k$, i.e. $\sum_{p=1}^v 2^{v-p} \alpha_p^k = 2^v - k$.) is non-trivial if and only if $u_p^{ij} = \alpha_p^k$ for $p = 1, \dots, v$. If $2^v > k_{ij}$ then the inequality

$$\sum_{p=1}^v 2^{v-p} u_p^{ij} \geq 2^v - k_{ij}$$

ensures that only these values $(u_1^{ij}, \dots, u_v^{ij})$ are feasible which are required.

The characterization of the non-overlapping of T_i and T_j using (17) yields in general a reducing of the number of restrictions. A similar result can be achieved if the inner hodograph is used to characterize the containment of T_i within B .

The **inner hodograph** is defined to be the boundary of the set

$$H_{0i} = \{(x, y) \in \mathbb{R}^2 : (x, y) + T_i \subset B\}.$$

H_{0i} is the set of all feasible allocation points (x, y) of T_i which possibly is empty. H_{0i} is a convex polygon if B and T_i are convex polygons. If k_{0i} denotes the number of corner points of H_{0i} then $k_{0i} \leq k_0$ holds. Therefore, H_{0i} can be described as follows:

$$H_{0i} = \{(x, y) \in \mathbb{R}^2 : h_k^{0i}(x, y) \leq 0, k = 1, \dots, k_{0i}\}.$$

Hence, $T_i(v^i) \subset B$ if and only if

$$h_k^{0i}(x_i, y_i) \leq 0, \quad k = 1, \dots, k_{0i}. \quad (19)$$

In the case $k_{0i} < k_0$, the system of inequalities (19) contains fewer inequalities than the system (10) does.

Using the inequalities of type (18) and (19) the model (11)–(13) is equivalent to:

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max \quad (20)$$

subject to

$$\begin{aligned} h_k^{0i}(x_i, y_i) &\leq M(1 - a_i), \quad k = 1, \dots, k_{0i}, \quad i = 1, \dots, m; \\ a_i &\in \{0, 1\}, \quad i = 1, \dots, m; \end{aligned} \quad (21)$$

$$h_k^{ij}(x_j - x_i, y_j - y_i) + M \left(\sum_{p=1}^{v_{ij}} \alpha_p^{ijk} + \sum_{p=1}^{v_{ij}} (-1)^{\alpha_p^{ijk}} u_p^{ij} \right) \geq 0, \quad (22)$$

with $v_{ij} = \lceil \log_2(k_{ij}) \rceil$; $i = 1, \dots, m-1$, $k = 1, \dots, k_{ij}$, $j = i+1, \dots, m$;

$$\sum_{p=1}^{v_{ij}} 2^{v_{ij}-p} u_p^{ij} \geq 2^{v_{ij}} - k_{ij}, \quad \text{if } 2^{v_{ij}} > k_{ij}; \quad (23)$$

$$u_1^{ij}, \dots, u_{v_{ij}}^{ij} \in \{0, 1\}, \quad i = 1, \dots, m-1, \quad j = i+1, \dots, m.$$

The model (20)–(23) contains $2m$ continuous, $m + \sum_{i=1}^{m-1} \sum_{j=i+1}^m v_{ij}$ 0/1-variables and $\sum_{i=1}^m k_{0i} + \sum_{i=1}^{m-1} \sum_{j=i+1}^m k_{ij}$ inequalities and at most furthermore $m(m-1)/2$ inequalities. Since

$$\sum_{i=1}^{m-1} \sum_{j=i+1}^m v_{ij} = \sum_{i=1}^{m-1} \sum_{j=i+1}^m \lceil \log_2(k_{ij}) \rceil < \sum_{i=1}^{m-1} \sum_{j=i+1}^m k_{ij} \leq (m-1) \sum_{i=1}^m k_i,$$

in the model above are used essentially fewer 0/1-variables in comparison to model (11)–(13). Because of $\sum_{i=1}^m k_{0i} \leq mk_0$, also fewer inequalities are needed in general.

4.2.4 Example

The region and objects of the example in 4.2.2 are used. In Fig 2 the hodograph of T_1 and T_2 and the set H_{12} are shown.

It holds:

$$\begin{aligned} H_{12} = \{(x, y) \in R^2 : & h_1^{12}(x, y) = -2x + y - 2 \leq 0, \\ & h_2^{12}(x, y) = x + y - 2 \leq 0, \\ & h_3^{12}(x, y) = x - 1 \leq 0, \\ & h_4^{12}(x, y) = x - y - 2 \leq 0, \\ & h_5^{12}(x, y) = -y - 2 \leq 0\}. \end{aligned}$$




Figure 2 Hodograph of T_1 and T_2 and H_{12}

Because of $v = \lceil \log_2(5) \rceil = 3$, three 0/1-variables u_1, u_2, u_3 are needed to describe the condition $\text{int}(T_1(v^1)) \cap \text{int}(T_2(v^2)) = \emptyset$.

$$\begin{aligned} -2(x_2 - x_1) + y_2 - y_1 - 2 + M(3 - u_1 - u_2 - u_3) &\geq 0, \\ x_2 - x_1 + y_2 - y_1 - 2 + M(2 - u_1 - u_2 + u_3) &\geq 0, \\ x_2 - x_1 - 1 + M(2 - u_1 + u_2 - u_3) &\geq 0, \\ x_2 - x_1 - (y_2 - y_1) - 2 + M(1 - u_1 + u_2 + u_3) &\geq 0, \\ -(y_2 - y_1) - 2 + M(2 + u_1 - u_2 - u_3) &\geq 0, \end{aligned}$$

and

$$4u_1 + 2u_2 + u_3 \geq 2^3 - 5 = 3.$$

The last inequality secures that exact one of the first five inequalities is non-trivial.

4.3. PACKING OF ARBITRARY POLYGONS

4.3.1 Modeling

For problems of packing convex objects it was shown in section 4.2 that the containment of T_i within B can be described in a good manner by using the inner hodograph,

and in a similarly good way the non-overlapping of T_i and T_j can be handled using the hodograph of these two objects. Now, these ideas will be translated to the case of arbitrary polygonal objects.

4.3.1.1 Containment of T_i within B

The set

$$H_{0i} := \{ (x, y) \in R^2 : (x, y) + T_i \subset B \}$$

is the set of all feasible allocation points (x, y) of T_i . In contrast to the convex case, H_{0i} is in general not connected (see the example in 4.3.2). But H_{0i} is always a union of convex polygons. Hence, if H_{0i} consists of k_{0i} convex polygons then it can be described as follows:

$$\begin{aligned} H_{0i} = & \left\{ (x, y) \in R^2 : ((h_{11}^{0i}(x, y) \leq 0) \wedge \cdots \wedge (h_{1l_{0i}^{0i}}^{0i}(x, y) \leq 0)) \right. \\ & \left. \vee \cdots \vee ((h_{k_{0i}1}^{0i}(x, y) \leq 0) \wedge \cdots \wedge (h_{k_{0i}l_{k_{0i}}^{0i}}^{0i}(x, y) \leq 0)) \right\} \\ = & \left\{ (x, y) \in R^2 : \bigvee_{j=1}^{k_{0i}} \bigwedge_{l=1}^{l_j^{0i}} (h_{jl}^{0i}(x, y) \leq 0) \right\}. \end{aligned}$$

Moreover, $T_i(v^i) \subset B$ if and only if $v^i = (x_i, y_i)$ is an element of one of the k_{0i} convex polygons of H_{0i} . Or, if (x_i, y_i) fulfills one of the k_{0i} systems of inequalities

$$h_{j1}^{0i}(x_i, y_i) \leq 0, \dots, h_{jl_j^{0i}}^{0i}(x_i, y_i) \leq 0, \quad j = 1, \dots, k_{0i}, \quad (24)$$

then $T_i(x_i, y_i)$ is contained in B .

As demonstrated in 4.2.3 the “or”-conditions can be handled by using of 0/1-variables. Let $\lceil \log_2(k_{0i}) \rceil = v_{0i}$ and let $(\alpha_1^{0ij}, \dots, \alpha_{v_{0i}}^{0ij})$ be the binary coefficients of $2^{v_{0i}} - j$ in accordance to $\sum_{p=1}^{v_{0i}} 2^{v_{0i}-p} \alpha_p^{0ij} = 2^{v_{0i}} - j$, $j = 1, \dots, k_{0i}$.

Using the 0/1-variables $u_1^{0i}, \dots, u_{v_{0i}}^{0i}$ the following system of inequalities results from (24):

$$h_{jl}^{0i}(x_i, y_i) \leq M \left(\sum_{p=1}^{v_{0i}} \alpha_p^{0ij} + \sum_{p=1}^{v_{0i}} (-1)^{\alpha_p^{0ij}} u_p^{0i} \right), \quad l = 1, \dots, l_j^{0i}, \quad (25)$$

$$u_p^{0i} \in \{0, 1\}, \quad p = 1, \dots, v_{0i}.$$

Only if $u_p^{0i} = \alpha_p^{0ij}$, $p = 1, \dots, v_{0i}$, then (25) is non-trivial. M has to be sufficiently large, e.g. $M \geq \max_{i=1, \dots, m} \max_{l=1, \dots, l_j^{0i}} \max_{(x, y) \in H_{0i}} h_{jl}^{0i}(x, y)$.

In analogy to previous models 0/1-variables a_i are used defined in (2). Now, $T_i(v^i) \subset B$, $i = 1, \dots, m$, can be modelled as follows:

$$h_{jl}^{0i}(x_i, y_i) - M \left(\sum_{p=1}^{v_{0i}} \alpha_p^{0ij} + \sum_{p=1}^{v_{0i}} (-1)^{\alpha_p^{0ij}} u_p^{0i} \right) \leq M(1 - a_i), \quad (26)$$

$$l = 1, \dots, l_j^{0i}, \quad j = 1, \dots, k_{0i},$$

$$u_p^{0i} \in \{0, 1\}, \quad p = 1, \dots, v_{0i} = \lceil \log_2(k_{0i}) \rceil,$$

and, if $2^{v_{0i}} > k_{0i}$ then additionally

$$\sum_{p=1}^{v_{0i}} 2^{v_{0i}-p} u_p^{0i} \geq 2^{v_{0i}} - k_{0i},$$

$$a_i \in \{0, 1\}, \quad i = 1, \dots, m.$$

4.3.1.2 Non-overlapping of T_i and T_j

The hodograph of the dense packing of T_j and T_i (in normalized position) is again defined to be the boundary of the set

$$H_{ij} := \{(x, y) \in R^2 : ((x, y) + T_j) \cap T_i \neq \emptyset\}.$$

The set H_{ij} is, in contrast to H_{0i} , always connected, but not necessarily convex.

The description of non-convex polygons T can be done in different ways e.g. as a union of convex polygons. Here, another possibility is chosen. The basis is the convex hull $\text{conv}(T)$ of T . Any sequence of bounding edges of T between two neighbourhood corner points of $\text{conv}(T)$ can be described by the corresponding straight lines resp. half-planes. After suitable transformations of the linear systems of inequalities which are connected by “ \wedge ”- and “ \vee ”-relations, T can be represented in the following form:

$$T = \left\{ (x, y) \in R^2 : \left((g_{11}(x, y) \leq 0) \vee \dots \vee (g_{1l_1}(x, y) \leq 0) \right) \right. \\ \left. \wedge \dots \wedge \left((g_{kl_1}(x, y) \leq 0) \vee \dots \vee (g_{kl_k}(x, y) \leq 0) \right) \right\}.$$

Hence, $(x, y) \notin \text{int}(T)$ if and only if

$$\left((g_{11}(x, y) \geq 0) \wedge \dots \wedge ((g_{1l_1}(x, y) \geq 0)) \right) \\ \vee \dots \vee \left((g_{kl_1}(x, y) \geq 0) \wedge \dots \wedge ((g_{kl_k}(x, y) \geq 0)) \right).$$

Let

$$T = H_{ij} = \left\{ (x, y) \in R^2 : (h_{11}^{ij}(x, y) \leq 0) \vee \cdots \vee (h_{1l_1^j}^{ij}(x, y) \leq 0) \right. \\ \left. \wedge \cdots \wedge (h_{k_{ij}1}^{ij}(x, y) \leq 0) \vee \cdots \vee (h_{k_{ij}l_{k_{ij}}^{ij}}^{ij}(x, y) \leq 0) \right\}.$$

Then, the non-overlapping of $T_i(v^i)$ and $T_j(v^j)$ is equivalent to:

$$\left((h_{11}^{ij}(x_j - x_i, y_j - y_i) \geq 0) \wedge \cdots \wedge (h_{1l_1^j}^{ij}(x_j - x_i, y_j - y_i) \geq 0) \right) \\ \vee \cdots \vee \left((h_{k_{ij}1}^{ij}(x_j - x_i, y_j - y_i) \geq 0) \wedge \cdots \wedge (h_{k_{ij}l_{k_{ij}}^{ij}}^{ij}(x_j - x_i, y_j - y_i) \geq 0) \right) \\ = \bigvee_{r=1}^{k_{ij}} \bigwedge_{l=1}^{l_r^{ij}} (h_{rl}^{ij}(x_j - x_i, y_j - y_i) \geq 0),$$

i.e., if (x_j, y_j) fulfills one of the above systems of inequalities then $T_i(v^i)$ and $T_j(v^j)$ do not overlap each other.

The handling of the “ \vee ”-conditions can be done by using 0/1-variables in a similar way as in 4.3.1.1 for the description of H_{0i} . It results k_{ij} systems of inequalities of the type (25). Hence, the non-overlapping of $T_i(v^i)$ and $T_j(v^j)$, $i = 1, \dots, m-1$, $j = i+1, \dots, m$, can be characterized as follows:

$$h_{r1}^{ij}(x_j - x_i, y_j - y_i) + M \left(\sum_{p=1}^{v_{ij}} \alpha_p^{ijr} + \sum_{p=1}^{v_{ij}} (-1)^{\alpha_p^{ijr}} u_p^{ij} \right) \geq 0, l = 1, \dots, l_r^{ij}, \quad (27)$$

$$r = 1, \dots, k_{ij}, u_1^{ij}, \dots, u_{v_{ij}}^{ij} \in \{0, 1\}, i = 1, \dots, m-1, j = i+1, \dots, m,$$

and if $2^{v_{ij}} - k_{ij} > 0$ then additionally

$$\sum_{p=1}^{v_{ij}} 2^{v_{ij}-p} u_p^{ij} \geq 2^{v_{ij}} - k_{ij}$$

where $v_{ij} = \lceil \log_2(k_{ij}) \rceil$, $\sum_{p=1}^{v_{ij}} 2^{v_{ij}-p} \alpha_p^{ijr} = 2^{v_{ij}} - r$. The constant M has to be sufficiently large, e.g.

$$M \geq - \min_{r=1, \dots, k_{ij}} \min_{q=1, \dots, l_r^{ij}} \min_{(x, y) \in H_{ij}} h_{rq}^{ij}(x, y).$$




Figure 3 a) Region B and objects T_1, T_2 and T_3 ; b) set H_{01}

Hence, the problem of packing arbitrary polygons can be written as follows:

$$z = \sum_{i=1}^m c_i a_i \rightarrow \max \quad \text{subject to (26) and (27) .}$$

Here, $2m$ continuous, $m + \sum_{i=1}^m v_{0i} + \sum_{i=1}^{m-1} \sum_{j=i+1}^m v_{ij}$ 0/1-variables and $\sum_{i=1}^m \sum_{r=1}^{k_{0i}} l_r^{0i} + \sum_{i=1}^{m-1} \sum_{j=i+1}^m \sum_{r=1}^{k_{ij}} l_r^{ij}$ inequalities and at most further $\sum_{i=1}^m k_{0i} + \sum_{i=1}^{m-1} \sum_{j=i+1}^m k_{ij}$ inequalities are used.

The computation of the sets H_{0i} and H_{ij} can be done in analogy to the algorithm given in [8] to compute a hodograph.

4.3.2 Example

Let be given a region B and three objects T_1, T_2 and T_3 as shown in Fig. 3a). Figures 3b), 4 and 5 show the sets H_{01} (not connected), H_{02} (convex), H_{03} (non-convex), H_{12} (convex), H_{13} (non-convex) and H_{23} (non-convex).

It holds:

$$H_{01} = \left\{ (x, y) \in R^2 : \left[(h_{11}^{01}(x, y) = -x \leq 0) \wedge (h_{12}^{01}(x, y) = x + y - 2 \leq 0) \wedge (h_{13}^{01}(x, y) = x - 2y + 1 \leq 0) \right] \vee \left[(h_{21}^{01}(x, y) = -x + 4 \leq 0) \wedge (h_{22}^{01}(x, y) = x - 3 \leq 0) \wedge (h_{23}^{01}(x, y) = -x + y + 3 \leq 0) \wedge (h_{24}^{01}(x, y) = x - y - 3 \leq 0) \right] \right\}.$$




Figure 4 a) H_{02} b) H_{03}




Figure 5 a) H_{12} b) H_{13} c) H_{23}

Since $k_{01} = 2$ only one 0/1-variable u_1^{01} is to introduce. H_{02} is connected, moreover it is convex. An additional 0/1-variable is not necessary. For H_{03} it holds:

$$H_{03} = \left\{ (x, y) \in R^2 : \left[(h_{11}^{03}(x, y) = -x \leq 0) \wedge (h_{12}^{03}(x, y) = y - 2 \leq 0) \wedge (h_{13}^{03}(x, y) = x - 2y + 2 \leq 0) \right] \vee \left[(h_{21}^{03}(x, y) = -x + 2 \leq 0) \wedge (h_{22}^{03}(x, y) = x - 3 \leq 0) \wedge (h_{23}^{03}(x, y) = y - 2 \leq 0) \wedge (h_{24}^{03}(x, y) = -y + 2 \leq 0) \right] \vee \left[(h_{31}^{03}(x, y) = -x + 3 \leq 0) \wedge (h_{32}^{03}(x, y) = -x + y + 1 \leq 0) \wedge (h_{33}^{03}(x, y) = x - 4 \leq 0) \wedge (h_{34}^{03}(x, y) = x - y - 2 \leq 0) \right] \right\}.$$

Since $k_{03} = 3$, two 0/1-variables u_1^{03}, u_2^{03} are required. Hence, (26) has the following

form:

$$\begin{array}{llll} -x_1 & -M(1 - u_1^{01}) & \leq & M(1 - a_1), \\ x_1 + y_1 - 2 & -M(1 - u_1^{01}) & \leq & M(1 - a_1), \\ x_1 - 2y_1 + 1 & -M(1 - u_1^{01}) & \leq & M(1 - a_1), \\ -x_1 + 4 & -Mu_1^{01} & \leq & M(1 - a_1), \\ x_1 - 3 & -Mu_1^{01} & \leq & M(1 - a_1), \\ -x_1 + y_1 + 3 & -Mu_1^{01} & \leq & M(1 - a_1), \\ x_1 - y_1 - 3 & -Mu_1^{01} & \leq & M(1 - a_1), \\ \\ -x_2 & & \leq & M(1 - a_2), \\ x_2 - 1 & & \leq & M(1 - a_2), \\ y_2 - 3 & & \leq & M(1 - a_2), \\ -y_2 + 2 & & \leq & M(1 - a_2), \\ \\ -x_3 & -M(2 - u_1^{03} - u_2^{03}) & \leq & M(1 - a_3), \\ y_3 - 2 & -M(2 - u_1^{03} - u_2^{03}) & \leq & M(1 - a_3), \\ x_3 - 2y_3 + 2 & -M(2 - u_1^{03} - u_2^{03}) & \leq & M(1 - a_3), \\ -x_3 + 2 & -M(1 - u_1^{03} + u_2^{03}) & \leq & M(1 - a_3), \\ x_3 - 3 & -M(1 - u_1^{03} + u_2^{03}) & \leq & M(1 - a_3), \\ y_3 - 2 & -M(1 - u_1^{03} + u_2^{03}) & \leq & M(1 - a_3), \end{array}$$

$$\begin{array}{rcl}
-y_3 + 2 & -M(1 - u_1^{03} + u_2^{03}) & \leq M(1 - a_3), \\
-x_3 + y_3 + 1 & -M(1 + u_1^{03} - u_2^{03}) & \leq M(1 - a_3), \\
x_3 - y_3 - 2 & -M(1 + u_1^{03} - u_2^{03}) & \leq M(1 - a_3), \\
-x_3 + 3 & -M(1 + u_1^{03} - u_2^{03}) & \leq M(1 - a_3), \\
x_3 - 4 & -M(1 + u_1^{03} - u_2^{03}) & \leq M(1 - a_3), \\
& 2u_1^{03} + u_2^{03} & \geq 1.
\end{array}$$

For H_{12} it holds:

$$H_{12} = \left\{ (x, y) \in R^2 : (h_{11}^{12}(x, y) = -x - 5 \leq 0) \wedge (h_{21}^{12}(x, y) = y - 3 \leq 0) \wedge \right. \\
\left. (h_{31}^{12}(x, y) = x - 1 \leq 0) \wedge (h_{41}^{12}(x, y) = -y - 1 \leq 0) \right\}.$$

Since $k_{12} = 4$, two 0/1-variables u_1^{12}, u_2^{12} are necessary. Hence, (27) can be written as follows for $i = 1, j = 2$ (non-overlapping of $T_1(v_1)$ and $T_2(v_2)$):

$$\begin{array}{rcl}
-(x_2 - x_1) - 5 & +M(2 - u_1^{12} - u_2^{12}) & \geq 0 \\
(y_2 - y_1) - 3 & +M(1 - u_1^{12} + u_2^{12}) & \geq 0 \\
(x_2 - x_1) - 1 & +M(1 + u_1^{12} - u_2^{12}) & \geq 0 \\
-(y_2 - y_1) - 1 & +M(u_1^{12} + u_2^{12}) & \geq 0.
\end{array}$$

Furthermore, it holds:

$$H_{13} = \left\{ (x, y) \in R^2 : (h_{11}^{13}(x, y) = -x - 2 \leq 0) \wedge (h_{21}^{13}(x, y) = y - 3 \leq 0) \wedge \right. \\
\left. [(h_{31}^{13}(x, y) = x \leq 0) \vee (h_{32}^{13}(x, y) = y - 2 \leq 0)] \wedge (h_{41}^{13}(x, y) = x - 1 \leq 0) \wedge \right. \\
\left. (h_{51}^{13}(x, y) = x - y - 2 \leq 0) \wedge (h_{61}^{13}(x, y) = -y - 2 \leq 0) \right\}.$$

Because of $k_{13} = 6$, variables $u_1^{13}, u_2^{13}, u_3^{13} \in \{0, 1\}$ are to introduce. Hence, the non-overlapping of $T_1(v^1)$ and $T_3(v^3)$ can be described as follows (according to (27) one has $i = 1, j = 3$):

$$\begin{array}{rcl}
-(x_3 - x_1) - 2 & +M(3 - u_1^{13} - u_2^{13} - u_3^{13}) & \geq 0 \\
(y_3 - y_1) - 3 & +M(2 - u_1^{13} - u_2^{13} + u_3^{13}) & \geq 0 \\
x_3 - x_1 & +M(2 - u_1^{13} + u_2^{13} - u_3^{13}) & \geq 0 \\
(y_3 - y_1) - 2 & +M(2 - u_1^{13} + u_2^{13} - u_3^{13}) & \geq 0 \\
x_3 - x_1 - 1 & +M(1 - u_1^{13} + u_2^{13} + u_3^{13}) & \geq 0 \\
x_3 - x_1 - (y_3 - y_1) - 2 & +M(2 + u_1^{13} - u_2^{13} - u_3^{13}) & \geq 0 \\
-(y_3 - y_1) - 2 & +M(1 + u_1^{13} - u_2^{13} + u_3^{13}) & \geq 0 \\
& 4u_1^{13} + 2u_2^{13} + u_3^{13} & \geq 2.
\end{array}$$

In a similar way, the corresponding inequalities can be given to secure the non-overlapping of $T_2(v^2)$ and $T_3(v^3)$.

5. GENERALIZATION AND REMARKS

The generalization of the investigations for the two-dimensional case is possible to the three-dimensional case in a direct manner. Especially, the generalization of the packing of rectangular objects in section 4.1 yields the well-known container loading problem. The modeling of the packing problem of convex (three-dimensional) polyhedrons within a convex polyhedron can be done similarly to section 4.2.1 or to section 4.2.3 when using (three-dimensional) inner and outer hodographs.

The dimensions $n = 1, 2, 3$ are to be distinguished because of their practical relevance. Formally, packing problems of higher dimensions can also be defined and modelled as linear mixed-integer optimization problems with 0/1-variables.

In the investigations made only translations of the objects were allowed, but not rotations. Models for that more general case are not linear.

Acknowledgement

The authors would like to acknowledge the helpful comments of the anonymous referees.

References

- [1] Baker, B.S., Fortune, S.J. and Mahaney, S.R. (1986) Polygon containment under translation. *Journal of Algorithms* 7 pp. 532–548.
- [2] Beasley, J.E. (1985) An exact two-dimensional non-guillotine cutting tree search procedure. *Oper. Res.* 33 pp. 49–65.
- [3] Chen, C.S., Ram, B. and Sarin, S. (March 1989). *An Integer Programming Model for a Class of Assortment Problems*. Report, Dept. of Ind. Eng., North Carolina, A & T State Univ., Greensboro.
- [4] Dyckhoff, H. (1990). A typology of cutting and packing problems. *Europ. J. OR.* 44 pp. 145–160.

- [5] Dyckhoff, H., Wäscher, G. (1988). *Bibliography of Cutting and Packing*. Working Paper RWTH Aachen.
- [6] Stoyan, U.G., Gil, N.I. (1976). *Methods and Algorithms for the Arrangement of Planar Geometric Objects*. Kijev (russ).
- [7] Sweeney, P.E., and Ridenour, E.I. *Cutting and Packing Problems: A categorized application-oriented research bibliography*. Working Paper 610, School of Business Administration, University of Michigan.
- [8] Terno, J., Lindemann, R., Scheithauer, G. (1987). *Zuschnittprobleme und ihre praktische Lösung*. Verlag Harry Deutsch, Thun und Frankfurt/Main.