

Calculation of nmr parameters in paramagnetic metal-organic materials

Jure Lapajne

10. februar 2019

Presentation plan

- 1 MOFs = Metal—organic frameworks

Presentation plan

- 1 MOFs = Metal—organic frameworks
- 2 NMR — what and why

Presentation plan

- 1 MOFs = Metal—organic frameworks
- 2 NMR — what and why
- 3 NMR parameter calculation

Presentation plan

- 1 MOFs = Metal—organic frameworks
- 2 NMR — what and why
- 3 NMR parameter calculation
- 4 DFT


Presentation plan

- 1 MOFs = Metal—organic frameworks
- 2 NMR — what and why
- 3 NMR parameter calculation
- 4 DFT
- 5 Preliminary results

MOF = METAL-ORGANIC FRAMEWORK


MOF = METAL-ORGANIC FRAMEWORK

- crystal structure: central metallic ions + organic ligands




MOF = METAL-ORGANIC FRAMEWORK

- crystal structure: central metallic ions + organic ligands
- diversity: large number of possible combinations




MOF = METAL-ORGANIC FRAMEWORK

- crystal structure: central metallic ions + organic ligands
- diversity: large number of possible combinations
- wide usage: gas storage, clean energy applications, nonlinear optics, catalysts




MOF = METAL-ORGANIC FRAMEWORK

- crystal structure: central metallic ions + organic ligands
- diversity: large number of possible combinations
- wide usage: gas storage, clean energy applications, nonlinear optics, catalysts
- metallic ions commonly feature unpaired electrons





MOF = METAL-ORGANIC FRAMEWORK

- crystal structure: central metallic ions + organic ligands
- diversity: large number of possible combinations
- wide usage: gas storage, clean energy applications, nonlinear optics, catalysts
- metallic ions commonly feature unpaired electrons
- NMR spectra feature large shifts caused by unpaired electrons



NMR spectra of MOFs

- usual organic molecules display shifts in range $[-200, 200]$ ppm
- MOF spectra display large shifts rarely seen in purely organic molecules




Nuclei in external magnetic field

Nuclei in external magnetic field

- strong external magnetic field: several T


Nuclei in external magnetic field

- strong external magnetic field: several T
- nuclei with magnetic moment: lowest energy state splits




Nuclei in external magnetic field

- strong external magnetic field: several T
- nuclei with magnetic moment: lowest energy state splits
- two new states ΔE apart




Nuclei in external magnetic field

- strong external magnetic field: several T
- nuclei with magnetic moment: lowest energy state splits
- two new states ΔE apart
- radio frequency spectrum: excitations from low to high energy states



Nuclei in external magnetic field

- strong external magnetic field: several T
- nuclei with magnetic moment: lowest energy state splits
- two new states ΔE apart
- radio frequency spectrum: excitations from low to high energy states
- absorption peak at $\Delta E = \hbar\omega_{res}$
- ω_{res} depends on $B_{eff}(\text{observed nucleus})$



Effective magnetic field

Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

Effective magnetic field

Several parameters affect $B_{eff}(nucleus)$ and ω_{res} :

- electronic structure – shielding of external magnetic field

Effective magnetic field

Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

- electronic structure – shielding of external magnetic field
- spin–spin coupling to nearby nuclei and unpaired electrons

Effective magnetic field


Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

- electronic structure – shielding of external magnetic field
- spin–spin coupling to nearby nuclei and unpaired electrons
- unpaired electrons: large paramagnetic shifts

Effective magnetic field

Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

- electronic structure – shielding of external magnetic field
- spin–spin coupling to nearby nuclei and unpaired electrons
- unpaired electrons: large paramagnetic shifts
- calculation of nmr parameters: accurate knowledge of electronic wavefunction




- MOFs: metal-organic frameworks

Effective magnetic field

Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

- electronic structure – shielding of external magnetic field
- spin–spin coupling to nearby nuclei and unpaired electrons
- unpaired electrons: large paramagnetic shifts
- calculation of nmr parameters: accurate knowledge of electronic wavefunction




- MOFs: metal-organic frameworks
- transition metal atoms: unpaired electrons

Effective magnetic field

Several parameters affect $B_{\text{eff}}(\text{nucleus})$ and ω_{res} :

- electronic structure – shielding of external magnetic field
- spin–spin coupling to nearby nuclei and unpaired electrons
- unpaired electrons: large paramagnetic shifts
- calculation of nmr parameters: accurate knowledge of electronic wavefunction



- MOFs: metal-organic frameworks
- transition metal atoms: unpaired electrons

Chemical and hyperfine shifts

Chemical and hyperfine shifts

Definition

Total shift tensor $\underline{\underline{\sigma}}$ is defined by:

$$\vec{B}_{eff} = \vec{B}_0 \left(\underline{\underline{I}} - \underline{\underline{\sigma}} \right).$$

Two sources of chemical shifts:

Chemical and hyperfine shifts

Definition

Total shift tensor $\underline{\underline{\sigma}}$ is defined by:

$$\vec{B}_{eff} = \vec{B}_0 \left(\underline{\underline{I}} - \underline{\underline{\sigma}} \right).$$

Two sources of chemical shifts:

- electron density change caused by applied external magnetic field
- depends on electron density $n(\vec{r})$
- coupling between unpaired electron and observed nuclei
- depends on spin density $n_{\uparrow}(\vec{r}_{nuclei}) - n_{\downarrow}(\vec{r}_{nuclei})$

Calculation of NMR parameters:
accurate electronic wave function needed!

electronic wave function calculation

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed
- highly customizable - suitable for various molecules

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed
- highly customizable - suitable for various molecules

Alternatives

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed
- highly customizable - suitable for various molecules

Alternatives

- * Hartree—Fock

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed
- highly customizable - suitable for various molecules

Alternatives

- * Hartree—Fock
- * Quantum Monte Carlo

electronic wave function calculation

$$\hat{H} = T_n + T_e + W_{n-n} + W_{e-n} + W_{e-e} + V_{ext}$$

- Born-Oppenheimer approximation
- direct solution of coupled pde not feasible
- given accuracy level: time grows exponentially as a function of number of particles

Approximations

DFT

- most widely used
- good tradeoff between accuracy and speed
- highly customizable - suitable for various molecules


Alternatives

- * Hartree—Fock
- * Quantum Monte Carlo
- * Coupled-Cluster methods
- ⋮

density functional theory — DFT

N -particle problem:

- DFT effectively reduces N -particle problem to 1-particle problem
- Kohn-Sham theorems: transition from $3N$ to 3 coordinates.



System of N -particles:

- bijection between the set of external potentials and corresponding non-degenerate ground states
- bijection between the set of ground states and the set of ground states electron densities

energy as a functional of density

energy as a functional of density

Ground state $|\psi_0\rangle$:

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

Using bijection from previous slide, one can write:

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

Using bijection from previous slide, one can write:

$$|\psi_0\rangle = |\Psi_0[n_0(\vec{r})]\rangle,$$

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

Using bijection from previous slide, one can write:

$$|\psi_0\rangle = |\Psi_0[n_0(\vec{r})]\rangle,$$

and

$$E_0[n_0(\vec{r})] = \langle\Psi[n_0(\vec{r})]|\hat{H}|\Psi[n_0(\vec{r})]\rangle.$$

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

Using bijection from previous slide, one can write:

$$|\psi_0\rangle = |\Psi_0[n_0(\vec{r})]\rangle,$$

and

$$E_0[n_0(\vec{r})] = \langle\Psi[n_0(\vec{r})]|\hat{H}|\Psi[n_0(\vec{r})]\rangle.$$

Unfortunately the functional $\Psi[n_0(\vec{r})]$ not known!

energy as a functional of density

Ground state $|\psi_0\rangle$:

$$\hat{H}|\psi_0\rangle = E_0|\psi_0\rangle, \quad \text{with} \quad \langle\psi_0|\psi_0\rangle = 1.$$

Using bijection from previous slide, one can write:

$$|\psi_0\rangle = |\Psi_0[n_0(\vec{r})]\rangle,$$

and

$$E_0[n_0(\vec{r})] = \langle\Psi[n_0(\vec{r})]|\hat{H}|\Psi[n_0(\vec{r})]\rangle.$$

Unfortunately the functional $\Psi[n_0(\vec{r})]$ not known!

$E[n(\vec{r})]$ is modelled empirically.

Energy functional

$E[n(\vec{r})]$ should contain:

- * kinetic energy $T[n(\vec{r})]$
- * exchange – correlation term $E_{xc}[n(\vec{r})]$
- * coulomb interaction
- * external potential

$$E_H[n] = \frac{1}{2} \int \frac{n(\vec{r})n(\vec{r}')}{|\vec{r} - \vec{r}'|} d\vec{r}d\vec{r}'$$

$$V_{\text{ext}}[n] = \frac{1}{2} \int n(\vec{r})v_{\text{ext}}(\vec{r})d\vec{r}$$

Issues:

- kinetic energy term
- Calculation on exchange-correlation part

Solution:

- use of Slater determinant
- various approaches, no general rule/solution

Calculation procedure

Calculation procedure

Construct potential of nuclei and include it in $V_{ext}(\vec{r})$.

Calculation procedure

Construct potential of nuclei and include it in $V_{ext}(\vec{r})$.



Compute $E_H[n(\vec{r})]$, $E_{xc}[n(\vec{r})]$ and $E_{ext}[n(\vec{r})]$

Calculation procedure

Construct potential of nuclei and include it in $V_{ext}(\vec{r})$.



Compute $E_H[n(\vec{r})]$, $E_{xc}[n(\vec{r})]$ and $E_{ext}[n(\vec{r})]$



solve $(T[n(\vec{r})] + E_H[n(\vec{r})] + E_{xc}[n(\vec{r})] + E_{ext}[n(\vec{r})]) \psi_i = \epsilon_i \psi_i$

Calculation procedure

Construct potential of nuclei and include it in $V_{ext}(\vec{r})$.



Compute $E_H[n(\vec{r})]$, $E_{xc}[n(\vec{r})]$ and $E_{ext}[n(\vec{r})]$




solve $(T[n(\vec{r})] + E_H[n(\vec{r})] + E_{xc}[n(\vec{r})] + E_{ext}[n(\vec{r})]) \psi_i = \epsilon_i \psi_i$




compute $n(\vec{r}) = \sum_i \psi_i$, where $\epsilon_i < \epsilon_f$


Calculation procedure




Calculation procedure



Calculation procedure



Calculation procedure



DFT flexibility

Large number of various exchange-correlation functionals:

- hybrid: GGA + hartree-fock exchange

$$-\frac{1}{2} \sum_{k,l} \int \frac{\phi_k^*(\vec{r}, \sigma) \phi_l(\vec{r}, \sigma) \phi_l^*(\vec{r}', \sigma') \phi_k(\vec{r}', \sigma')}{|\vec{r} - \vec{r}'|} d\vec{r} d\vec{r}'$$


- meta-GGA: higher order derivatives

- GGA:

$$E_{xc}^{gga} = \int n(\vec{r})^{4/3} F(|\nabla n(\vec{r})|/n(\vec{r})^{4/3})$$

- LDA: uniform gas

$$E_{xc}^{lda}[n] = -C \int n(\vec{r})^{4/3} d\vec{r}$$



My work

My work

- various functionals,

My work



- various functionals,
- various basis sets,

My work

- various functionals,
- various basis sets,
- boundary conditions: periodic vs single molecule,

My work

- various functionals,
- various basis sets,
- boundary conditions: periodic vs single molecule,
- full electron dft vs frozen core dft



Conclusion

- 1 investigation of nmr