

Machine Learning HW8

Student ID: 111652015

Problem 1

Definition (Notes on 10/17)

$$L_{SSM}(\theta) = \mathbb{E}_{x \sim p(x)}[\|S(x; \theta)\|^2] + \mathbb{E}_{x \sim p(x)} \mathbb{E}_{v \sim p(v)} \left[2 v^\top \nabla_x (v^\top S(x; \theta)) \right].$$

Goal Show that

$$L_{SSM}(\theta) = \mathbb{E}_{x \sim p(x)} \mathbb{E}_{v \sim p(v)} \left[\|v^\top S(x; \theta)\|^2 + 2 v^\top \nabla_x (v^\top S(x; \theta)) \right].$$

Lemma Assume $\mathbb{E}_v[vv^\top] = I$. Then for any vector a ,

$$\mathbb{E}_v[(v^\top a)^2] = a^\top \mathbb{E}_v[vv^\top] a = \|a\|^2.$$

Proof Apply the identity with $a = S(x; \theta)$:

$$\mathbb{E}_v[\|v^\top S(x; \theta)\|^2] = \|S(x; \theta)\|^2.$$

Taking \mathbb{E}_x on both sides and substituting into the given definition yields

$$\begin{aligned} L_{SSM}(\theta) &= \mathbb{E}_x \mathbb{E}_v[\|v^\top S(x; \theta)\|^2] + \mathbb{E}_x \mathbb{E}_v[2 v^\top \nabla_x (v^\top S(x; \theta))] \\ &= \mathbb{E}_x \mathbb{E}_v \left[\|v^\top S(x; \theta)\|^2 + 2 v^\top \nabla_x (v^\top S(x; \theta)) \right]. \end{aligned}$$

Proof of Lemma Note that

$$(v^\top a)^2 = (a^\top v)(v^\top a) = a^\top (vv^\top) a.$$

Taking expectations and using linearity,

$$\mathbb{E}[(v^\top a)^2] = \mathbb{E} \left[a^\top (vv^\top) a \right] = a^\top \mathbb{E}[vv^\top] a = a^\top I a = \|a\|^2.$$

Problem 2 (Briefly explain SDE)

Stochastic Differential Equation (SDE) A noisy dynamical system can be modeled by the SDE

$$dx_t = f(x_t, t) dt + G(x_t, t) dW_t, \quad x(0) = x_0. \quad (1)$$

Here $x_t \in R^d$ is the unknown stochastic process; W_t is a Brownian motion ; f is the drift that determines the average trend , and G is the diffusion matrix that injects noise into the system.

Itô integral form Equation (1) is equivalent to the Itô integral equation

$$x_t = x_0 + \int_0^t f(x_s, s) ds + \int_0^t G(x_s, s) dW_s. \quad (2)$$

Definition of the Itô integral The stochastic integral in (2) is defined as the mean-square limit

$$\int_0^t G(x_s, s) dW_s = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} G(x(t_k), t_k) [W(t_{k+1}) - W(t_k)],$$

for partitions $0 = t_0 < t_1 < \dots < t_n = t$, where the increments satisfy

$$W(t_{k+1}) - W(t_k) \sim \mathcal{N}(0, (t_{k+1} - t_k)I),$$

and increments over disjoint intervals are independent.

Existence and Uniqueness (Common Conditions) f, G globally Lipschitz + linear growth \Rightarrow unique strong solution.

Euler–Maruyama Method Let the step size be $\Delta t = T/N$ and define grid points $t_n = n \Delta t$ for $n = 0, 1, \dots, N$. Set the initial value $X_0 = x_0$. The Euler–Maruyama iteration is

$$X_{n+1} = X_n + f(X_n, t_n) \Delta t + G(X_n, t_n) \Delta W_n, \quad n = 0, 1, \dots, N-1,$$

where the noise increments satisfy

$$\Delta W_n \sim \mathcal{N}(0, \Delta t I), \quad \text{independently for each } n.$$

Problem 3 Unanswered Questions

1. What is the difference between a strong solution and a weak solution of an SDE?
2. Does the SDE have an analytic (closed-form) solution?