

Product Datasheet | H2-Db easYmer® kit

Application	easYmers® are peptide loadable HLA class I monomers.
-------------	--

easYmers® can be used to generate peptide-HLA monomers with your choice of peptide. The monomers can easily be tetramerized with fluorophore conjugated streptavidin and used to analyse T cells by flow cytometry. The easYmer reagent can also be used to evaluate specific

peptide-HLA I binding.

Catalog no. 5001-01

Size	Test	Vol.	Folding Buffer	Control Peptide
	20	40 μl	1 vial (0.5 ml)	1 vial
	50	100 μΙ	1 vial (1 ml)	1 vial
	150	300 μl	1 vial (1 ml)	2 vials
	500	1000 µl	1 vial (3 ml)	3 vials

Allotype H2-Db

peptide receptive, biotinylated in Tris/Maleate pH 7, 30% Glycerol

Storage Store at -20°C. Avoid repeated freze-thaw cycles.

Shelf life easYmers® are stable at -20°C for 12 months.

See expiration on vial

Folding buffer Tris/Maleate pH 7

Control Peptide ASNENMETM (20 nmol lyophilized peptide per vial)

H2-Db binder and positive control for evaluation of peptide-HLA folding. Dissolve each peptide vial in 20 μ l DMSO resulting in a peptide

conc.of 1 mM. Store at -20°C

Peptide source Influenza NP366-374

For Research Use Only (RUO)