Neonatal Oxygen Management and the Cord Transition: An Extension of the Gestational Oxygen-Timing Alignment Framework

Extension Research Track 5

Author: Jay Andrew Carpenter **Affiliation:** Independent Researcher, Sydney, Australia **Contact:** jay.andrew.carpenter@gmail.com | jay@secs.observer **Date:** March 2026 **License:** CC-BY 4.0 International

Extends: Carpenter, J.A. (2026). *Gestational Oxygen-Timing Alignment: A Unifying Framework Linking Placental Insufficiency, Oligodendrocyte Progenitor Vulnerability, and Neurodevelopmental Outcome.* medRxiv preprint.

Abstract

The gestational oxygen-timing alignment model establishes that placental insufficiency during critical oligodendrocyte progenitor cell (OPC) maturation windows (gestational weeks 23–32) produces permanent white matter injury patterns determining neurodevelopmental outcome. This extension investigates the **neonatal transition period** — the moment of cord clamping and the hours-to-weeks that follow — as the final modifiable intervention window before hypoxic white matter damage becomes irreversible. The cord cut moment represents a physiological inflection point: the switch from placental to pulmonary gas exchange. For infants already compromised by in utero placental insufficiency, this transition constitutes a potential **second hypoxic insult** during an already-vulnerable developmental period. This document synthesises evidence from delayed cord clamping trials, neonatal resuscitation oxygen protocols, NICU oxygen targeting studies (SUPPORT, BOOST II, COT, NEOPROM), cerebral near-infrared spectroscopy (NIRS) monitoring, and caffeine therapy research to construct an integrated framework for neonatal oxygen management optimised for OPC survival. The analysis extends to multi-parity severity effects — explaining why successive children of mothers with recurrent preeclampsia may exhibit progressively worse neurodevelopmental outcomes — and proposes a novel connection between gestational OPC pool depletion and the characteristic memory patterns observed in age-related dementia. A preeclampsia-flagged neonatal pathway with enhanced oxygenation monitoring is proposed as a practical clinical translation of the alignment model. Seventy-eight references are cited spanning neonatology, placental biology, oligodendrocyte neuroscience, and neurodegenerative disease.

---

1. Introduction

1.1 The Final Intervention Window

The gestational oxygen-timing alignment model positions placental insufficiency as the upstream determinant of neurodevelopmental injury through disruption of fetal oxygenation during critical periods of oligodendrocyte progenitor cell (OPC) maturation [1]. The published synthesis framework identifies gestational weeks 23–32 as the primary vulnerability window, during which OPCs are actively proliferating, migrating, and initiating differentiation programs that are exquisitely sensitive to hypoxic disruption [2, 3].

What the core model establishes as the prenatal mechanism, however, does not terminate at birth. The transition from intrauterine to extrauterine life represents the most dramatic physiological shift a human being ever undergoes — and the period surrounding cord clamping and the initiation of pulmonary respiration constitutes the **last modifiable intervention window** before white matter injury patterns become permanent [4].

For the healthy term infant born after an uncomplicated pregnancy, this transition is a well-orchestrated physiological process. For the preterm infant — particularly one whose placenta has already been failing — the transition represents a critical juncture where existing vulnerability can be either mitigated or catastrophically compounded [5].

1.2 The Double-Hit Hypothesis

The present extension proposes a **double-hit model** of neonatal white matter injury:

- **Hit 1 (Prenatal):** Placental insufficiency from preeclampsia or related conditions → chronic fetal hypoxia → OPC maturation arrest during gestational vulnerability windows
- **Hit 2 (Perinatal/Postnatal):** Suboptimal oxygen management during and after birth → acute-on-chronic hypoxic injury to already-compromised OPC populations

This double-hit framework explains why two infants born at the same gestational age after preeclamptic pregnancies can have markedly different neurodevelopmental outcomes: the quality of perinatal oxygen management determines whether the second hit completes the injury or allows partial rescue [6, 7].

1.3 Scope

This document examines:

1. The physiology of the cord clamping transition and its implications for OPC survival
2. Evidence for delayed cord clamping as a neuroprotective strategy
3. Neonatal resuscitation oxygen management and the FiO_2 debate
4. Ongoing NICU oxygen targeting and its relationship to white matter outcomes
5. Cerebral monitoring technologies for real-time neuroprotection
6. Risk stratification and the concept of developmental clearance
7. Multi-parity effects on successive offspring
8. The lifetime implications of OPC pool depletion for age-related neurodegeneration

---

2. The Cord Clamping Transition: Physiology and Vulnerability

2.1 Placental Gas Exchange: The In Utero Baseline

Throughout gestation, the placenta serves as the fetal lung. Oxygen transfer across the villous membrane is driven by the maternal-fetal oxygen gradient, with maternal arterial pO_2 of approximately 100 mmHg producing an umbilical venous pO_2 of approximately 28–35 mmHg in the healthy pregnancy [8]. This seemingly low oxygen tension is adequate for normal fetal development because fetal haemoglobin (HbF) has a left-shifted oxygen-dissociation curve relative to adult haemoglobin, allowing efficient oxygen extraction at low partial pressures [9].

In preeclampsia, this system is already compromised. Deficient spiral artery remodelling reduces utero-placental blood flow, and the anti-angiogenic imbalance (elevated sFlt-1, reduced PlGF) impairs villous vascular development [10, 11]. The result is chronic fetal hypoxaemia — the "first hit" of the alignment model — with umbilical venous pO_2 potentially reduced to 20–25 mmHg or lower [12].

2.2 The Moment of Cord Clamping

When the umbilical cord is clamped and cut, the placental oxygen supply is abruptly terminated. In the physiological sequence, this should coincide with the establishment of pulmonary gas exchange through the infant's first breaths. However, the temporal coupling between cord clamping and the onset of effective ventilation is not guaranteed — and in the compromised preterm infant, it frequently fails [13].

The critical vulnerability period is the interval between the cessation of placental gas exchange and the establishment of adequate pulmonary ventilation. During this interval, the infant receives oxygen from **neither** source. Even in the healthy term infant, this transition produces a transient fall in oxygen saturation, with SpO_2 values of 60–65% at 1 minute of life being normal [14]. For the already-hypoxaemic preterm infant born after placental insufficiency, this nadir represents an acute hypoxic insult superimposed on chronic hypoxic exposure.

2.3 Haemodynamic Consequences

Cord clamping also produces immediate haemodynamic changes relevant to cerebral oxygen delivery:

Preload reduction: Clamping the umbilical vein eliminates the low-resistance placental vascular bed from the circulation, abruptly increasing systemic vascular resistance (SVR) and reducing venous return to the heart [15]. In the preterm infant, whose myocardium has limited contractile reserve, this can produce a transient reduction in cardiac output.

Ductal shunting: The ductus arteriosus remains patent in the first hours of life. The sudden increase in SVR relative to pulmonary vascular resistance (PVR) can shift ductal shunting from right-to-left to left-to-right, potentially producing a steal phenomenon that reduces systemic (including cerebral) blood flow [16].

Cerebral autoregulation immaturity: Preterm infants have limited cerebral autoregulatory capacity, meaning cerebral blood flow is pressure-passive across a significant range of mean arterial pressures [17]. Haemodynamic instability during the cord transition therefore translates directly into cerebral oxygen delivery fluctuations.

2.4 Implications for OPC Populations

The OPC populations that have survived the prenatal "first hit" of placental insufficiency are already in a metabolically stressed state. These cells have adapted to chronic hypoxia through upregulation of HIF-1α signalling pathways, which maintain cell survival but arrest differentiation into mature myelinating oligodendrocytes [18, 19].

The cord transition produces a paradoxical challenge for these cells:

1. **Transient deepening of hypoxia** during the ventilation gap may push surviving OPCs past their tolerance threshold
2. **Abrupt reoxygenation** once ventilation is established can trigger oxidative stress through generation of reactive oxygen species (ROS), which are directly toxic to OPCs [20]
3. **Haemodynamic instability** produces fluctuating cerebral oxygen delivery, and cycling between hypoxia and relative hyperoxia is more damaging to white matter than sustained mild hypoxia [21]

This establishes the cord clamping transition as fundamentally relevant to the alignment model: it is the moment where the prenatal injury can be either stabilised or completed.

---

3. Delayed Cord Clamping: Evidence and Mechanism

3.1 Physiological Rationale

Delayed cord clamping (DCC) — defined as deferring cord clamping for at least 30–60 seconds after birth, or until cord pulsation ceases — allows continued placental-fetal blood transfer (placental transfusion) during the period when the infant is establishing pulmonary ventilation [22]. The physiological logic is straightforward: rather than simultaneously terminating placental oxygen supply and demanding

that pulmonary gas exchange immediately compensate, DCC allows a graduated transition where both systems operate in parallel.

In the preterm infant, placental transfusion provides approximately 10–30 mL/kg of additional blood volume over 60–180 seconds [23]. This blood transfer carries several potential neuroprotective components:

- **Red blood cells:** Additional haemoglobin mass for oxygen carrying capacity during the transition period
- **Plasma volume:** Improved circulating volume supporting cardiac output and cerebral perfusion pressure
- **Stem cells:** Haematopoietic and mesenchymal stem cells of placental origin with potential neurotrophic properties [24]
- **Immunoglobulins:** Passive immune factors that may modulate the inflammatory response
- **Iron stores:** Critical for ongoing haemoglobin synthesis and neurodevelopmental processes (iron is essential for oligodendrocyte metabolism and myelin synthesis) [25]

3.2 Clinical Evidence for Neuroprotection

The evidence supporting DCC in preterm infants has accumulated substantially over the past two decades.

The Australian Placental Transfusion Study (APTS): This landmark multicentre trial randomised 1,566 infants born before 30 weeks' gestation to DCC (≥60 seconds) versus immediate cord clamping (ICC, <10 seconds). While the primary outcome of death or major morbidity at 36 weeks did not reach statistical significance, DCC was associated with reduced mortality (6.4% vs 9.0%, p=0.03) in the intention-to-treat analysis [26]. A pre-specified subgroup analysis demonstrated that the mortality benefit was concentrated in infants born at 24–27 weeks — the gestational ages corresponding to the peak OPC vulnerability window identified in the alignment model.

Cochrane Systematic Review (2024 update): The most recent Cochrane analysis of 48 trials (6,967 infants) concluded that DCC ≥30 seconds for preterm infants reduces hospital mortality (RR 0.73, 95% CI 0.56–0.95) and reduces the need for blood transfusion in the first 6 weeks (RR 0.65, 95% CI 0.54–0.78) [27]. Importantly, DCC was not associated with increased risks of intraventricular haemorrhage (IVH) or hypothermia.

Andersson et al. (2015): A follow-up study of term infants randomised to DCC ≥180 seconds versus ICC found improved fine motor and social domain scores at 4 years of age, particularly in boys [28]. While these were term infants, the finding suggests that even in uncomplicated pregnancies, the cord transition carries neurodevelopmental significance.

Mercer et al. (2016): An RCT of 208 infants born at 24–31 weeks found that DCC (30–45 seconds) was associated with higher Motor Composite scores on the Bayley-III at 18–22 months corrected age, with the strongest effect in male infants [29].

3.3 DCC in Preeclamptic Pregnancies

A critical gap in the DCC literature is the relative paucity of data specifically examining DCC outcomes in preeclamptic pregnancies. This is a significant limitation because preeclampsia creates a unique physiological context:

The compromised placenta as a limiting factor: In severe preeclampsia, the placental vasculature is already impaired. The assumption that DCC will provide adequate placental transfusion may not hold when the uteroplacental circulation is restricted by deficient spiral artery remodelling [30]. Placental blood flow in severe preeclampsia may be reduced by 50% or more, meaning that the "transfusion volume" available during DCC is significantly less than in uncomplicated pregnancies.

Potential counterargument — quality over quantity: Even if the volume of placental transfusion is reduced in preeclampsia, the continued oxygen supply during the ventilation establishment period may be proportionally MORE important for the already-hypoxaemic infant. Any supplemental oxygen delivery during the transition gap, however modest, represents a relative improvement over the zero-oxygen state produced by immediate clamping.

Placental anti-angiogenic factors: The preeclamptic placenta releases elevated levels of sFlt-1 and soluble endoglin into the fetal circulation [31]. It is theoretically possible that prolonged cord clamping delay in the context of preeclampsia could increase fetal exposure to these circulating anti-angiogenic factors. However, the half-life of sFlt-1 is approximately 4-6 hours, meaning that a 60–120 second extension of exposure is biologically negligible relative to the accumulated prenatal exposure.

Clinical urgency: Preeclampsia deliveries frequently occur in the context of clinical urgency (emergency caesarean section for maternal or fetal distress), and the obstetric team's focus on maternal stabilisation may override DCC protocols. This is a practical barrier to implementation that must be addressed in clinical pathway design [32].

3.4 Umbilical Cord Milking

Umbilical cord milking (UCM) — the manual expression of blood from the cord toward the infant — has been investigated as an alternative to DCC, particularly in emergency situations where DCC is impractical [33].

Katheria et al. (2019): The PREMOD2 trial randomised 474 infants born at 23–31 weeks to UCM versus DCC ≥60 seconds. The trial was stopped early for the 23–27 week subgroup due to a higher rate of severe IVH in the UCM arm (22% vs 6%, $p<0.001$) [34]. This finding is directly relevant to the alignment model: cord milking delivers a rapid bolus of blood that may cause cerebral haemodynamic instability, whereas DCC provides a gradual, physiological transfusion. In the very preterm infant with immature cerebral autoregulation, the distinction matters profoundly.

Implication for OPC protection: The PREMOD2 finding suggests that the manner of placental blood transfer is as important as the volume. Gentle, physiological transfusion (DCC) supports cerebral haemodynamic stability, while rapid bolus transfusion (UCM) may cause pressure surges in the fragile germinal matrix vasculature. Since IVH and periventricular white matter injury share anatomical and pathophysiological overlap, UCM-associated IVH directly threatens the OPC populations the alignment model identifies as vulnerable [35].

3.5 Physiological-Based Cord Clamping

An emerging approach — **physiological-based cord clamping (PBCC)** — delays cord clamping specifically until the establishment of spontaneous respiration, rather than for a fixed time interval [36]. This approach directly addresses the "ventilation gap" identified in Section 2.2.

Fogarty et al. (2018): A feasibility trial of PBCC in preterm infants demonstrated that clamping the cord only after the onset of regular breathing resulted in smoother cardiovascular and respiratory transitions, with higher SpO_2 at 5 minutes and more stable cerebral oxygenation measured by NIRS [37]. The physiological logic is compelling: rather than imposing an arbitrary time delay, PBCC ensures that pulmonary gas exchange is at least partially established before placental supply is discontinued.

Relevance to preeclampsia-exposed infants: PBCC may be particularly beneficial for infants born after placental insufficiency, who have reduced physiological reserve to tolerate the ventilation gap. However, these same infants are more likely to require active resuscitation at birth, creating a tension between waiting for spontaneous respiration and initiating assisted ventilation.

---

4. Neonatal Resuscitation: The FiO_2 Debate

4.1 Historical Context

The history of neonatal resuscitation oxygen management is a cautionary tale about the harm that can result from well-intentioned intervention. For decades, 100% oxygen was considered the standard for neonatal resuscitation, based on the assumption that maximal oxygen delivery could only be beneficial for a hypoxaemic newborn [38].

This assumption was fundamentally challenged by the work of Saugstad and colleagues, who demonstrated in the 1990s and 2000s that high-concentration oxygen resuscitation produced oxidative stress, paradoxically worsening outcomes in asphyxiated neonates [39, 40]. The recognition that supplemental oxygen is a double-edged sword — capable of rescuing from hypoxia but also of inflicting oxidative damage — has profound implications for the alignment model's framework of OPC vulnerability.

4.2 Current Guidelines

The International Liaison Committee on Resuscitation (ILCOR) and the Neonatal Resuscitation Program (NRP) now recommend [41, 42]:

- **Term infants ≥35 weeks:** Initiate resuscitation with 21% FiO_2 (room air)
- **Preterm infants <35 weeks:** Initiate resuscitation with 21–30% FiO_2
- **Titrate FiO_2** based on pulse oximetry targeting pre-ductal SpO_2 values that approximate the normal postnatal transition

The target SpO_2 values during the first 10 minutes of life are based on normative nomograms derived from healthy term and near-term infants [43]:

Time after birth	Target SpO_2 range
1 minute	60–65%
2 minutes	65–70%
3 minutes	70–75%
4 minutes	75–80%
5 minutes	80–85%
10 minutes	85–95%

4.3 OPC-Specific Considerations in Oxygen Targeting

The standard SpO_2 targets were derived from observations of healthy newborns in the first minutes of life [43, 44]. They represent the normal physiological transition and are applied uniformly to all neonates. However, the alignment model raises a critical question: **should infants born after confirmed placental insufficiency have different oxygen targets during resuscitation?**

The rationale for modified targets rests on the metabolic state of OPCs in the hypoxia-exposed fetus:

Chronic hypoxia adaptation: OPCs that have survived prenatal hypoxia have upregulated HIF-1α and shifted toward glycolytic metabolism [18]. These cells are adapted to a low-oxygen environment. Abrupt exposure to high oxygen concentrations produces a burst of reactive oxygen species through mitochondrial complex dysfunction, directly damaging OPC membranes, DNA, and differentiation-critical transcription factors [20, 45].

OPC sensitivity to ROS: Pre-oligodendrocytes (late OPCs, the predominant oligodendroglial cell in the preterm brain) are exceptionally vulnerable to oxidative stress. They have relatively low levels of

antioxidant enzymes (superoxide dismutase, glutathione peroxidase, catalase) compared to mature oligodendrocytes or astrocytes [46, 47]. The maturation arrest imposed by chronic hypoxia keeps these cells locked in their most oxidant-vulnerable state.

The reoxygenation paradox: For the hypoxia-adapted OPC, the resumption of normal oxygenation is itself a form of oxidative stress. This is analogous to ischaemia-reperfusion injury in stroke: the return of blood flow (and oxygen) to ischaemic tissue produces a burst of oxidative damage that can exceed the injury from the ischaemic period itself [48].

4.4 The Goldilocks Zone

The alignment model thus predicts a **narrow optimal oxygenation window** for resuscitation of the preeclampsia-exposed preterm infant:

- **Too little oxygen (persistent hypoxia):** Deepens OPC maturation arrest, cell death through energy failure
- **Too much oxygen (hyperoxia/rapid reoxygenation):** Oxidative stress-mediated OPC injury, exacerbated by low antioxidant capacity
- **Optimal ("Goldilocks"):** Gradual restoration of oxygenation that supports cellular metabolism without triggering oxidative cascades

This Goldilocks zone may be narrower for preeclampsia-exposed infants than for those born preterm without placental insufficiency, implying that current one-size-fits-all SpO_2 targets may be inadequate for the highest-risk population.

4.5 Evidence from Oxygenation Studies in Preterm Resuscitation

Vento et al. (2009): A randomised trial of 78 preterm infants (24–28 weeks) compared resuscitation with 30% vs 90% FiO_2. The low-oxygen group had significantly lower markers of oxidative stress (oxidised glutathione, 8-isoprostanes) and reduced rates of bronchopulmonary dysplasia [49]. While white matter outcomes were not specifically reported, the reduction in systemic oxidative markers implies reduced oxidative exposure for cerebral OPCs.

Rabi et al. (2011): The To2rpido trial compared 100% vs 21% FiO_2 for delivery room resuscitation of term and near-term infants with birth depression. Room air resuscitation was associated with faster time to first breath and reduced mortality [50].

Oei et al. (2017): A meta-analysis of individual patient data from 768 preterm infants (<32 weeks) found that initial high FiO_2 (>0.6) was associated with increased mortality, while the lowest mortality was observed with initial FiO_2 of 0.21–0.30 [51].

These studies consistently demonstrate that lower initial FiO_2 produces better outcomes in preterm infants, supporting the Goldilocks zone concept for OPC survival during the cord transition.

---

5. NICU Oxygen Targeting: The Ongoing Challenge

5.1 The Great Oxygen Targeting Debate

Once the acute resuscitation period is complete, the preterm infant enters a phase of ongoing oxygen management in the NICU that may last weeks to months. The optimal SpO_2 target range for very preterm infants has been one of the most extensively studied and bitterly debated questions in neonatology [52].

5.2 The NeOProM Collaboration

Five large randomised controlled trials — collectively enrolling nearly 5,000 extremely preterm infants — compared lower (85–89%) versus higher (91–95%) SpO_2 targeting:

SUPPORT (USA, 2010): 1,316 infants, 24–27 weeks. Lower target associated with increased mortality (19.9% vs 16.2%), no difference in severe retinopathy [53].

BOOST II (UK/Australia/New Zealand, 2013): 2,448 infants, <28 weeks. Lower target associated with increased mortality (23.1% vs 15.9% in UK; similar trends in Australia/NZ). Trial stopped early for the UK cohort due to safety concerns [54].

COT (Canada, 2013): 1,201 infants, 23–27 weeks. Non-significant trend toward increased death or disability in the lower target group [55].

NEOPROM meta-analysis (Askie et al., 2018): Individual patient data meta-analysis of all five trials (4,965 infants) concluded that the lower SpO_2 target (85–89%) was associated with significantly higher mortality (RR 1.17, 95% CI 1.04–1.31) and higher rates of necrotising enterocolitis, while the higher target (91–95%) was associated with higher rates of severe retinopathy of prematurity [56].

5.3 Interpretation Through the Alignment Model Lens

The NeOProM results take on additional significance when viewed through the OPC vulnerability framework:

The mortality signal: The increased mortality in the lower SpO_2 target arm is consistent with the prediction that sustained marginal oxygenation pushes already-compromised OPC populations past their survival threshold. While OPCs are adapted to chronic mild hypoxia in utero, the postnatal context is fundamentally different: the placenta is gone, and the infant's lungs are the sole oxygen source. Sustained SpO_2 of 85–89% in the postnatal period may not provide the same tissue oxygenation as an equivalent saturation in the fetal circulation (where HbF provides enhanced oxygen extraction) [57].

White matter outcomes in NeOProM: Surprisingly, the NeOProM meta-analysis did not find a significant difference in white matter injury (as assessed by cranial ultrasound) between the two SpO_2 target groups. However, this may reflect the crude sensitivity of cranial ultrasound for detecting diffuse white matter injury — MRI studies consistently demonstrate white matter abnormalities not visible on ultrasound [58].

The missing variable — fluctuation: The NeOProM trials compared TARGET ranges, but what may matter more for OPC survival is the **stability of oxygenation within whatever range is targeted.** Oxygen fluctuation — cycling between hypoxia and relative hyperoxia — is potentially more damaging than sustained exposure to either extreme [21, 59].

5.4 Intermittent Hypoxia Episodes (IHE)

Preterm infants in the NICU experience frequent intermittent hypoxia episodes (IHE) — transient falls in SpO_2 below 80% lasting 10 seconds or more. These episodes are extremely common: very preterm infants may experience hundreds of IHE per week, particularly in the first weeks of life [60, 61].

Cumulative burden: Martin et al. (2011) demonstrated that the cumulative burden of IHE in the first weeks of life is associated with adverse neurodevelopmental outcomes at 18 months, independent of gestational age and other confounders [62]. The relationship was dose-dependent: more IHE episodes predicted worse outcomes.

Mechanism — cycling oxidative stress: Each IHE followed by reoxygenation constitutes a miniature ischaemia-reperfusion event in the brain. For OPCs already in a vulnerable state, the cumulative effect of hundreds of such cycles may be more damaging than a single sustained hypoxic episode of equivalent

total duration [63]. Each cycle generates a pulse of ROS that damages OPC membranes and disrupts differentiation signalling.

Apnoea of prematurity as a mediator: IHE in preterm infants are primarily caused by apnoea of prematurity — the immature respiratory control that produces frequent pauses in breathing [64]. This is directly relevant to the alignment model: apnoea of prematurity is a consequence of brainstem immaturity, and its severity may itself be related to gestational hypoxic exposure (hypoxia impairs the development of brainstem respiratory centres, creating a self-reinforcing cycle of apnoea → IHE → white matter injury → impaired brainstem development → more apnoea) [65].

5.5 Caffeine as Indirect Neuroprotection

Caffeine citrate therapy for apnoea of prematurity has emerged as one of the most effective neonatal interventions, with substantial evidence of neuroprotective benefit that extends well beyond its respiratory stimulant effect.

The CAP Trial (Schmidt et al., 2006, 2007, 2012): The Caffeine for Apnea of Prematurity trial randomised 2,006 very low birth weight infants to caffeine citrate versus placebo. Caffeine therapy reduced the rate of bronchopulmonary dysplasia, reduced the duration of positive pressure ventilation, and — most remarkably — improved neurodevelopmental outcomes at 18 months corrected age, with reduced rates of cerebral palsy and cognitive delay [66, 67]. At 5-year follow-up, the motor benefit persisted, and gross motor function was significantly better in the caffeine group [68].

Mechanism through the alignment model: Caffeine's neuroprotective effect can be understood as indirect OPC protection through multiple pathways:

1. **Reduced IHE burden:** By stimulating respiratory drive, caffeine reduces the frequency and severity of apnoeic episodes and associated IHE, reducing cumulative cycling oxidative stress to white matter [69]
2. **Adenosine A_2 receptor antagonism:** Caffeine blocks adenosine receptors, which modulates neuroinflammation. During hypoxia, extracellular adenosine levels rise dramatically, and adenosine A_2A receptor activation on microglia promotes pro-inflammatory cytokine release that is toxic to OPCs [70]
3. **Direct antioxidant properties:** Caffeine has been shown to scavenge free radicals and reduce oxidative stress markers in preterm infants, potentially protecting OPCs from ROS-mediated injury during reoxygenation events [71]
4. **Timing of initiation:** Retrospective analyses suggest that early caffeine initiation (within the first 2 days of life) is associated with better outcomes than later initiation, consistent with the alignment model's emphasis on time-critical intervention during OPC vulnerability windows [72]

5.6 The Case for Continuous Monitoring

Current NICU practice monitors oxygen saturation continuously via pulse oximetry, but the clinical response to desaturation events is often reactive rather than proactive. The alignment model supports a paradigm shift toward:

1. **Prospective risk stratification:** Infants born after confirmed placental insufficiency should be flagged for enhanced monitoring protocols from admission
2. **Automated oxygen controllers:** Devices that automatically titrate FiO_2 in response to SpO_2 changes have been shown to reduce time outside target range and, critically, to reduce the frequency of oxygen fluctuations [73]
3. **Cerebral oxygenation monitoring (NIRS):** Pulse oximetry measures arterial oxygen saturation, but cannot determine cerebral tissue oxygenation. Near-infrared spectroscopy provides continuous

measurement of regional cerebral oxygen saturation ($rcSO_2$), which is the variable that directly determines OPC oxygen supply [74] (see Section 6)

4. **Duration of monitoring:** Rather than discontinuing intensive monitoring at an arbitrary postmenstrual age, the alignment model supports continued monitoring until developmental milestones indicate that the OPC vulnerability window has closed (see Section 7)

---

6. Sampling, Measurement, and Monitoring

6.1 Cord Blood Gas Analysis

Paired umbilical cord blood gases (arterial and venous) obtained at delivery provide the most direct available assessment of in utero oxygenation immediately preceding birth [75].

What cord gases reveal about OPC exposure:

Parameter	Normal range	Significance for alignment model
UA pH	7.18–7.38	Acidaemia reflects sustained hypoxia prior to delivery
UA pO_2	10–25 mmHg	Directly measures fetal tissue oxygenation
UA pCO_2	40–55 mmHg	Elevated in respiratory acidosis (acute hypoxia)
UA Base excess	-2 to -8 mEq/L	Severity of metabolic acidosis (chronic hypoxia duration)
UA Lactate	<3.5 mmol/L	Marker of anaerobic metabolism under hypoxic conditions

Interpretation for risk stratification: An umbilical arterial pH <7.10, base excess more negative than -12 mEq/L, or lactate >6 mmol/L in the context of a preeclamptic pregnancy indicates significant in utero hypoxic stress and should trigger enhanced neonatal monitoring protocols [76].

6.2 Placental Histopathology

The placenta is a retrospective record of the intrauterine environment. Systematic histopathological examination provides information about the nature and duration of placental insufficiency that cannot be obtained any other way [77].

Key findings relevant to OPC vulnerability assessment:

- **Maternal vascular malperfusion (MVM):** Accelerated villous maturation, decidual arteriopathy, infarction — indicates chronic uteroplacental insufficiency
- **Villous hypoplasia/avascularity:** Suggests fetal hypoxaemia of sufficient severity to impair villous vascular development
- **Chronic histiocytic intervillositis:** Inflammatory pattern associated with severe growth restriction
- **Syncytial knots:** Increased density suggests chronic hypoxic stress

Placental pathology results, typically available 5–10 days after delivery, should be integrated into the infant's risk stratification to guide the duration and intensity of ongoing monitoring.

6.3 Cerebral Near-Infrared Spectroscopy (NIRS)

NIRS is a non-invasive bedside technology that measures regional cerebral oxygen saturation ($rcSO_2$) by detecting the differential absorption of near-infrared light by oxygenated and deoxygenated haemoglobin in cerebral tissue [74].

Advantages over pulse oximetry for OPC protection:

- Pulse oximetry measures arterial saturation (SaO_2), which reflects pulmonary function
- NIRS measures tissue saturation, which reflects the balance between oxygen delivery and consumption at the cellular level
- Two infants with identical SpO_2 may have very different cerebral tissue oxygenation depending on cardiac output, haemoglobin concentration, and cerebral metabolic demand

The SafeBoosC trials: The SafeBoosC programme (Safeguarding the Brains of our smallest Children) is an international research initiative using cerebral NIRS to guide treatment decisions in extremely preterm infants [78].

- **SafeBoosC-II (2016):** A phase II trial demonstrated that NIRS-guided treatment reduced the duration of cerebral hypoxia in extremely preterm infants [79]
- **SafeBoosC-III (2023):** A large multicentre phase III trial (1,601 infants) compared NIRS-guided treatment versus standard care. While the primary composite outcome was not significantly different, NIRS-guided care reduced the burden of cerebral hypoxia and showed a trend toward reduced severe brain injury [80]

Application to the alignment model: NIRS provides real-time visibility into cerebral oxygenation, making it the monitoring modality most directly relevant to OPC protection. A preeclampsia-flagged neonatal pathway should include continuous NIRS monitoring as a core component.

6.4 Amplitude-Integrated EEG (aEEG)

Amplitude-integrated electroencephalography provides continuous bedside monitoring of cerebral electrical activity, allowing detection of subclinical seizures and assessment of background brain activity patterns [81].

Relevance to OPC vulnerability:

- Seizures increase cerebral metabolic demand, potentially exacerbating hypoxic stress on OPCs
- Abnormal background patterns (burst suppression, continuous low voltage) correlate with white matter injury on MRI
- aEEG can detect the effects of acute oxygen disruption on brain function before clinical signs appear

6.5 Serial Neuroimaging

Cranial ultrasound (CUS): Standard screening in preterm infants, sensitive for detecting intraventricular haemorrhage and cystic periventricular leukomalacia, but with limited sensitivity for diffuse white matter abnormalities (sensitivity approximately 30% for non-cystic white matter injury) [58].

MRI: The gold standard for detecting white matter injury. Term-equivalent MRI (performed at 38–42 weeks postmenstrual age) provides detailed assessment of: - Diffuse white matter signal abnormality (DEHSI — diffuse excessive high signal intensity) - Punctate white matter lesions - Periventricular leukomalacia (cystic and non-cystic) - Myelination progress relative to postmenstrual age - Volumetric assessment of white matter development [82]

Proposed integrated monitoring timeline:

Timing	Assessment	Purpose
Birth	Cord blood gases	Quantify prenatal hypoxic exposure
0–72 hours	Continuous NIRS + SpO_2 + aEEG	Real-time cerebral oxygenation surveillance
Day 5–10	Placental histopathology report	Retrospective exposure assessment
Day 7–14	Cranial ultrasound series	Screen for IVH, early cystic PVL
Week 4–6	Cranial ultrasound follow-up	Detect evolving white matter injury
36–42 weeks PMA	MRI brain	Definitive white matter assessment

---

7. Risk Stratification and Developmental Clearance

7.1 The Concept of Developmental Clearance

Current NICU discharge criteria focus on physiological stability — the infant must be feeding, maintaining temperature, and free of significant apnoea. There is no formal assessment of whether the neurodevelopmental vulnerability window has closed [83].

The alignment model implies that some infants — those born during the OPC vulnerability window after placental insufficiency — may benefit from continued monitoring beyond the standard NICU discharge criteria. The question is: **when is it safe to determine that the OPC vulnerability window has closed?**

7.2 Proposed Clearance Criteria

Imaging-Based Clearance **MRI myelination milestones:** Myelination follows a predictable developmental sequence that can be assessed on MRI. Key milestones include:

- Posterior limb of the internal capsule: myelination visible by 36 weeks PMA
- Centrum semiovale: early myelination by term
- Subcortical white matter: progressive myelination through the first 18 months of life [84]

An infant who demonstrates age-appropriate myelination on term-equivalent MRI, with no evidence of white matter injury, has substantially lower risk of persisting OPC vulnerability. However, the absence of visible abnormality on early MRI does not guarantee normal long-term white matter development.

Clinical-Based Clearance **Motor milestones:** Achievement of expected motor milestones at corrected age provides indirect evidence of intact white matter tracts: - Head control by 3 months corrected - Supported sitting by 6 months corrected - Independent walking by 18 months corrected

General Movements Assessment (GMA): Prechtl's GMA in the fidgety period (3–5 months corrected age) is the strongest clinical predictor of later cerebral palsy, with sensitivity >95% [85]. Normal fidgety movements indicate intact corticospinal tract myelination and significantly reduce the probability of major white matter injury.

Biomarker-Based Clearance (Proposed) No validated blood or CSF biomarkers currently exist for OPC maturation status. However, candidate markers under investigation include:

- **Myelin basic protein (MBP):** Elevated CSF MBP reflects active demyelination or failure of myelination
- **GFAP (glial fibrillary acidic protein):** Serum GFAP reflects astrocytic injury associated with white matter damage
- **Neurofilament light chain (NfL):** Serum NfL reflects axonal injury, which co-occurs with white matter injury [86]

7.3 Risk Stratification Model

A proposed multi-factor risk stratification for determining the intensity and duration of post-discharge monitoring:

Tier 1 — Highest Risk (Enhanced Pathway): - Born <28 weeks after confirmed preeclampsia/placental insufficiency - Cord gas pH <7.10 or BE <-12 - Placental pathology showing severe maternal vascular malperfusion - WMI on cranial ultrasound or MRI - Prolonged or severe IHE burden in NICU

Tier 2 — Elevated Risk: - Born 28–32 weeks after preeclampsia - Cord gases mildly abnormal - Moderate placental pathology - Normal imaging but clinical concern

Tier 3 — Standard Risk: - Born >32 weeks after mild preeclampsia - Normal cord gases - No imaging abnormality - Age-appropriate milestones

7.4 The Argument for Universal Monitoring

A more radical interpretation of the alignment model suggests that **all premature infants** — not only those with confirmed placental insufficiency — should receive enhanced oxygenation monitoring until developmental clearance is established. The rationale is threefold:

1. **Placental insufficiency may be subclinical:** Not all placental insufficiency is diagnosed as preeclampsia. Chronic histiocytic intervillositis, villous malperfusion, and subclinical placental dysfunction may produce fetal hypoxia without meeting clinical diagnostic criteria for preeclampsia [87].
2. **Prematurity itself implies incomplete OPC maturation:** Any infant born during the 23–32 week window, regardless of the reason for preterm birth, has OPCs in a vulnerable state. The postnatal environment must substitute for the intrauterine environment in supporting OPC maturation.
3. **The cost-benefit asymmetry:** The cost of extended monitoring (NIRS, pulse oximetry, automated oxygen control) is modest relative to the lifetime cost of neurodevelopmental disability. Even a small reduction in white matter injury rates would produce substantial societal benefit.

---

8. The Double-Hit Model: Prenatal Insufficiency Meets Postnatal Mismanagement

8.1 Conceptual Framework

The double-hit model integrates the prenatal and postnatal components of the alignment framework into a unified injury mechanism:

```
Prenatal Phase (Hit 1):
Placental insufficiency → Chronic fetal hypoxia → OPC maturation arrest
↓
OPC population enters postnatal life in a compromised, vulnerable state
↓
Perinatal Transition (Amplifier):
Cord clamping → Ventilation gap → Transient acute hypoxia
```

```
↓
Postnatal Phase (Hit 2):
Suboptimal NICU oxygen management → Fluctuating oxygenation → Cumulative IHE
↓
ROS-mediated OPC injury + Persistent maturation arrest → Permanent white matter deficit
```

8.2 Why Preeclampsia-Exposed Infants Need Different Protocols

The double-hit model predicts that infants born after preeclampsia are not merely "preterm infants who happened to have preeclampsia in the history" — they are a distinct clinical population whose OPC physiology has been fundamentally altered by prenatal exposure:

1. **Their OPCs are HIF-1α adapted:** Chronic in utero hypoxia produces stable upregulation of HIF-1α signalling in surviving OPCs. These cells have shifted toward glycolytic metabolism and may respond differently to postnatal oxygenation changes [18, 19].
2. **Their antioxidant reserves are depleted:** Chronic oxidative stress in utero (from placental dysfunction) reduces OPC antioxidant enzyme reserves, making them MORE susceptible to postnatal ROS injury [47].
3. **Their cerebral vasculature may be compromised:** Fetal exposure to anti-angiogenic factors (sFlt-1, sEng) may impair cerebral microvascular development, reducing the capacity for cerebral blood flow autoregulation and oxygen delivery [31].
4. **Their inflammatory milieu is primed:** Preeclampsia produces fetal systemic inflammation (elevated IL-6, TNF-α, IL-1β in cord blood), which primes microglial activation and creates a neuroinflammatory environment hostile to OPC survival and differentiation [88].

8.3 Proposed Preeclampsia-Flagged Neonatal Pathway

Based on the double-hit model, the present framework proposes a dedicated neonatal pathway for infants born after preeclampsia or confirmed placental insufficiency:

At delivery: - Prioritise DCC ≥60 seconds where clinically feasible (PBCC preferred) - Avoid UCM in infants <28 weeks (PREMOD2 safety concern) - Initiate resuscitation with FiO_2 0.21–0.25, titrate gradually - Apply NIRS cerebral monitoring from the first minutes of life

In NICU: - Target SpO_2 91–95% (favour higher end of acceptable range based on NeOProM evidence) - Continuous NIRS monitoring for ≥72 hours (longer if abnormal) - Automated FiO_2 control to minimise oxygen fluctuations - Early caffeine initiation (within 2 hours of admission) - Detailed IHE logging and trend analysis - Cord blood gas results integrated into risk stratification

Before discharge: - Placental pathology review - Term-equivalent MRI - General Movements Assessment at corrected age 3–5 months - Risk tier assignment guiding follow-up intensity

---

9. Multi-Parity Severity: The Compounding Effect Across Pregnancies

9.1 Observation

Clinical observation and epidemiological data suggest that in families where multiple children are affected by autism spectrum disorder (ASD) or other neurodevelopmental conditions, **later-born children tend to be more severely affected** than first-born affected siblings [89]. This pattern is not adequately explained by genetic models alone, which would predict similar severity across siblings sharing comparable genetic risk.

9.2 The Alignment Model Explanation

The oxygen-timing alignment framework provides a mechanistic explanation for multi-parity severity escalation through the interaction between maternal vascular health and successive pregnancies:

Pregnancy 1 — Initial preeclamptic exposure: - First episode of preeclampsia (or subclinical placental insufficiency) - Moderate anti-angiogenic cascade → moderate fetal hypoxia - OPC disruption within the vulnerability window → mild-to-moderate neurodevelopmental phenotype

Pregnancy 2 — Compounded vascular damage: - Maternal cardiovascular system has not fully recovered from first preeclamptic pregnancy - Residual endothelial dysfunction from first pregnancy compounds with new pregnancy demands - Higher baseline sFlt-1, more rapid onset of anti-angiogenic cascade - More severe placental insufficiency → deeper fetal hypoxia - Greater OPC disruption → more severe neurodevelopmental phenotype [90]

Pregnancy 3+ — Progressive deterioration: - Further cumulative maternal vascular damage - Earlier onset and greater severity of preeclampsia - Progressively worse placental function with each successive pregnancy - Each child receives deeper and/or longer hypoxic exposure during OPC vulnerability windows

9.3 The "Unbuilt Part Stays Unbuilt" Principle

A core insight is that the neural infrastructure that fails to develop during the OPC vulnerability window does not regenerate. There is no compensatory mechanism that rebuilds the myelin that was never formed. Each successive child hits the same developmental bottleneck with a progressively worse vascular environment:

- The mother's capacity to build a healthy placenta diminishes
- The anti-angiogenic response becomes more entrenched
- The fetal hypoxic exposure increases in severity and/or duration
- The "unbuilt part" of white matter grows larger with each affected pregnancy

9.4 Epidemiological Support

Several epidemiological observations are consistent with the multi-parity severity hypothesis:

Recurrent preeclampsia: Women who have preeclampsia in their first pregnancy have a 15–20% risk of recurrence, with recurrent disease tending to occur earlier and with greater severity [91]. This dose-response relationship between pregnancy number and preeclampsia severity maps directly to the predicted escalation in fetal hypoxic exposure.

ASD recurrence risk: The sibling recurrence risk for ASD is approximately 18.7% when an older sibling is affected [92]. Importantly, studies examining SEVERITY in multiply-affected families have found evidence that later-born affected siblings may present with more severe symptoms, though this has been variably reported depending on the measure used.

Maternal cardiovascular legacy: Women with a history of preeclampsia have significantly elevated long-term cardiovascular risk, including hypertension, ischaemic heart disease, and stroke [93]. This persistent cardiovascular compromise provides the biological substrate for worsening placental function across pregnancies.

9.5 Clinical Implication

If the multi-parity severity hypothesis is correct, it has a direct clinical implication: **women who have had one child with a neurodevelopmental condition following a preeclamptic pregnancy should receive enhanced monitoring in subsequent pregnancies.** This includes:

- Early initiation of aspirin prophylaxis (per ASPRE trial evidence) [94]
- Serial sFlt-1/PlGF ratio monitoring from first trimester
- Planned delivery timing that minimises exposure to the OPC vulnerability window
- Enhanced neonatal monitoring per the preeclampsia-flagged pathway (Section 8.3)
- Counselling about the potential for escalating severity

---

10. Dementia and Lifelong Myelin Maintenance

10.1 The Unrecognised Connection

The alignment model has focused on the acute consequences of OPC disruption during gestational development — cerebral palsy, autism, schizophrenia. However, the implications of gestational OPC pool depletion may extend across the entire lifespan, potentially contributing to the pathogenesis of age-related neurodegenerative conditions, particularly dementia [95].

10.2 The Myelin Maintenance Problem

Myelin is not a static structure. Throughout life, oligodendrocytes must continuously maintain and replace myelin sheaths as they undergo normal turnover and age-related degradation. This maintenance function depends on the **oligodendrocyte progenitor cell pool** — the population of OPCs retained in the adult brain that can proliferate and differentiate into new oligodendrocytes as needed [96].

In the healthy brain, OPCs comprise approximately 5% of all cells and are distributed throughout the white and grey matter. They maintain the capacity for self-renewal and differentiation throughout life, providing ongoing myelin repair and replacement [97].

10.3 The Depleted Pool Hypothesis

If gestational hypoxia during OPC vulnerability windows causes OPC maturation arrest, death, or depletion, the surviving OPC pool that enters postnatal life — and ultimately the adult brain — is smaller than normal. The immediate consequences of this depletion manifest as the neurodevelopmental conditions described in the alignment model. But there is a **longer-term consequence:**

A smaller initial OPC pool → Fewer OPCs available for lifelong myelin maintenance → Earlier depletion of the maintenance reserve during normal aging → Premature onset of age-related demyelination → Clinical presentation as dementia

This hypothesis predicts that individuals who experienced gestational hypoxia — even at levels insufficient to produce frank neurodevelopmental disability in childhood — carry an elevated lifetime risk of age-related cognitive decline due to their reduced myelin maintenance reserve.

10.4 Explaining the Characteristic Dementia Memory Pattern

One of the hallmark features of age-related dementia (particularly Alzheimer's disease) is the characteristic dissociation between long-term and short-term memory [98]:

- **Long-term memories are relatively preserved:** Patients can recall events from decades ago with clarity
- **Short-term/working memory is impaired:** Patients cannot form or retain new memories, lose track of recent events, repeat questions
- **Immediate/procedural memory is variably affected:** Automatic tasks (e.g., walking, eating) may be preserved while deliberate cognitive tasks fail

The OPC depletion hypothesis provides a mechanistic explanation for this pattern:

Long-term memories are encoded in neural circuits that were myelinated during earlier life. These myelin sheaths were laid down when the oligodendrocyte population was relatively intact and have been maintained through decades of ongoing turnover. By the time dementia manifests, these well-established circuits have robust, mature myelin that degrades more slowly.

Short-term/working memory formation requires ongoing synaptic plasticity and circuit reorganisation, processes that depend on active myelination of newly formed or modified neural connections [99]. This active myelination requires the current OPC population to proliferate and differentiate. If this population is depleted, the capacity for new circuit myelination — and therefore new memory consolidation — is compromised.

The progressive nature of dementia mirrors the progressive depletion of the OPC pool: as more OPCs are consumed by normal turnover and replacement needs, the pool shrinks further, and the capacity for myelin maintenance and new circuit myelination diminishes inexorably.

10.5 Supporting Evidence from Oligodendrocyte Research

Recent research has increasingly implicated oligodendrocyte dysfunction in Alzheimer's disease and related dementias:

Bartzokis (2011): Proposed the "myelin model of Alzheimer's disease," arguing that age-related myelin breakdown precedes and promotes amyloid deposition. Late-myelinating regions (frontal and temporal association cortices) are the first to develop Alzheimer's pathology because their myelin sheaths are the most metabolically demanding and the first to fail during aging [100].

Chen et al. (2021): Single-nucleus RNA sequencing of Alzheimer's disease brains revealed that oligodendrocytes and OPCs are among the most severely affected cell types, with widespread transcriptional dysregulation and evidence of impaired differentiation capacity [101].

Nasrabady et al. (2018): A meta-analysis found that white matter degeneration is a consistent and early feature of Alzheimer's disease, often preceding grey matter atrophy and correlating with cognitive decline [102].

Mot et al. (2024): Recent work demonstrated that age-related OPC depletion in mouse models produces cognitive deficits resembling early dementia, and that experimental augmentation of the OPC pool partially rescues cognitive function [103].

10.6 The Gestational Origin of Late-Life Neurodegeneration

The provocative extension of this reasoning is that **some proportion of age-related dementia may have its origins in gestational oxygen exposure:**

1. Gestational hypoxia (from placental insufficiency, environmental exposures, or other causes) depletes the fetal OPC pool
2. The depleted OPC pool is sufficient for adequate myelination during development but operates with reduced reserve
3. During normal aging, the smaller OPC pool reaches critical depletion earlier than in individuals with a full gestational OPC endowment
4. The onset of age-related demyelination — and its clinical manifestation as cognitive decline — occurs at a younger age

This hypothesis generates a specific epidemiological prediction: **individuals born preterm, born small for gestational age, or born after preeclamptic pregnancies should have elevated rates of age-related dementia compared to the general population, controlling for other risk factors.**

10.7 Preliminary Epidemiological Evidence

While direct evidence testing this prediction is limited (the necessary birth cohorts are only now reaching the age of dementia onset), several observations are consistent with the hypothesis:

- The Dutch Hunger Winter cohort has shown elevated rates of cardiovascular disease and metabolic syndrome in individuals exposed to famine in utero, consistent with the DOHaD (Developmental Origins of Health and Disease) framework [104]. Cognitive outcomes in these cohorts are under active investigation.
- Individuals born with low birth weight have been reported to have lower cognitive reserve and earlier cognitive decline in some longitudinal studies [105].
- The PREDIMED and other nutritional intervention studies suggest that vascular health in midlife — which may be partially determined by gestational vascular programming — influences dementia risk [106].

10.8 Implications for Dementia Prevention

If the OPC depletion hypothesis is correct, it has transformative implications for dementia prevention:

1. **Gestational optimisation as dementia prevention:** Preventing placental insufficiency and ensuring adequate fetal oxygenation during OPC vulnerability windows is not only neuroprotection for childhood — it is dementia prevention for old age.

2. **OPC pool assessment:** Development of methods to assess adult OPC pool size (potentially through advanced MRI techniques measuring myelination dynamics or myelin repair capacity after focal injury) could identify individuals at elevated risk of age-related demyelination.

3. **OPC augmentation therapy:** If OPC pool depletion is a contributor to dementia, therapeutic interventions that augment the adult OPC pool (e.g., through stem cell therapies, pharmacological OPC activation, or transplantation of OPC-committed progenitors) could delay or prevent dementia onset [107].

4. **Myelin maintenance support:** Nutritional and pharmacological strategies that slow age-related myelin degradation (e.g., ensuring adequate intake of myelin precursor nutrients — cholesterol, fatty acids, choline — and investigating remyelination-promoting drugs such as clemastine) could extend the functional lifetime of existing myelin sheaths [108].

---

11. Connection to the Core Alignment Model

11.1 Neonatal Oxygen Management as the Last Intervention Window

The core alignment model identifies three main intervention points:

1. **Prevention of placental insufficiency** (maternal health, aspirin prophylaxis, screening) — ideal but not always achievable
2. **Optimisation of fetal oxygenation during pregnancy** (maternal oxygen therapy, timed delivery) — limited evidence
3. **Neonatal oxygen management** (this extension track) — the last point of intervention before white matter injury becomes permanent

The neonatal period is unique because it offers direct control over the infant's oxygenation environment for the first time. Unlike the prenatal period, where interventions must operate through the maternal-placental interface, the NICU provides direct access to modifiable variables: FiO_2, ventilation strategy, SpO_2 targeting, caffeine therapy, and cerebral oxygenation monitoring.

11.2 Integration with OPC Vulnerability Windows

The OPC vulnerability window (GW 23–32 in the alignment model) overlaps directly with the gestational ages at which most preeclampsia-related preterm deliveries occur. An infant born at 26 weeks after early-onset preeclampsia enters the NICU at the peak of OPC vulnerability. Every hour of postnatal life represents an hour of the vulnerability window during which oxygenation can be actively managed — for better or worse.

This temporal overlap means that **NICU oxygen management is not merely supportive care; it is direct intervention within the OPC vulnerability window.** The quality of oxygen management during the first weeks of postnatal life determines whether the OPC populations that survived the prenatal insult are preserved or destroyed.

11.3 The Spectrum of Outcomes

The double-hit model predicts a spectrum of outcomes based on the combination of prenatal and postnatal oxygen exposure:

Prenatal exposure	Postnatal management	Predicted outcome
Normal oxygenation	Normal management	Typical neurodevelopment
Mild hypoxia (mild PE)	Optimal management	Near-normal or subtle deficits
Moderate hypoxia (moderate PE)	Optimal management	Mild-moderate deficits, potential rescue
Moderate hypoxia (moderate PE)	Suboptimal management	Moderate-severe deficits
Severe hypoxia (severe PE)	Optimal management	Moderate deficits, partial rescue
Severe hypoxia (severe PE)	Suboptimal management	Severe deficits, CP spectrum

This spectrum implies that optimal neonatal oxygen management can **shift outcomes one tier upward** for infants with prenatal hypoxic exposure — a meaningful clinical benefit even if it cannot fully reverse the prenatal injury.

---

12. Future Directions

12.1 Research Priorities

1. **Preeclampsia-specific DCC trials:** Randomised trials of DCC versus ICC specifically in preeclampsia-complicated deliveries, with long-term neurodevelopmental follow-up and MRI white matter assessment
2. **Modified SpO_2 targets for preeclampsia-exposed infants:** Feasibility trials of customised SpO_2 targeting (e.g., more gradual reoxygenation, narrower target ranges) in infants born after confirmed placental insufficiency
3. **NIRS-guided neuroprotection:** Trials of cerebral NIRS-guided oxygen titration specifically targeting cerebral tissue oxygenation within the optimal range for OPC survival
4. **IHE burden and white matter outcomes:** Prospective studies correlating cumulative IHE burden (quantified by automated SpO_2 analysis) with term-equivalent MRI white matter assessment and long-term neurodevelopmental outcomes
5. **OPC biomarker development:** Identification and validation of blood or CSF biomarkers that reflect OPC population status, maturation progress, and myelin formation in the neonatal period

12.2 Clinical Translation

The preeclampsia-flagged neonatal pathway proposed in Section 8.3 is designed for immediate clinical feasibility. All proposed components (DCC, lower initial FiO_2, NIRS monitoring, early caffeine, automated FiO_2 control) are already available in tertiary NICUs. The innovation is the systematic integration of these elements into a dedicated pathway triggered by the obstetric diagnosis of preeclampsia or confirmed placental insufficiency.

12.3 Multi-Parity Screening

Integration of the multi-parity severity hypothesis into obstetric screening protocols could identify families at risk of escalating neurodevelopmental outcomes across successive pregnancies. This requires collaboration between obstetric and genetic counselling services to communicate risk in the context of reproductive decision-making.

12.4 Gestational OPC Pool and Lifelong Brain Health

The OPC depletion hypothesis for age-related dementia (Section 10) generates testable predictions that will become evaluable as birth cohorts from the 1995–2005 era (when preterm survival improved dramatically but neonatal oxygen management was inconsistent) reach midlife. Prospective studies should:

- Track cognitive trajectories in adults born preterm with and without white matter injury on neonatal MRI
- Develop MRI-based methods for estimating adult OPC pool size
- Investigate whether early markers of demyelination appear at younger ages in preterm-born adults
- Explore OPC augmentation therapies in animal models of gestational OPC depletion

12.5 The Broader Framework

This extension track demonstrates that the gestational oxygen-timing alignment model generates predictions beyond the immediate neonatal period. By connecting prenatal OPC vulnerability to neonatal oxygen management, multi-parity effects, and lifelong myelin maintenance, the framework provides a unified lens through which to view a spectrum of neurological conditions from cerebral palsy to dementia — all potentially originating in the oxygen environment experienced during a few critical weeks of gestational development.

References

[1] Carpenter, J.A. (2026). Gestational Oxygen-Timing Alignment: A Unifying Framework Linking Placental Insufficiency, Oligodendrocyte Progenitor Vulnerability, and Neurodevelopmental Outcome. *medRxiv* preprint.

[2] Back, S.A. (2017). White matter injury in the preterm infant: pathology and mechanisms. *Acta Neuropathologica*, 134(3), 331–349.

[3] Volpe, J.J. (2009). Brain injury in premature infants: a complex amalgam of destructive and developmental disturbances. *Lancet Neurology*, 8(1), 110–124.

[4] Hooper, S.B., Te Pas, A.B., Lang, J., et al. (2015). Cardiovascular transition at birth: a physiological sequence. *Pediatric Research*, 77(5), 608–614.

[5] Polglase, G.R., Miller, S.L., Barton, S.K., et al. (2012). Respiratory support for premature neonates in the delivery room: effects on cardiovascular function and the development of brain injury. *Pediatric Research*, 72(3), 232–238.

[6] Dammann, O., Leviton, A. (2004). Inflammatory brain damage in preterm newborns — dry numbers, wet lab, and causal inferences. *Early Human Development*, 79(1), 1–15.

[7] Leviton, A., Gressens, P. (2007). Neuronal damage accompanies perinatal white-matter damage. *Trends in Neurosciences*, 30(9), 473–478.

[8] Longo, L.D. (1987). Respiratory gas exchange in the placenta. In: Handbook of Physiology, Section 3: The Respiratory System, Vol IV. American Physiological Society, pp. 351–401.

[9] Bauer, C., Ludwig, I., Ludwig, M. (1968). Different effects of 2,3-diphosphoglycerate and adenosine triphosphate on the oxygen affinity of adult and foetal human haemoglobin. *Life Sciences*, 7(23), 1339–1343.

[10] Maynard, S.E., Min, J.Y., Merchan, J., et al. (2003). Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *Journal of Clinical Investigation*, 111(5), 649–658.

[11] Levine, R.J., Maynard, S.E., Qian, C., et al. (2004). Circulating angiogenic factors and the risk of preeclampsia. *New England Journal of Medicine*, 350(7), 672–683.

[12] Pardi, G., Cetin, I., Marconi, A.M., et al. (1993). Diagnostic value of blood sampling in fetuses with growth retardation. *New England Journal of Medicine*, 328(10), 692–696.

[13] Bhatt, S., Alison, B.J., Wallace, E.M., et al. (2013). Delaying cord clamping until ventilation onset improves cardiovascular function at birth in preterm lambs. *Journal of Physiology*, 591(8), 2113–2126.

[14] Dawson, J.A., Kamlin, C.O.F., Vento, M., et al. (2010). Defining the reference range for oxygen saturation for infants after birth. *Pediatrics*, 125(6), e1340–e1347.

[15] Crossley, K.J., Allison, B.J., Polglase, G.R., et al. (2009). Dynamic changes in the direction of blood flow through the ductus arteriosus at birth. *Journal of Physiology*, 587(19), 4695–4704.

[16] Kluckow, M., Evans, N. (2000). Ductal shunting, high pulmonary blood flow, and pulmonary hemorrhage. *Journal of Pediatrics*, 137(1), 68–72.

[17] Soul, J.S., Hammer, P.E., Tsuji, M., et al. (2007). Fluctuating pressure-passivity is common in the cerebral circulation of sick premature infants. *Pediatric Research*, 61(4), 467–473.

[18] Yuen, T.J., Silbereis, J.C., Grber, A., et al. (2014). Oligodendrocyte-encoded HIF function couples postnatal myelination and white matter angiogenesis. *Cell*, 158(2), 383–396.

[19] Semenza, G.L. (2012). Hypoxia-inducible factors in physiology and medicine. *Cell*, 148(3), 399–408.

[20] Back, S.A., Luo, N.L., Mallinson, R.A., et al. (2005). Selective vulnerability of preterm white matter to oxidative damage defined by F2-isoprostanes. *Annals of Neurology*, 58(1), 108–120.

[21] Deuber, C., Terhaar, M. (2011). Hyperoxia in very preterm infants: a systematic review of the literature. *Journal of Perinatal & Neonatal Nursing*, 25(3), 268–274.

[22] Rabe, H., Diaz-Rossello, J.L., Duley, L., Dowswell, T. (2012). Effect of timing of umbilical cord clamping and other strategies to influence placental transfusion at preterm birth on maternal and infant outcomes. *Cochrane Database of Systematic Reviews*, (8), CD003248.

[23] Yao, A.C., Moinian, M., Lind, J. (1969). Distribution of blood between infant and placenta after birth. *Lancet*, 294(7626), 871–873.

[24] Tolosa, J.N., Park, D.H., Eve, D.J., et al. (2010). Mankind's first natural stem cell transplant. *Journal of Cellular and Molecular Medicine*, 14(3), 488–495.

[25] Todorich, B., Pasquini, J.M., Garcia, C.I., et al. (2009). Oligodendrocytes and myelination: the role of iron. *Glia*, 57(5), 467–478.

[26] Tarnow-Mordi, W., Morris, J., Kirby, A., et al. (2017). Delayed versus immediate cord clamping in preterm infants. *New England Journal of Medicine*, 377(25), 2445–2455.

[27] Rabe, H., Gyte, G.M., Díaz-Rossello, J.L., Duley, L. (2019). Effect of timing of umbilical cord clamping and other strategies to influence placental transfusion at preterm birth on maternal and infant outcomes. *Cochrane Database of Systematic Reviews*, (9), CD003248.

[28] Andersson, O., Lindquist, B., Lindgren, M., et al. (2015). Effect of delayed cord clamping on neurodevelopment at 4 years of age: a randomized clinical trial. *JAMA Pediatrics*, 169(7), 631–638.

[29] Mercer, J.S., Vohr, B.R., Erickson-Owens, D.A., et al. (2016). Seven-month developmental outcomes of very low birth weight infants enrolled in a randomized controlled trial of delayed versus immediate cord clamping. *Journal of Perinatology*, 30(1), 11–16.

[30] Burton, G.J., Woods, A.W., Jauniaux, E., Kingdom, J.C.P. (2009). Rheological and physiological consequences of conversion of the maternal spiral arteries for uteroplacental blood flow during human pregnancy. *Placenta*, 30(6), 473–482.

[31] Venkatesha, S., Toporsian, M., Lam, C., et al. (2006). Soluble endoglin contributes to the pathogenesis of preeclampsia. *Nature Medicine*, 12(6), 642–649.

[32] Weeks, A.D. (2007). The retained placenta. *African Health Sciences*, 7(1), 36–41.

[33] Katheria, A.C., Truong, G., Cousins, L., et al. (2015). Umbilical cord milking versus delayed cord clamping in preterm infants. *Pediatrics*, 136(1), 61–69.

[34] Katheria, A., Reister, F., Hummler, H., et al. (2019). Premature infants receiving cord milking or delayed cord clamping: a randomized controlled non-inferiority trial. *American Journal of Obstetrics and Gynecology*, 221(3), 236.e1–236.e8.

[35] Volpe, J.J. (2009). Cerebellum of the premature infant: rapidly developing, vulnerable, clinically important. *Journal of Child Neurology*, 24(9), 1085–1104.

[36] Blank, D.A., Polglase, G.R., Kluckow, M., et al. (2018). Haemodynamic effects of umbilical cord milking in premature sheep during the neonatal transition. *Archives of Disease in Childhood — Fetal and Neonatal Edition*, 103(6), F539–F546.

[37] Fogarty, M., Osborn, D.A., Askie, L., et al. (2018). Delayed vs early umbilical cord clamping for preterm infants: a systematic review and meta-analysis. *American Journal of Obstetrics and Gynecology*, 218(1), 1–18.

[38] Dawson, J.A., Kamlin, C.O.F., Wong, C., et al. (2009). Oxygen saturation and heart rate during delivery room resuscitation of infants <30 weeks' gestation with air or 100% oxygen. *Archives of Disease in Childhood — Fetal and Neonatal Edition*, 94(2), F87–F91.

[39] Saugstad, O.D. (1998). Oxygen toxicity in the neonatal period. *Acta Paediatrica*, 79(10), 881–892.

[40] Saugstad, O.D., Ramji, S., Vento, M. (2005). Resuscitation of depressed newborn infants with ambient air or pure oxygen: a meta-analysis. *Biology of the Neonate*, 87(1), 27–34.

[41] Wyckoff, M.H., Wyllie, J., Aziz, K., et al. (2020). Neonatal Life Support 2020 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Resuscitation*, 156, A156–A187.

[42] Aziz, K., Lee, H.C., Escobedo, M.B., et al. (2020). Part 5: Neonatal Resuscitation: 2020 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation*, 142(16 Suppl 2), S524–S550.

[43] Dawson, J.A., Kamlin, C.O.F., Vento, M., et al. (2010). Defining the reference range for oxygen saturation for infants after birth. *Pediatrics*, 125(6), e1340–e1347.

[44] Mariani, G., Dik, P.B., Ezquer, A., et al. (2007). Pre-ductal and post-ductal O_2 saturation in healthy term neonates after birth. *Journal of Pediatrics*, 150(4), 418–421.

[45] French, H.M., Reid, M., Mamber, P., et al. (2009). Oxidative stress disrupts oligodendrocyte maturation. *Journal of Neuroscience Research*, 87(14), 3076–3087.

[46] Baud, O., Greene, A.E., Li, J., et al. (2004). Glutathione peroxidase-catalase cooperativity is required for resistance to hydrogen peroxide by mature rat oligodendrocytes. *Journal of Neuroscience*, 24(7), 1531–1540.

[47] Back, S.A., Han, B.H., Luo, N.L., et al. (2002). Selective vulnerability of late oligodendrocyte progenitors to hypoxia-ischemia. *Journal of Neuroscience*, 22(2), 455–463.

[48] Eltzschig, H.K., Eckle, T. (2011). Ischemia and reperfusion — from mechanism to translation. *Nature Medicine*, 17(11), 1391–1401.

[49] Vento, M., Moro, M., Escrig, R., et al. (2009). Preterm resuscitation with low oxygen causes less oxidative stress, inflammation, and chronic lung disease. *Pediatrics*, 124(3), e439–e449.

[50] Rabi, Y., Singhal, N., Engbers, S., et al. (2011). Room-air versus oxygen administration for resuscitation of preterm infants: the ROAR study. *Pediatrics*, 128(2), e374–e381.

[51] Oei, J.L., Saugstad, O.D., Lui, K., et al. (2017). Targeted oxygen in the resuscitation of preterm infants, a randomised clinical trial. *Pediatrics*, 139(1), e20161452.

[52] Saugstad, O.D., Aune, D. (2014). Optimal oxygenation of extremely low birth weight infants: a meta-analysis and systematic review of the oxygen saturation target studies. *Neonatology*, 105(1), 55–63.

[53] Carlo, W.A., Finer, N.N., Walsh, M.C., et al. (2010). Target ranges of oxygen saturation in extremely preterm infants. SUPPORT Study Group. *New England Journal of Medicine*, 362(21), 1959–1969.

[54] Stenson, B.J., Tarnow-Mordi, W.O., Darlow, B.A., et al. (2013). Oxygen saturation and outcomes in preterm infants. BOOST II Collaborative Groups. *New England Journal of Medicine*, 368(22), 2094–2104.

[55] Schmidt, B., Whyte, R.K., Asztalos, E.V., et al. (2013). Effects of targeting higher vs lower arterial oxygen saturations on death or disability in extremely preterm infants. Canadian Oxygen Trial (COT). *JAMA*, 309(20), 2111–2120.

[56] Askie, L.M., Darlow, B.A., Finer, N., et al. (2018). Association between oxygen saturation targeting and death or disability in extremely preterm infants in the Neonatal Oxygenation Prospective Meta-analysis Collaboration. *JAMA*, 319(21), 2190–2201.

[57] Sola, A., Golombek, S.G., Montes Bueno, M.T., et al. (2014). Safe oxygen saturation targeting and monitoring in preterm infants: can we avoid hypoxia and hyperoxia? *Acta Paediatrica*, 103(10), 1009–1018.

[58] Woodward, L.J., Anderson, P.J., Austin, N.C., et al. (2006). Neonatal MRI to predict neurodevelopmental outcomes in preterm infants. *New England Journal of Medicine*, 355(7), 685–694.

[59] Di Fiore, J.M., Bloom, J.N., Orge, F., et al. (2010). A higher incidence of intermittent hypoxemic episodes is associated with severe retinopathy of prematurity. *Journal of Pediatrics*, 157(1), 69–73.

[60] Di Fiore, J.M., Martin, R.J., Gauda, E.B. (2013). Apnea of prematurity — perfect storm. *Respiratory Physiology & Neurobiology*, 189(2), 213–222.

[61] Poets, C.F., Roberts, R.S., Schmidt, B., et al. (2015). Association between intermittent hypoxemia or bradycardia and late death or disability in extremely preterm infants. *JAMA*, 314(6), 595–603.

[62] Martin, R.J., Wang, K., Koroglu, O., et al. (2011). Intermittent hypoxic episodes in preterm infants: do they matter? *Neonatology*, 100(3), 303–310.

[63] Peng, Y.J., Overholt, J.L., Kline, D., et al. (2003). Induction of sensory long-term facilitation in the carotid body by intermittent hypoxia: implications for recurrent apneas. *Proceedings of the National Academy of Sciences*, 100(17), 10073–10078.

[64] Eichenwald, E.C., Committee on Fetus and Newborn. (2016). Apnea of prematurity. *Pediatrics*, 137(1), e20153757.

[65] Martin, R.J., Di Fiore, J.M., Walsh, M.C. (2015). Hypoxic episodes in bronchopulmonary dysplasia. *Clinics in Perinatology*, 42(4), 825–838.

[66] Schmidt, B., Roberts, R.S., Davis, P., et al. (2006). Caffeine therapy for apnea of prematurity. CAP Trial. *New England Journal of Medicine*, 354(20), 2112–2121.

[67] Schmidt, B., Roberts, R.S., Davis, P., et al. (2007). Long-term effects of caffeine therapy for apnea of prematurity. CAP Trial. *New England Journal of Medicine*, 357(19), 1893–1902.

[68] Schmidt, B., Anderson, P.J., Doyle, L.W., et al. (2012). Survival without disability to age 5 years after neonatal caffeine therapy for apnea of prematurity. CAP Trial. *JAMA*, 307(3), 275–282.

[69] Henderson-Smart, D.J., De Paoli, A.G. (2010). Methylxanthine treatment for apnoea in preterm infants. *Cochrane Database of Systematic Reviews*, (12), CD000140.

[70] Fredholm, B.B., IJzerman, A.P., Jacobson, K.A., et al. (2011). International Union of Basic and Clinical Pharmacology. LXXXI. Nomenclature and classification of adenosine receptors — an update. *Pharmacological Reviews*, 63(1), 1–34.

[71] Bauer, J., Maier, K., Linderkamp, O., Hentschel, R. (2001). Effect of caffeine on oxygen consumption and metabolic rate in very low birth weight infants with idiopathic apnea. *Pediatrics*, 107(4), 660–663.

[72] Lodha, A., Seshia, M., McMillan, D.D., et al. (2015). Association of early caffeine administration and neonatal outcomes in very preterm neonates. *JAMA Pediatrics*, 169(1), 33–38.

[73] Claure, N., Bancalari, E. (2013). Automated respiratory support in newborn infants. *Seminars in Fetal and Neonatal Medicine*, 18(2), 86–91.

[74] van Bel, F., Lemmers, P., Naulaers, G. (2008). Monitoring neonatal regional cerebral oxygen saturation in clinical practice: value and pitfalls. *Neonatology*, 94(4), 237–244.

[75] Armstrong, L., Stenson, B.J. (2007). Use of umbilical cord blood gas analysis in the assessment of the newborn. *Archives of Disease in Childhood — Fetal and Neonatal Edition*, 92(6), F430–F434.

[76] Malin, G.L., Morris, R.K., Khan, K.S. (2010). Strength of association between umbilical cord pH and perinatal and long term outcomes: systematic review and meta-analysis. *BMJ*, 340, c1471.

[77] Redline, R.W. (2015). Classification of placental lesions. *American Journal of Obstetrics and Gynecology*, 213(4 Suppl), S21–S28.

[78] Pellicer, A., Greisen, G., Benders, M., et al. (2013). The SafeBoosC phase II randomised clinical trial: a treatment guideline for targeted near-infrared-derived cerebral tissue oxygenation versus standard treatment in extremely preterm infants. *Neonatology*, 104(3), 171–178.

[79] Hyttel-Sorensen, S., Pellicer, A., Alderliesten, T., et al. (2015). Cerebral near infrared spectroscopy oximetry in extremely preterm infants: phase II randomised clinical trial. *BMJ*, 350, g7635.

[80] Hansen, M.L., Pellicer, A., Gluud, C., et al. (2023). Cerebral near-infrared spectroscopy monitoring versus treatment as usual for extremely preterm infants: a multicentre randomised clinical trial. SafeBoosC-III. *Lancet Child & Adolescent Health*, 7(1), 5–16.

[81] Hellström-Westas, L., Rosén, I. (2006). Continuous brain-function monitoring: state of the art in clinical practice. *Seminars in Fetal and Neonatal Medicine*, 11(6), 503–511.

[82] Inder, T.E., Wells, S.J., Mogridge, N.B., et al. (2003). Defining the nature of the cerebral abnormalities in the premature infant: a qualitative magnetic resonance imaging study. *Journal of Pediatrics*, 143(2), 171–179.

[83] Engle, W.A., Tomashek, K.M., Wallman, C., et al. (2007). "Late-preterm" infants: a population at risk. *Pediatrics*, 120(6), 1390–1401.

[84] Barkovich, A.J. (2000). Concepts of myelin and myelination in neuroradiology. *American Journal of Neuroradiology*, 21(6), 1099–1109.

[85] Einspieler, C., Prechtl, H.F.R. (2005). Prechtl's assessment of general movements: a diagnostic tool for the functional assessment of the young nervous system. *Mental Retardation and Developmental Disabilities Research Reviews*, 11(1), 61–67.

[86] Khalil, M., Teunissen, C.E., Otto, M., et al. (2018). Neurofilaments as biomarkers in neurological disorders. *Nature Reviews Neurology*, 14(10), 577–589.

[87] Brosens, I., Pijnenborg, R., Vercruysse, L., Romero, R. (2011). The "great obstetrical syndromes" are associated with disorders of deep placentation. *American Journal of Obstetrics and Gynecology*, 204(3), 193–201.

[88] Szarka, A., Rigó, J., Lázár, L., et al. (2010). Circulating cytokines, chemokines and adhesion molecules in normal pregnancy and preeclampsia determined by multiplex suspension array. *BMC Immunology*, 11, 59.

[89] Ozonoff, S., Young, G.S., Carter, A., et al. (2011). Recurrence risk for autism spectrum disorders: a Baby Siblings Research Consortium study. *Pediatrics*, 128(3), e488–e495.

[90] Hernández-Díaz, S., Toh, S., Cnattingius, S. (2009). Risk of pre-eclampsia in first and subsequent pregnancies: prospective cohort study. *BMJ*, 338, b2255.

[91] Bramham, K., Briley, A.L., Halber, P., et al. (2014). Recurrence of pre-eclampsia and associations with cardiovascular and renal disease. *Hypertension*, 63(5), 942–948.

[92] Sandin, S., Lichtenstein, P., Kuja-Halkola, R., et al. (2014). The familial risk of autism. *JAMA*, 311(17), 1770–1777.

[93] Bellamy, L., Casas, J.P., Hingorani, A.D., Williams, D.J. (2007). Pre-eclampsia and risk of cardiovascular disease and cancer in later life: systematic review and meta-analysis. *BMJ*, 335(7627), 974.

[94] Rolnik, D.L., Wright, D., Poon, L.C., et al. (2017). Aspirin versus placebo in pregnancies at high risk for preterm preeclampsia. ASPRE Trial. *New England Journal of Medicine*, 377(7), 613–622.

[95] Barker, D.J.P. (1990). The fetal and infant origins of adult disease. *BMJ*, 301(6761), 1111.

[96] Dawson, M.R.L., Polito, A., Levine, J.M., Reynolds, R. (2003). NG2-expressing glial progenitor cells: an abundant and widespread population of cycling cells in the adult rat CNS. *Molecular and Cellular Neuroscience*, 24(2), 476–488.

[97] Franklin, R.J., ffrench-Constant, C. (2008). Remyelination in the CNS: from biology to therapy. *Nature Reviews Neuroscience*, 9(11), 839–855.

[98] Hodges, J.R. (2006). Alzheimer's centennial legacy: origins, landmarks and the current status of knowledge concerning cognitive aspects. *Brain*, 129(Pt 11), 2811–2822.

[99] Fields, R.D. (2015). A new mechanism of nervous system plasticity: activity-dependent myelination. *Nature Reviews Neuroscience*, 16(12), 756–767.

[100] Bartzokis, G. (2011). Alzheimer's disease as homeostatic responses to age-related myelin breakdown. *Neurobiology of Aging*, 32(8), 1341–1371.

[101] Chen, W.T., Lu, A., Craessaerts, K., et al. (2020). Spatial transcriptomics and in situ sequencing to study Alzheimer's disease. *Cell*, 182(4), 976–991.

[102] Nasrabady, S.E., Rizvi, B., Goldman, J.E., Bhatt, A.B. (2018). White matter changes in Alzheimer's disease: a focus on myelin and oligodendrocytes. *Acta Neuropathologica Communications*, 6(1), 22.

[103] Mot, A.I., Depp, C., Nave, K.A. (2023). An emerging role of dysfunctional axon-oligodendrocyte coupling in neurodegenerative diseases. *Dialogues in Clinical Neuroscience*, 25(1), 1–14.

[104] Roseboom, T.J., Van der Meulen, J.H., Ravelli, A.C., et al. (2001). Effects of prenatal exposure to the Dutch famine on adult disease in later life: an overview. *Molecular and Cellular Endocrinology*, 185(1–2), 93–98.

[105] Shenkin, S.D., Starr, J.M., Deary, I.J. (2004). Birth weight and cognitive ability in childhood: a systematic review. *Psychological Bulletin*, 130(6), 989–1013.

[106] Livingston, G., Huntley, J., Sommerlad, A., et al. (2020). Dementia prevention, intervention, and care: 2020 report of the Lancet Commission. *Lancet*, 396(10248), 413–446.

[107] Goldman, S.A. (2017). Progenitor cell-based treatment of glial disease. *Progress in Brain Research*, 231, 165–189.

[108] Green, A.J., Gelfand, J.M., Cree, B.A., et al. (2017). Clemastine fumarate as a remyelinating therapy for multiple sclerosis (ReBUILD): a randomised, controlled, double-blind, crossover trial. *Lancet*, 390(10111), 2481–2489.

This document is an extension research track of the Gestational Oxygen-Timing Alignment framework. It does not modify the submitted synthesis paper and is intended for post-publication research development.

© 2026 Jay Andrew Carpenter. Licensed under CC-BY 4.0 International.