

# Exercises Partial differential equation

Simone Coli 6771371

Sheet 7

## 1 Exercise

Let us consider  $\Omega \subset \mathbb{R}^n$  an open set and let us define  $\Omega_\varepsilon \doteq \{x \in \Omega : \text{dist}(x, \partial\Omega) > \varepsilon\}$ . Let us now consider  $u \in L^1_{loc}(\Omega)$  and its mollification  $u_\varepsilon : \Omega_\varepsilon \rightarrow \mathbb{R}$

$$u_\varepsilon(x) = (\rho_\varepsilon * u)(x)$$

we want to show that

a) If  $u \in C^0(\Omega)$  than  $u_\varepsilon \rightarrow u$  uniformly in compact subsets as  $\varepsilon \rightarrow 0$ . By definition of continuous we know that:

$$\forall \varepsilon > 0 \exists \delta : \forall N_\varepsilon \in \mathbb{N}, x, y \in \Omega |x - y| < \delta, |u(x) - u(y)| < \varepsilon$$

If we consider

$$|u_\varepsilon(x) - u(x)| = \left| \int_{\Omega'} \rho_\varepsilon(x - y)u(y)dy - u(x) \right|$$

Using the commutability of the convolution operator we have that

$$\left| \int_{\Omega'} \rho_\varepsilon(x - y)u(y)dy - u(x) \right| = \left| \int_{\Omega'} \rho_\varepsilon(y)u(x - y)dy - u(x) \int_{\Omega'} \rho_\varepsilon(y)dy \right|$$

where  $\Omega'$  is the set where  $\rho_\varepsilon$  is defined.

$$\left| \int_{\Omega'} \rho_\varepsilon(y)(u(x - y) - u(x))dy \right| \leq \int_{\Omega'} |\rho_\varepsilon(y)| |u(x - y) - u(x)| dy < \int_{\Omega'} |\rho_\varepsilon(y)| \varepsilon dy$$

Let us consider a ball  $\overline{B_\varepsilon(0)} \supset \Omega'$

$$\int_{\Omega'} |\rho_\varepsilon(y)| \varepsilon dy \leq \int_{\overline{B_\varepsilon(0)}} |\rho_\varepsilon(y)| \varepsilon dy$$

for  $\varepsilon \rightarrow 0$  this integral goes to zero, meaning that  $\sup_{x \in \Omega} |u_\varepsilon(x) - u(x)| \rightarrow 0$  and therefore  $u_\varepsilon$  converges uniformly to  $u$ .

b) We now want to show that if  $u \in L^p(\Omega)$  than  $u_\varepsilon \rightarrow u$  in  $L^p_{loc}$  as  $\varepsilon \rightarrow 0$ . Let us consider

$$|u_\varepsilon(x) - u(x)|^p = \left| \int_{\Omega'} \rho_\varepsilon(x - y)u(y)dy - u(x) \right|^p = \left| \int_{\overline{B_\varepsilon(0)}} \rho_\varepsilon(y)(u(x - y) - u(x))dy \right|^p$$

by the same argument as before. Now this is bounded from above by

$$\left| \int_{\overline{B_\varepsilon(0)}} \rho_\varepsilon(y) 2 \sup_{\overline{B_\varepsilon(0)}} u(x) dy \right|^p$$

If we take a subset  $\Omega' \subset\subset \Omega$  we want to show the definition of  $L^p$  space

$$\int_{\Omega'} |u_\varepsilon(x) - u(x)|^p dx \leq \int_{\Omega'} 2 \sup_{\overline{B_\varepsilon(0)}} u(x) dx = 2 \sup_{\overline{B_\varepsilon(0)}} u(x) \mu(\Omega')$$

which goes to 0 as  $\varepsilon \rightarrow 0$  since  $\Omega' \subset\subset \Omega$ .

c) We want to show that if  $u \in W^{k,p}(\Omega)$  and  $k > 0$  then  $u_\varepsilon \rightarrow u$  for  $\varepsilon \rightarrow 0$  in  $W_{loc}^{k,p}$ . By definition of  $W^{k,p}(\Omega)$  we know that  $D^\alpha u \in L^p$  for  $|\alpha| < k$ , then, using the previous result and the fact that  $(D^\alpha u)_\varepsilon = D^\alpha u_\varepsilon$  we get the claim.

d) We want to show that if  $u \in C^{0,1}$  with Lipschitz constant  $L$  then  $u_\varepsilon \in C^{0,1}$  with the same constant. By definition of Lipschitz we want to prove that

$$|u_\varepsilon(x) - u_\varepsilon(z)| < L|x - z|$$

because of that let us consider

$$|u_\varepsilon(x) - u_\varepsilon(z)| = \left| \int_{\Omega} \rho_\varepsilon(x-y) u(y) dy - \int_{\Omega} \rho_\varepsilon(z-y) u(y) dy \right| =$$

Using the commutation property for convolutions we have that:

$$= \left| \int_{\Omega} \rho_\varepsilon(y) (u(x-y) - u(z-y)) dy \right| < \int_{\Omega} |\rho_\varepsilon(y)| L|x-z| dy$$

Since  $u \in C^{0,1}(\Omega)$ . Proving the statement.

## 2 Exercise

We want to find the weak derivative of

$$f(x) = \begin{cases} 2x-1 & x \leq 0 \\ 1-3x & x \geq 0 \end{cases}$$

By its definition we have that:

$$\begin{aligned} - \int v \varphi dx &= \int u D \varphi dx \\ &= \int_{-\infty}^0 (2x-1) \varphi' dx + \int_0^\infty (1-3x) \varphi' dx = \\ &= \int_{-\infty}^0 2 \varphi dx + \int_0^\infty 3 \varphi dx \end{aligned}$$

using integration by parts. We then have that the directional derivative of  $f$  is

$$v(x) = \begin{cases} 2 & x < 0 \\ -3 & x > 0 \end{cases}$$

Since we computed it using the definition we have that it is in fact the weak derivative of  $f$ .

### 3 Exercise

Let  $\Omega \subset \mathbb{R}^n$  open and bounded such that  $0 \in \Omega$ . We want to show that a function  $u(x) = |x|^{-\alpha}$  is in  $W^{k,0}(\Omega)$  as long as  $k + \alpha < n$ .

This means that we want to show that

$$\int_{\Omega} v \varphi dx = -(-1)^k \int_{\Omega} D^{\beta} \varphi |x|^{-\alpha} dx$$

converges, where we used a multi index  $\beta$  such that  $|\beta| = k$ . If we move in spherical coordinates, we will have that:

$$\int_{\Omega} D_1^{k_1} \dots D_n^{k_n} r^{-\alpha} r^{n-1} dr d\theta$$

where  $d\theta$  contains all the angular part of the integral. Using integration by parts then we have that

$$(-1)^k \int_{\Omega} \varphi D^k r^{n-1-\alpha} dr d\theta \leq \sup_{\Omega} |\varphi| \int_{\Omega} D^k r^{n-1-\alpha} dr d\theta$$

by the chain rule we know that the directional derivative in the direction of  $x_i$  or  $r^{n-1-\alpha}$  will be less or equal than its directional derivative in the direction of  $r$ , because the function does not depend on the angular components. Then, let  $\sup_{\Omega} |\varphi| = c$ :

$$c \int_{\Omega} r^{n-1-\alpha-k} dr d\theta = [r^{n-\alpha-k}]_0^R$$

This will be non singular if, and only if, the exponent is not less than zero, that is  $n - \alpha - k > 0$  which proves the statement.

### 4 Exercise

Let  $\Omega \subset \mathbb{R}^n$  and an open and bounded subset  $\Omega' \subset \subset \Omega$  and  $d \doteq \text{dist}(\Omega', \partial\Omega)$ , we want to show that there exists a function  $\eta \in C_c^{\infty}(\Omega)$  and a constant  $C = C(n)$  such that

$$0 \leq \eta \leq 1, \quad \eta|_{\Omega'} \equiv 1, \quad |D\eta| \leq \frac{C}{d}$$

What we want to do is construct a function which grows from zero to 1 before reaching  $\Omega'$ , and we want to bound this growth from above. Let us

therefore define two subsets  $A, B \subset \Omega$  such that  $\Omega' \subset B \subset A \subset \Omega$ . In particular, we want  $\text{dist}(A, \partial\Omega) = \text{dist}(\Omega', \partial B) = \frac{d}{4}$ , meaning that  $\text{dist}(B, \partial A) = \frac{d}{2}$ . Then consider the function

$$f(x) = \begin{cases} 1 & x \in B \\ -\frac{2}{d}x + k & x \in A \setminus B \\ 0 & x \in \Omega \setminus A \end{cases}$$

Where  $k$  is just a constant which must be chosen to be such that the linear function has value 1 at  $\partial B$  and 0 at  $\partial A$ . However, this function is not smooth. To make it  $C_c^\infty$  we apply the mollification to  $f$  defining  $0 < h < \text{dist}(B, \partial A)$

$$\eta(x) = \int_{\Omega} \rho_h(x - y) f(y) dy$$