Spatial sensitivity analysis of social simulation models

J. Raimbault^{1,2,3,4,*} and J. Perret¹
* juste.raimbault@ign.fr

¹LASTIG, IGN-ENSG ²CASA, UCL ³UPS CNRS 3611 ISC-PIF ⁴UMR CNRS 8504 Géographie-cités

CCS 2023
Online session - Auditorium 2
October 17th 2023

Validation of simulation models

Typology of validation methods for simulation models from a systematic review [Raimbault, 2023]

- prediction
- sensitivity analysis
- uncertainty
- multiple methods
- benchmark
- calibration
- optimisation

- visualisation
- Pattern Oriented Modelling
- participatory
- exploration
- mixed
- surrogate

Methods, standards and definitions strongly depend on disciplines and model functions [Raimbault, 2019b]

Specificities of socio-spatial systems

- ightarrow Spatio-temporal non-stationarity and non-ergodicity [Raimbault, 2019a]
- \rightarrow Fuzzy and noisy data [Olteanu-Raimond et al., 2015] \rightarrow Genericity/specific of patterns and processes [Raimbault et al., 2020] \rightarrow Modifiable Areal Unit Problem [Wong, 2004] \rightarrow Multiscalar systems [Raimbault, 2021b]

Spatial sensitivity analysis

Spatial configurations are model parameters too

- "Space matters": impact of spatial configuration on model behavior
- Model behaviours which are robust to spatial configuration
- Model behaviours which are robust to noise in real datasets

⇒ Construction of a generic library for spatial sensitivity analysis, including the generation of synthetic data, perturbation of real data and indicators

Synthetic data: general context

- \rightarrow coupling models with spatial configuration generators (spatial synthetic data) gives model sensitivity to space through sensitivity analysis of the coupled model
- \rightarrow synthetic urban forms resembling real configurations
- \rightarrow at different scales: microscopic (buildings), mesoscopic (population distribution), macroscopic (system of cities)

Synthetic building layouts

At the microscopic scale (district): generating building layouts

Raimbault, J., & Perret, J. (2019, July). Generating urban morphologies at large scales. In Artificial Life Conference Proceedings (pp. 179-186). MIT Press.

- systematic comparison of simple processual generators
- introduction of morphological indicators
- calibration on sampled layouts from OpenStreetMap

Synthetic population grids

At the mesoscopic scale: population grids

- a reaction-diffusion model for population distributions
- urban form measures at the mesoscopic scale
- synthetic generators coupling population and road networks

Raimbault, J. (2018). Calibration of a density-based model of urban morphogenesis. PloS one, 13(9), e0203516.

Raimbault, J. (2019). An urban morphogenesis model capturing interactions between networks and territories. In The Mathematics of Urban Morphology (pp. 383-409). Birkhäuser, Cham.

Synthetic systems of cities


At the macroscopic scale: synthetic systems of cities

- Evolutive urban theory: systems of cities follow general stylised facts [Pumain, 2018]
- Rank-size law [Pumain et al., 2006]
- Central place theory
- ightarrow Cities-network co-evolution model explored on synthetic systems of cities
- Raimbault, J. (2021). Modeling the co-evolution of cities and networks. In Handbook of Cities and Networks (pp. 166-193). E. Elgar.

Method flowchart

General workflow to test the spatial sensitivity of simulation models

Raimbault, J., Cottineau, C., Le Texier, M., Le Nechet, F., & Reuillon, R. (2019). Space Matters: Extending Sensitivity Analysis to Initial Spatial Conditions in Geosimulation Models. Journal of Artificial Societies and Social Simulation, 22(4).



Implementation: integration into OpenMOLE

Library implemented in scala: advantages of functional and object programming; Apache Spark; no widely used GIS library in scala.

https://github.com/openmole/spatialdata

 \rightarrow integration into the OpenMOLE model exploration open source software [?]



Enables seamlessly (i) model embedding; (ii) access to HPC resources; (iii) exploration and optimization algorithms

https://openmole.org/

Benchmarking generators Noise propagation and real data perturbation Variance-based indicators Interdisciplinary systematic review

Real data perturbation

- \rightarrow How does noise in real data impacts the result ?
 - Impact of missing elements
 - Impact of imprecise coordinates or topology
 - Optimal matching between spatial datasets
- → How does perturbation of real data allows to explore scenario

Examples:

- simulating urban projects by modifying population of areas with a given spatial correlation structure
- simulating network disruptions or new transportation lines

Benchmarking generators Noise propagation and real data perturbation Variance-based indicators Interdisciplinary systematic review

Benchmarking generators Noise propagation and real data perturbation Variance-based indicators Interdisciplinary systematic review

[Koo et al., 2020]

Discussion

Developments

- more spatial network generative models [?], correlated synthetic data
 [?]
- domain models: LUTI, urban dynamics
- other disciplines, ecology, geosciences [?]?
- interaction with data driven disciplines? (planning, architecture, spatio-temporal datamining)
- genericity of some models? (reaction-diffusion)
- synthetic data generation methods (synthetic populations)
- synthetic data at the core of applied statistics methodology (less in spatial statistics?)
- port the library to more classic languages (python, R)

Conclusion

- ightarrow **Space matters:** relevance of spatially-explicit models and spatial sensitivity analysis.
- ightarrow Synthetic data: first experimental samplings included in OpenMOLE, soon more to come.
- → **Disciplinary context**: strong contingency on included models.

Get the library at https://github.com/openmole/spatialdata

Open issues at https://github.com/openmole/spatialdata/issues

References I



Environmental modelling & software, 134:104857.

Olteanu-Raimond, A.-M., Mustiere, S., and Ruas, A. (2015).

Knowledge formalization for vector data matching using belief theory.

Journal of Spatial Information Science, (10):21-46.

References II



Pumain, D. (2018).

An evolutionary theory of urban systems.

International and transnational perspectives on urban systems, pages 3–18.



Pumain, D., Paulus, F., Vacchiani-Marcuzzo, C., and Lobo, J. (2006).

An evolutionary theory for interpreting urban scaling laws.

Cybergeo: European Journal of Geography.



Raimbault, J. (2018).

Calibration of a density-based model of urban morphogenesis.

PloS one, 13(9):e0203516.

References III




An urban morphogenesis model capturing interactions between networks and territories.

The mathematics of urban morphology, pages 383-409.



Validation levels and standards depending on models types and functions.

In CCS 2019, Singapore. NTU Singapore.



Modeling the co-evolution of cities and networks.

In Handbook of Cities and Networks, pages 166–193.

References IV

Raimbault, J. (2021b).

Strong coupling between scales in a multi-scalar model of urban dynamics.

arXiv preprint arXiv:2101.12725.

Raimbault, J. (2023).

Validation of geosimulation models: a systematic review.

In ECTQG 2023, Braga. Coimbra University.

Raimbault, J., Denis, E., and Pumain, D. (2020).

Empowering urban governance through urban science: Multi-scale dynamics of urban systems worldwide.

Sustainability, 12(15):5954.

References V



Wong, D. W. (2004).

The modifiable areal unit problem (maup).

In WorldMinds: geographical perspectives on 100 problems: commemorating the 100th anniversary of the association of American geographers 1904–2004, pages 571–575. Springer.