

Spatial sensitivity of the swarm chemistry model

Author(s)¹

¹Institution

corresponding-author@email.com

Abstract

Artificial life simulation models are most of the time spatially explicit, yet there are not much dedicated studies of how spatial structure and spatial initial conditions influence model outcomes. We propose in this contribution to study the role of spatial structure in the classical model of swarm chemistry. We couple the evolutionary swarm chemistry model with spatial structure generators, and quantify the impact of space on diversity outcomes by using a diversity search algorithm. We find a significant influence of the type of spatial structure on model outcomes.

Introduction

The role of spatial structure and dynamics in evolution has been highlighted (Lion and Baalen, 2008), but remains an auxiliary focus in the literature as most models do not consider space or in very simplified settings. An evolutionary niche is not principally defined by its spatial extent, but indeed corresponds to some geographical territories, and it has furthermore been shown that spatial effects may favour niche construction (Silver and Di Paolo, 2006). From the viewpoint of Holland (2012)’s theory for complex adaptive systems, boundaries are at the core of the niche concept and thus the spatial structure of complex systems defines their dynamics. Also, biogeography optimisation algorithms have a good performance precisely because evolution is embedded in space (Simon, 2008). Therefore, a better understanding of the role of space is an important theoretical component for the discipline of Artificial Life.

Regarding Artificial Societies, and more particularly socio-spatial simulation models, a recent stream of work investigated this aspect. Raimbault et al. (2019) showed that spatial initial conditions could have as much impact on model outcomes than the parameters of the model itself. Raimbault and Perret (2023) detailed how diverse spatial configuration generators can be coupled with simulation models to quantify such effects. The term of Spatial Sensitivity Analysis was coined in that context, and a specific scala library was developed to implement these methods (Raimbault et al., 2020). A question remaining open is to what extent such effects are specific to social systems, or

do they hold in other disciplines. Furthermore in terms of broader methodological aspects, sensitivity analysis of ALife models is not widely applied despite a few examples such as (Alden et al., 2014).

This contribution reports on preliminary results aimed at investigating the spatial sensitivity in ALife simulation models. More particularly, we focus on an iconic model of ALife which is the Swarm Chemistry model (Sayama, 2009). It is a well studied model of self-organisation, and it was shown by (Sayama, 2018) that introducing evolutionary dynamics between particles could generate swarm patterns in an open-ended way. While (Sayama, 2018) uses space to introduce evolutionary pressure through perturbations and shows that evolution outcomes are strongly influenced by the magnitude of the perturbation, it remains a simple splitting of the world in half. We couple the model with more elaborate spatial structure generators for the spatial distribution of competition rules, and investigate the role of the spatial structure on diversity outcomes.

Methods

We use the evolutionary version of Swarm Chemistry described by (Sayama, 2018). The base of the model are particles which self-organise based on local interaction rules (on speed and rotation e.g.) ruled by particle-level parameters. Some mutation in parameter values and transmission between particles at collision allows some evolutionary dynamics to be introduced. As no global fitness landscape rules this evolution, different rules can be used for this “selection” step at transmission: faster particle transmits its genome, slower transmits, the majority within a local neighbourhood is taken, etc. These rules can be changed in space and time to foster evolutionary dynamics as shown by (Sayama, 2018).

We introduce a spatial context in which swarm dynamics and evolution will take place. Given a grid decomposition of the two dimensional space (in practice with a lower resolution than coordinates), each cell of the grid will contain the transmission rule for collisions occurring in this cell. We consider the rules “Faster”, “Slower”, “Behind”, “Major-

ity” and “Majority Relative” only (see (Sayama, 2018) for details), and organise the spatial distribution of rules with some structure. We use the following spatial contexts: (i) a totally random attribution of cell rule, to act as a null model; (ii) a uniform rule chosen randomly among all rules (which should also be a null model in some sense); (iii) four rectangular quadrants, with the space being split at a random x coordinate and at a random y coordinate, and with a uniform random rule within each quadrant - this configuration corresponds to neighbour regions with different evolution dynamics and exchanging particles, in a similar way to biogeography algorithms; (iv) a more realistic hierarchical organisation, inspired by the spatial organisation of systems of cities (Batty, 2008): a certain number of “urban centres” ($C = 10$ in practice) with a scaling law distribution of sizes (Zipf’s law, with a scaling exponent $\alpha_C = 1$ (Pumain et al., 2006b)) are randomly distributed in space, an exponential kernel with a radius proportional to the size is applied for each centre (monocentric cities (Lemoy and Caruso, 2020)), the distribution is rescaled as a probability distribution, a threshold ($\theta_C = 0.5$ in practice) is applied to discriminate between inside and outside of centres, and finally a randomly chosen rule is applied inside and an other random rule outside. Although inspired by urban science and patterns in social systems, we believe this last generator could also capture existing natural structures with similar hierarchical distributions, and have some genericity since hierarchy is recurrent in natural and social complex systems (Pumain et al., 2006a).

We focus on the diversity of particles as an outcome of coupling the spatial generators with swarm dynamics, as it is a key aspect for evolution and open-ended systems. We consider average and standard deviation of three particle parameters, namely c_1, c_2, c_3 which capture the main essence of each particle behaviour. This gives a 6-dimensional output, for which we seek to maximise diversity using the Pattern Search Exploration algorithm developed by (Chérel et al., 2015).

Swarm parameters are fixed at their default values used in (Sayama, 2018), except for mutation at normal time which was increased to 0.5, swarm size taken significantly smaller for performance reasons ($N = 100$), and final time $t_f = 10000$. The swarm is initialised with 10 random individuals and 90 inactive particles. The free parameters for the diversity search algorithm, run for 1000 generations, are the seed of the random number generator, and the type of spatial generator used among the four described above.

Results

We use the implementation of spatial structure generators provided by (Raimbault et al., 2020). For an easier integration with this scala library, the Swarm Chemistry model was reimplemented in scala, based on the open source Java code provided by (Sayama, 2018). Our




Figure 1: Scatterplot of PSE objectives, with points coloured by type of generator.

implementation is embedded into the OpenMOLE software for exploration and validation of simulation models (Reuillon et al., 2013), which includes the PSE diversity search algorithm. Source code and results are available on an open git repository at <https://github.com/AnonymousAuthor1/ALife2024>.

We show in Fig. 1 scatterplots of the final point cloud obtained with the PSE algorithm. The overall shape of each plot is not trivial, with reduced standard deviations for averages values of averages for each parameters. Correlation patterns are also diverse. When comparing across generators (point colour), statistical distribution are significantly changing for some dimensions, for example $\sigma(c_1)$ (which shows a unique bimodal distribution for Zipf only) and \bar{c}_2 . When looking at the higher and most significant correlations, between \bar{c}_1 and \bar{c}_3 , only uniform space is not significant and close to zero, while other are high. Altogether, these results confirm a significant effect of the spatial context on evolution outcomes in this model.

Discussion

We show the important influence of space with these preliminary results, and future work will have to focus on better understanding how qualitatively each structure influences evolution, and trying to observe recurrent evolutionary patterns such as potentially the emergence of a particle biogeography. Other types and more realistic generators for this particular system, but also the application to other disciplines such as ecology are also important future research directions.

References

- Alden, K., Timmis, J., and Coles, M. (2014). Easing parameter sensitivity analysis of netlogo simulations using spartan. In *Artificial Life Conference Proceedings*, pages 622–628. MIT Press One Rogers Street, Cambridge, MA 02142-1209, USA journals-info
- Batty, M. (2008). The size, scale, and shape of cities. *science*, 319(5864):769–771.
- Chérel, G., Cottineau, C., and Reuillon, R. (2015). Beyond corroboration: Strengthening model validation by looking for unexpected patterns. *PloS one*, 10(9):e0138212.
- Holland, J. H. (2012). *Signals and boundaries: Building blocks for complex adaptive systems*. Mit Press.
- Lemoy, R. and Caruso, G. (2020). Evidence for the homothetic scaling of urban forms. *Environment and Planning B: Urban Analytics and City Science*, 47(5):870–888.
- Lion, S. and Baalen, M. v. (2008). Self-structuring in spatial evolutionary ecology. *Ecology letters*, 11(3):277–295.
- Pumain, D., BATTY, M., GAUME, B., and David, L. (2006a). *Hierarchy in natural and social sciences*. Springer.
- Pumain, D., Paulus, F., Vacchiani-Marcuzzo, C., and Lobo, J. (2006b). An evolutionary theory for interpreting urban scaling laws. *Cybergeog: European Journal of Geography*.
- Raimbault, J., Cottineau, C., Le Texier, M., Le Néchet, F., and Reuillon, R. (2019). Space matters: Extending sensitivity analysis to initial spatial conditions in geosimulation models. *Journal of Artificial Societies & Social Simulation*, 22(4).
- Raimbault, J. and Perret, J. (2023). Spatial sensitivity analysis of social simulation models. In *CCS 2023*.
- Raimbault, J., Perret, J., and Reuillon, R. (2020). A scala library for spatial sensitivity analysis. In *28th GISRUK conference*, pages 32–36.
- Reuillon, R., Leclaire, M., and Rey-Coyrehourcq, S. (2013). Openmole, a workflow engine specifically tailored for the distributed exploration of simulation models. *Future Generation Computer Systems*, 29(8):1981–1990.
- Sayama, H. (2009). Swarm chemistry. *Artificial life*, 15(1):105–114.
- Sayama, H. (2018). Seeking open-ended evolution in swarm chemistry ii: Analyzing long-term dynamics via automated object harvesting. In *Artificial Life Conference Proceedings*, pages 59–66. MIT Press One Rogers Street, Cambridge, MA 02142-1209, USA journals-info
- Silver, M. and Di Paolo, E. (2006). Spatial effects favour the evolution of niche construction. *Theoretical Population Biology*, 70(4):387–400.
- Simon, D. (2008). Biogeography-based optimization. *IEEE transactions on evolutionary computation*, 12(6):702–713.