

# A multi-scalar model for system of cities

Juste Raimbault<sup>1,2,3</sup>

<sup>1</sup>UPS CNRS 3611 ISC-PIF, Paris, France, [juste.raimbault@polytechnique.edu](mailto:juste.raimbault@polytechnique.edu)

<sup>2</sup>CASA, UCL, London, UK

<sup>3</sup>UMR CNRS 8504 Gographie-cits, Paris, France

The modeling of urban growth is a crucial issue for the design of sustainable territorial policies, through the understanding of past urbanization processes and the forecasting of future urban trajectories. Several models have been proposed at different scales and integrating different dimensions of urban systems, such as land-use transport interaction models [1] or systems of cities models [2]. While multi-scalar models are recognized as crucial for the study of such systems [3], they remain in practice unexplored.

This contribution introduces a parsimonious multi-scalar model for systems of cities, based on simple dimensions (mainly populations) with stylized processes, but yielding an effective strong coupling between the metropolitan mesoscopic scale and the macroscopic scale of the system of cities. The model couples the spatial interaction model of [4] for the macro scale with the reaction-diffusion model for urban form studied by [5]. More precisely, urban areas viewed as a population grid are embedded into the macroscopic interaction model. To evolve populations and local urban forms, one time step consists of (i) population differences are computed by the interaction model; (ii) top-down feedback modifies parameters of mesoscopic models, given control parameters to capture typical scenarios (transit-oriented development or sprawl for diffusion, metropolization or uniformization for aggregation); (iii) local urban form are evolved at a given speed conditionally to the population variation; (iv) changes in urban form influence macroscopic interaction range (capturing the impact of local activity on global insertion), by integrating gravity flows in the area with a squared cost function making a compromise between congestion and flows.

The model is applied on synthetic systems of cities typical of a continental range (500km, hierarchy around 1, 20 cities), with initial local population grid configurations as monocentric. Parameter space is explored with the OpenMOLE model exploration software [6], eased by the implementation of the model in scala.

## References

- [1] Michael Wegener and Franz Fürst. Land-use transport interaction: state of the art. *Available at SSRN 1434678*, 2004.
- [2] Denise Pumain and Romain Reuillon. *Urban Dynamics and Simulation Models*. Springer International, 2017.
- [3] Celine Rozenblat and Denise Pumain. *Toward a Methodology for Multi-scalar Urban System Policies*, pages 385–393. Springer Singapore, Singapore, 2018.
- [4] Juste Raimbault. Indirect evidence of network effects in a system of cities. *Environment and Planning B: Urban Analytics and City Science*, page 2399808318774335, 2018.
- [5] Juste Raimbault. Calibration of a density-based model of urban morphogenesis. *PloS one*, 13(9):e0203516, 2018.
- [6] Romain Reuillon, Mathieu Leclaire, and Sebastien Rey-Coyrehourcq. Openmole, a workflow engine specifically tailored for the distributed exploration of simulation models. *Future Generation Computer Systems*, 29(8):1981–1990, 2013.