

Meta-analytic evidence that sexual selection improves population fitness

Justin G. Cally^{§*}, Devi Stuart-Fox[§] and Luke Holman[§]

[§]School of BioSciences, The University of Melbourne, Parkville, VIC, 3052, Australia

**justin.g.cally@gmail.com*

Abstract

Sexual selection has manifold ecological and evolutionary consequences, making its net effect on population fitness difficult to predict from first principles. Instead, one can empirically determine whether sexual selection increases or decreases population fitness by manipulating the opportunity for sexual selection in replicate populations, allow evolution to proceed, and then compare the evolved average fitness between the treatment groups. Here, we synthesise 459 effect sizes from 65 pertinent experimental evolution studies using meta-analysis. We find evidence that sexual selection on males tends to elevate the mean and reduce the variance for many fitness traits, especially in females evolving under stressful conditions. However, sexual selection had weaker positive effects on direct measures of population fitness, such as extinction rate and proportion of viable offspring, than on individual fitness measures with indirect or less clear relationships to population fitness. Overall, we conclude that the beneficial population-level consequences of sexual selection typically outweigh the harmful ones and that the effects of sexual selection on total selection, genetic variance, and/or evolvability can differ markedly between sexes and environments. We discuss the implications of these results for conservation and evolutionary biology.

INTRODUCTION

Sexual selection, defined as selection resulting from competition for mates or their gametes¹, is a ubiquitous evolutionary force that has profoundly shaped the natural world. As far back as Darwin², researchers have theorised that sexual selection can change the average absolute fitness of individuals in a population, henceforth termed ‘population fitness’^{3,4}. However, opinion is divided over whether the net effect on population fitness is positive or negative^{5–8}. *Prima facie*, one might predict that sexual selection would have no effect on population fitness, since it does not matter which individuals of the faster-reproducing sex (typically males) succeed in mating, so long as some do⁹. However, when genotypes with high mating or fertilisation success also have superior breeding values for traits that affect population fitness (*e.g.*, survival, parental care, female fecundity, or success in interspecific competition), sexual selection is predicted to elevate population fitness by causing a correlated response in these other traits⁸. In essence, the demographically-limiting sex (typically females) benefits from a gene pool that has been purged of harmful alleles through sexual selection on the non-limiting sex (typically males). Theoretically, the benefit to population fitness could be large^{10–12}.

Conversely, sexual selection can decrease population fitness if male sexually-selected traits are negatively genetically correlated with female fitness, producing intralocus sexual conflict^{13–18}. Additionally, sexual selection frequently favours phenotypes that reduce population fitness but benefit the individuals expressing them, such as harassment or infanticide by mate-seeking males^{19–21}, as well as investment in costly sexual signals and weaponry at the expense of parental care: *interlocus* sexual conflict^{22–24}. Given these conflicting theoretical expectations and empirical results, it remains unclear whether sexual selection tends to have a net benefit or cost to population fitness^{5–8}.

Researchers have investigated the population-level consequences of sexual selection using a range of approaches including macro-evolutionary studies^{25–27}, analysis of the fossil record²⁸, quantitative genetics^{13,29–32} and especially experimental evolution. In particular, many experimental evolution studies have manipulated the intensity of sexual selection in captive populations, allowed evolution to proceed, and then measured population fitness components such as lifespan, reproductive success, population extinction rate and mutation load. This approach facilitates direct measurement of the net effect of sexual selection on population fitness, at least in the specific populations and ecological conditions under study.

A number of factors might influence the strength and sign of the correlation between sexual selection and population fitness. First, the genetic correlation between female fecundity and male mating/fertilisation success varies in sign and magnitude between species³³ and even between conspecific populations³⁴, implying that sexual selection on males increases mean female fitness in some species and populations but not others. These inconsistencies could derive from differences in allele frequencies, or environmental differences that alter how genotype relates to phenotype and fitness. Second, it has been hypothesised that populations should display a more positive genetic correlation between male and female fitness – and thus potentially between mating/fertilisation success and population fitness – in novel or fluctuating environments, relative to stable environments^{13,32,35,36}. This is because stable environments create consistent selection, preferentially eroding genetic variation at sexually-concordant loci (*i.e.* loci where the fittest genotype is the same in both sexes) and leaving behind variation at sexually antagonistic loci. We know of no systematic reviews of this latter theory, though it has motivated several recent empirical tests^{13,29,31}, and is relevant to conservation genetics.

Here, we synthesise the empirical literature on sexual selection and population fitness using formal meta-analysis. We focus exclusively on experimental evolution studies that manipulated the presence or strength of sexual selection on males, and then measured some fitness component, since experiments provide a particularly strong test of the hypothesis that sexual selection affects the average fitness of populations. We find that sexual selection tends to improve population fitness; especially when fitness components were measured in females experiencing stressful rather than benign conditions. Additionally, we show that sexual selection tends to narrow phenotypic variance of fitness-related traits for females and mixed-sex samples in stressful conditions. These results suggest that sexual selection may be especially important for populations adapting to changing environments.

RESULTS

The effect size dataset

We retrieved 459 effect sizes from 65 studies. 92 effect sizes were collected from populations evolving under stressful conditions, while 337 were measured for those evolving in benign conditions. 189 of the effect sizes came from measurements made on males, 219 on females, and the remaining 51 from measurements of a mixed-sex sample of individuals. Most effect sizes in our dataset came from studies that manipulated sexual selection by completely removing it in one treatment via enforced random monogamy ($n = 241$); other effect sizes ($n = 218$) derived from alternative manipulations, such as changing the adult sex ratio. In total, we obtained effect sizes for 22 different fitness traits, with female reproductive success ($n = 102$) and offspring viability ($n = 56$) being the most commonly-measured traits. We classified 171 effect sizes as direct measures of population fitness, 141 as indirect, and the remaining 144 effect sizes as ambiguous (see Methods, Figure 1 and [Supplementary Table 1](#)). Specifically, we scored traits that are likely to correlate with population growth and persistence as direct (*e.g.*, female reproductive success, offspring viability, and extinction rate), those which are not necessarily correlated with population fitness but which do measure individual fitness as indirect (*e.g.*, lifespan, male mating success and ejaculate quality/production); and those for which the relationship to population fitness is unclear as ambiguous (*e.g.*, body size, mating duration, male reproductive success, and early fecundity). [Supplementary Tables 3,4](#) give a detailed description of our dataset.

Sexual selection is associated with higher mean values for most fitness components

The grand mean across all types of effect sizes (direct, indirect and ambiguous) was positive (REML $\beta = 0.24$, 95% CIs: 0.055 to 0.43, $p = 0.011$; Bayesian $\beta = 0.25$, 95% CIs: -0.0074 to 0.51, $BF_{>0} = 35$), indicating that sexual selection on males typically had a net positive effect on the majority of populations and fitness components so far studied. Moreover, the effect sizes associated with the manipulation of sexual selection varied between different fitness traits. Sexual selection had a beneficial effect on most fitness traits, but varied across the three relationships to fitness (Figure 1; [Supplementary Tables 5,6](#)). Sexual selection elevated fitness for traits that shared an ambiguous relationship to fitness (REML $\beta = 0.21$, 95% CIs: 0.058 to 0.093;

Bayesian $\beta = 0.20$, 95% CIs: -0.0016 to 0.39, $BF_{>0} = 38$, $n = 144$) and an indirect relationship to fitness (REML $\beta = 0.24$, 95% CIs: 0.13 to 0.36; Bayesian $\beta = 0.24$, 95% CIs: 0.033 to 0.43, $BF_{>0} = 59$, $n = 141$). Additionally, sexual selection elevated fitness components directly related to fitness, albeit at a lower magnitude (REML $\beta = 0.13$, 95% CIs: 0.019 to 0.24; Bayesian $\beta = 0.13$, 95% CIs: -0.079 to 0.31, $BF_{>0} = 11$, $n = 174$). A large forest plot with predicted effect sizes for each fitness component is presented in [Supplementary Figure 1](#), [Supplementary Table 7](#) and further detailed with model predictions and individual meta-analyses in [Supplementary Table 8,9](#). Sexual selection significantly reduced two fitness components, namely immunity (REML $\beta = -0.42$, 95% CIs: -0.64 to -0.20; Bayesian $\beta = -0.43$, 95% CIs: -0.70 to -0.15, $BF_{>0} = 0.0026$; $n = 35$) and body condition (REML $\beta = -1.2$, 95% CIs: -1.8 to -0.63; Bayesian $\beta = -1.2$, 95% CIs: -1.9 to -0.63, $BF_{>0} < 0.0001$, $n = 1$).

The roles of environmental stress and sex

We found that the sex of the individuals measured (male, female, or a mixture), and the conditions under which the population evolved (stressful or benign) interacted to affect the relationship between sexual selection and fitness (Table 1,2 and [Supplementary Tables 11-13](#)). Sexual selection on males significantly improved female fitness, and the beneficial effect of sexual selection was significantly stronger for females from populations evolving under stressful conditions (*e.g.* a food source to which they were not well-adapted) than under benign conditions (Figure 2a, Table 2, [Supplementary Table 14](#)). Sexual selection had a positive but non-significant effect on male fitness, and in contrast to females, fitness benefits were significantly weaker in stressful than benign environments (Figure 2a, Table 2). Consistent with the different consequences of sexual selection for female and male fitness, the mean effect size in mixed-sex samples was non-significantly positive, and there was no significant difference between benign versus stressful conditions (Figure 2a, Table 2). Overall, our results indicate that the positive effect of sexual selection on fitness is greater for females than males, and the difference between the sexes is magnified in stressful environments. When only fitness components directly related to fitness were used in the mixed-effects model the benefits of sexual selection were magnified in females evolving under stressful conditions ([Supplementary Figure 3](#) and [Supplementary Table 15](#)). Similarly, when we used an alternative measure of effect size (log response ratio or lnRR), the results aligned with those of Hedges' g in that sexual selection elevates population fitness, with its effect magnified for females evolving in stressful environments ([Supplementary Figure 4](#) and [Supplementary Table 16](#)).

Other moderator variables that we examined had minimal impacts on effect size ([Supplementary Table 9](#)). Specifically, effect size did not depend on whether or not the study was conducted blind ([Supplementary Figure 7](#)), nor on the number of generations for which the experimental evolution study was run ([Supplementary Figure 8,9](#)).

The effect size estimates we recovered were highly heterogeneous ($I^2 = 95.2\%$, 95% CIs: 94.4% to 95.9%), reflecting the large differences in experimental procedures, study species, and fitness components included in our meta-analysis^{37,38}. Heterogeneity stemmed mostly from between-study differences ($I^2_{study} = 36\%$, 95% CIs: 26.5% to 45.4%) rather than differences between fitness components and taxon ($I^2_{fitness\ components} = 0.4\%$, 95% CIs: 0.2% to 0.9%; $I^2_{taxon} = 1.4\%$, 95% CIs: 0.2% to 3.5%). Variation among taxa is explored further in the Supplementary Information ([Supplementary Figure 2](#), [Supplementary Table 10](#)).

Sexual selection reduces phenotypic variance, for female traits in stressful environments

By applying meta-analysis to log coefficient of variation ratios³⁹, we found evidence that sexual selection reduces phenotypic variation under certain conditions (Figure 2b). Specifically, phenotypic variance was significantly reduced by sexual selection for fitness components measured in females under stressful conditions ($\ln CVR = -0.78$, 95% CIs: -1.23 to -0.34, $n = 27$). By contrast, we found no significant effect of sexual selection on phenotypic variance in males, or for either sex under benign conditions (Figure 2b; [Supplementary Tables 17-20](#)). However, similar to the results in females, there was a non-significant trend for a reduction in phenotypic variance in mixed-sex samples measured under stressful conditions ($\ln CVR = -0.76$, 95% CIs: -1.22 to -0.31; Figure 2b). A meta-analysis using the log variability ratio (lnVR), which does not account for the mean-variance relationship present in the dataset ([Supplementary Figure 5](#)), suggested sexual selection reduces variance for mixed-sex samples in stressful conditions, but not females. Results of this meta-analysis can be found in the Supplementary Information ([Supplementary Figure 6 and Supplementary Table 21](#)).

As in the meta-analysis of trait means, there was high heterogeneity in the estimates of $\ln CVR$ ($I^2 = 98.9\%$, 95% CIs: 98.7% to 99.1%). Heterogeneity in the dataset was due to variability between studies fitness components ($I^2_{fitness\ components} = 12.7\%$, 95% CIs: 4.5% to 23.1%) and taxon ($I^2_{taxon} = 7.7\%$, 95% CIs: 1.3% to 18.4%), as well as large amounts of residual heterogeneity (78.6 %, 95 % CIs: 65.9% to 89%). Using the REML approach heterogeneity associated with the Study ID random effect was estimated at zero ($I^2_{study} = 0\%$), however the Bayesian approach suggests heterogeneity between studies may be zero or small ($sd_{study} = 0.07$, 95 % CIs: 0 to 0.19).

Publication bias

The funnel plot of effect sizes was asymmetrical, suggesting that some publication bias might be present (Figure 3a; Egger's test: $z = 5.9$, $p < 0.0001$). Specifically, there was a moderate excess of low-powered studies in which sexual selection had a more positive effect on the fitness component than average, implying that low-powered studies are more likely to be published if they report statistically significant fitness benefits of sexual selection (though funnel plots are not a decisive evidence of publication bias⁴⁰). Linear regressions show no significant relationship between effect size and journal impact factor (Figure 3b; $t_{437} = 1.2$, $p = 0.23$) or year of publication (Figure 3c; $t_{437} = -1.2$, $p = 0.24$); thus, we found no evidence that effect size dictates the likelihood of publication in high-profile journals, or that effect sizes have diminished as the field has matured^{41,42}.

DISCUSSION

Our meta-analysis revealed that populations evolving under sexual selection often have higher values for multiple fitness traits, relative to populations where sexual selection on males was experimentally removed or weakened. Sexual selection had beneficial effects on the majority of commonly measured individual fitness traits, which have indirect or unclear relationships to population fitness. Effects of sexual selection on direct measures of population fitness such as extinction rate, reproductive success (defined as the number of offspring produced), and the proportion of viable offspring, were smaller and variable (similar to the results of another meta-analysis⁴³), but nonetheless tended to be positive. Fitness traits related to immunocompetence were an exception: sexual selection typically resulted in weaker immunity. This result is interesting in light of the hypothesised trade-off between sexually-selected phenotypes and immunity, for example, due to immunosuppressive effects of sex hormones^{44,45}. Furthermore, the overall benefit of sexual selection was greater for females than males, and this sex difference was magnified in stressful environments. Consistent with stronger selection on female fitness under stress, female and mixed-sex samples showed reduced phenotypic variance when sexual selection was applied under stressful as opposed to benign conditions. These results suggest that sexual selection may contribute to population persistence under stressful conditions, such as fluctuating environmental change³⁵ or spatial variability⁴⁶, particularly since female reproductive output is often a limiting factor in population growth⁴⁷.

The results of the meta-analysis support predictions that sexual selection on males can improve population fitness and accelerate adaptation^{10–12,35,48–50}. One possible mechanism is that male mating success is positively genetically correlated with traits that contribute to population fitness, allowing females to benefit from a genome that has been purged of deleterious alleles through competition between males^{8,10,11}. A second (non-exclusive) mechanism is that experimental manipulation of sexual selection on males might directly alter the selective pressures acting on females, causing female traits to evolve. For example, removing sexual selection via enforced monogamy probably alters selection on females, because it alters the frequency of interactions with males (as well as the evolved genotype of those males). What is less clear is why the manipulation of sexual selection had a larger effect on female trait means and variances as opposed to males – this result is arguably the opposite of what one would expect, since it is males that experience stronger sexual selection. Below we discuss possible explanations for this result, in light of the core principle that the extent of adaptation depends on additive genetic (co)variance and the strength of selection^{51,52}.

Firstly, it is possible that female traits show more additive genetic variation than male traits, causing female traits to respond more strongly to a change in selection. This hypothesis is plausible because males frequently do experience stronger selection than females⁵³, and sustained strong selection reduces heritability. A systematic review found no overall difference in mean heritability between male and female traits⁵⁴, but did record numerous instances in which trait heritability was higher for females than males⁵⁴. The sex chromosomes provide another reason for sex-specific heritability. Males are heterogametic in most of the species in our sample (*i.e.* the species with XY or XO sex determination), which can reduce father-to-son heritability relative to mother-to-daughter, since sons do not inherit the larger sex chromosome from their fathers^{16,55–57}, potentially slowing the adaptation of male traits^{57,58}.

Secondly, selection on males might be weaker than selection on females, resulting in slower adaptation following the experimental manipulation of sexual selection. This explanation may initially seem implausible,

because net selection on males is often stronger than on females⁵³, due in part to the elevated importance of sexual selection in males as opposed to females^{59–62}. However, an oft-overlooked aspect is that selection might frequently be softer on males and harder on females⁶³, because the local competitive environment is usually more important for males than it is for females. For instance, a mediocre male genotype can have high fitness provided it outcompetes its local rivals, while low-fitness female genotypes might produce few offspring even when competing with other low-fitness females. Therefore, improvements in genetic quality might have stronger diminishing returns in males, possibly contributing to our finding that the genetic consequences of sexual selection lead to greater fitness benefits for females. Though this argument is speculative, we note that many experimental evolution designs exaggerate the sex difference in the softness of selection, relative to expectations for large, natural populations^{64–66}. For example, many studies^{67–71} have evolved insects in small sub-populations, each containing one female and multiple males, whose progeny are then mixed and randomly sampled to create the next generation; this design ensures that successful males simply needed to outcompete their rival(s) in the same sub-population (soft selection), while each female’s reproductive output is measured against the entire female population (hard selection).

Our results suggest that the greater benefit of sexual selection to females than males is magnified in stressful environments. Recent work has emphasised that environmental stress should reduce the strength of sexually antagonistic selection relative to selection that is concordant between sexes. Theoretical models reaching this conclusion^{35,72} have been supported by some empirical work^{13,36,73}; for example, one study found that high fitness males produced low fitness daughters under benign conditions but high fitness daughters under stress³⁶. However, other quantitative genetic studies have shown that stressful conditions do not always reduce sexual antagonism^{29,32,74}. Variation in effects of sexual selection in stressful environments may be due to potentially variable responses amongst taxa³³ and environments. Notably, Connallon & Hall³⁵ predict that the dynamics of environmental change alter the strength of sexual antagonism; for instance, gradual directional selection may facilitate indefinite sexual antagonism, while rapid cyclical change can swiftly remove it. Our meta-analysis suggests that under directional selection imposed by environmental stress, sexual antagonism is likely dampened; allowing sexual selection to facilitate adaptation and persistence.

Although our meta-analysis revealed an overall positive effect of sexual selection, the variation in effect size across the dataset is high, as is often the case for studies in ecology and evolution³⁷. Most of the heterogeneity was between studies (potentially due to differences between study designs and populations), while the taxon, number of generations of evolution, and use of blinding had less impact on effect size. Experimental evolution studies cover relatively few taxa, and most focus on easy-to-culture invertebrates with similar mating systems and sex determination. However, a meta-analysis of macroevolutionary studies on sexual selection and speciation rate found no significant taxon-based differences across a diverse sample of vertebrates and invertebrates (fish, insects, birds, spiders, reptiles, mammals)⁷⁵, perhaps suggesting that our results would generalise to other taxa. On the other hand, biological differences between taxa could change the relationship between sexual selection and population fitness. For example, species where males and females have radically different morphology might have a reduced inter-sexual genetic correlation for fitness, such that sexual selection has fewer pleiotropic benefits for females, while for species with sexually-selected male parental care, sexual selection might help the population by conferring high fitness to caring fathers.

Our findings have implications for fundamental and applied research. For example, the beneficial population-level consequences of sexual selection have been proposed as one possible resolution to the long-standing evolutionary puzzle regarding sexual reproduction⁷⁶. If sufficiently strong, these benefits can more than

compensate for the costs of sexual reproduction, and prevent sexual populations from being outcompeted by asexual mutants^{10,11,77}. Sexual selection is also important for conservation⁶ and captive breeding programs⁷⁸. Within captive breeding programs, genetic diversity is often managed through the enforced monogamy of a strategically selected (genetically diverse) breeding pair⁷⁸. Captive breeding programs may benefit from allowing sexual selection of ‘good genes’ or more compatible genes⁷⁹, or by increasing maternal investment by females paired with attractive males^{80–82}. Additionally, our findings imply that anthropogenic environmental changes that reduce the opportunity for sexual selection, such as eutrophication, pesticides, artificial light and noise pollution, could reduce the genetic quality of the population, and potentially compromise its long-term persistence^{83–87}. Equally, our results support recent evidence that human activities that directly counteract sexual selection, such as selective harvesting of the largest or most ornamented males, can lower population fitness⁸⁸. Based on the weight of evidence from experimental evolution, we suggest that sexually-selected populations may be more resilient to environmental change, including anthropogenic environmental pressures, over relevant time scales.

METHODS

Literature search

We searched *ISI Web of Science* and *Scopus* on 9th June 2017 for peer-reviewed, English language studies that manipulated the presence or strength of sexual selection using experimental evolution, and then measured some proxy of population fitness. A detailed list of search terms is given in the Supplementary Information (see [Supplementary Methods](#)).

After removing duplicates, we read the titles and abstracts of the remaining 1,015 papers and removed those that did not fit our inclusion criteria (typically because they did not present primary experimental evolution data). This left 130 papers, for which we read the full text and applied the inclusion criteria outlined in the PRISMA diagram (Figure 4). Briefly, we included studies that 1) were conducted in a dioecious animal, 2) experimentally manipulated the strength of sexual selection (*e.g.*, via experimentally-enforced random monogamy or an altered sex ratio) for at least one generation, and 3) measured a trait that we judged to be a potential correlate of population fitness. This third criterion is the most subjective, because there is rarely enough data to determine whether a particular trait is (or is not) correlated with population fitness. We therefore relied on our best judgement when deciding what outcomes were correlated with population fitness. We categorised the fitness outcomes into three categories: ambiguous, indirect and direct (detailed in [Supplementary Table 1](#)). Briefly, ambiguous measures of fitness were those that are reported to have an unclear or variable association with fitness (*e.g.*, body size, mating duration, early fecundity and male reproductive success). Indirect fitness components were those that are often used as a proxy of fitness but do not directly measure aspects of success in reproduction or population viability (*e.g.*, lifespan, mating success and ejaculate quality/production). Finally, direct measures of fitness (female/mixed sex reproductive success, offspring viability and extinction rate) are those that measure fitness through components of reproduction or long-term viability. The Supplementary Methods describe why each of the 130 papers was included or excluded ([Supplementary Table 2](#)).

Of these 130 papers, 62 were excluded based on the PRISMA criteria (Figure 4). Additionally, three papers

presented insufficient information to calculate effect size. In these cases, we contacted the authors and attempted to obtain the missing data, with partial success. The final meta-analysis included data from 65 papers.

Data extraction

From each paper, we first attempted to extract the arithmetic means, standard deviations and sample sizes of each of the different treatment groups, which facilitate calculation of effect size (see below). Typically, there were two or three treatments, which varied in the strength of sexual selection on males through manipulations to the adult sex ratio, in these cases we considered treatments with the greater male to female ratio to be the high sexual selection treatment group. For some papers, summary statistics were not written down, but were presented in a figure such as a bar chart: in these cases, we extracted the data using *WebPlotDigitizer* v.3.12⁸⁹. If the treatment means were not reported (and the raw data were unavailable), we instead calculated effect size from test statistics comparing treatment means (*e.g.*, F , t , z or χ^2 values), which we used to estimate effect size using several formulae (see below).

Where possible, we extracted data for each independent replicate or experimental evolution line within a study; otherwise, we used pooled treatment means. For studies that repeatedly measured the same population across multiple generations, we only extracted data for the last reported generation.

In addition to the data used to calculate effect size, we collected a set of moderator variables for each paper (see [the source data](#) and associated Supplementary Information). The moderators were selected due to their ready availability, and because we hypothesised that they might explain some of the observed heterogeneity in effect size. A key moderator was whether the environmental conditions that a population evolved under were stressful (*e.g.*, elevated mutation load, novel/sub-optimal food source, increased sub-lethal temperatures). Additionally, we collected details for each effect size on: sex (male, female or a mixed sample of both), taxon (flies, beetles, mice, nematodes, mites, crickets and guppies), the presence/absence of blind methodology, and number of generations a treatment group underwent experimental evolution. In the interests of creating a useful data resource, we also recorded details about each experiment that were not formally analysed due to a shortage of data, such as the type of sexual selection that was manipulated (pre-copulatory, post-copulatory, or both) and the male to female ratio, which is included in the Supplementary Information ([the source data](#)).

Effect size calculation

For each measurement of each pair of treatments, we estimated the standardised effect size Hedges' g ⁹⁰. Similar to Cohen's d , Hedges' g expresses the difference in means in terms of standard deviations (making it dimensionless), but it is more robust to unequal sampling and small sample sizes⁹¹. For comparisons of extracted treatment means, we calculated Hedges' g using the `mes` function in the *compute.es* R package⁹². To calculate Hedges' g from test statistics, we used the `fes`, `chies`, and `tes` functions in the *compute.es* package (for F , χ^2 and t statistics, respectively). The `propes` function was used to calculate effect size from a difference in proportions; in two cases^{93,94}, a proportion was equal to one (producing infinite effect sizes), and so we subtracted one from the numerator when estimating Hedges' g . In all cases, we selected a direction for the effect size calculation such that in our meta-analysis, negative effect sizes indicate that the removal

of sexual selection was associated with higher fitness trait values, and positive effect sizes indicate higher fitness when sexual selection was elevated or left intact. We also inverted the sign of effect sizes pertaining to measurements that are expected to be negatively related to population fitness (*e.g.*, parasite load, mutation load, extinction risk/rate, mating latency (males), and rate of senescence). Because many of our 65 papers measured multiple fitness outcomes, studied multiple replicate populations, or had three or more sexual selection treatments, we calculated a total of 459 effect sizes.

Additionally, using studies that presented means, standard deviations and sample sizes ($n = 352$) we were able to calculate an alternative measure of effect size: the log response ratio ($lnRR$)^{95,96}. The $lnRR$ was used as a supplement to Hedges' g because it relaxes the assumption in equal variances between control and treatment groups (homoscedasticity).

For the meta-analysis testing whether sexual selection affects phenotypic variance (as opposed to the mean), we estimated the difference in variance between each pair of treatments using the natural logarithm of the ratio between the coefficient of variation for each group (termed $lnCVR$)³⁹: $ln(CV fitness_{SS\ high}/CV fitness_{SS\ low})$. The use of $lnCVR$ allows us to determine the effects of sexual selection on phenotypic variance, with the coefficient of variation implicitly controlling for the mean-variance relationship seen in the dataset (Supplementary Figure 5). As a supplement, we also calculated the natural logarithm of the absolute ratio between the absolute variation for each group ($lnVR$) in order to assess the impact of sexual selection on trait variance, irrespective of their magnitudes³⁹. The calculation of $lnCVR$ and $lnVR$ relies on the availability of arithmetic means, standard deviations and sample sizes for the two treatment groups^{39,97}, and so we were only able to calculate $lnCVR$ and $lnVR$ for 354 of 459 comparisons.

Mixed-effects meta-analysis

Firstly, we obtained a weighted mean effect size (Hedges' g) for the entire dataset, using both Bayesian and restricted maximum likelihood (REML) approaches for completeness. The weighted mean was obtained by fitting a model with no moderator variables (*i.e.* fixed effects), but fitness component (*e.g.*, body size, female reproductive success), study ID, and taxon as random/group-level effects. That is, we separately model correlations between different effect sizes sourced from the same study, taxon, or pertaining to the same fitness component, and account for these interdependencies when estimating the overall effect. Given the small number of phylogenetically diverse species, we did not utilise phylogenetic corrections within the models. In our meta-analyses we report Bayes Factors (BF), giving the likelihood ratio that the focal effect size differs from zero $BF_{>0}$.

Secondly, we fixed the relationship to fitness class (Ambiguous, Indirect or Direct) as a moderator variable in Bayesian and REML models (whilst maintaining study and taxon as group-level effects) to derive predictions for effect size within each of the three fitness-relationship classes, using the relevant `predict` functions for each of the R packages used (see below). This meta-analysis was then supplemented by another model where we fixed fitness component as a moderator variable (*e.g.*, immunity, lifespan, offspring viability and female reproductive success); predictions for this model on the 22 fitness components were derived as above. Alternatively, to assess the impact of sexual selection on each fitness component independently of one another we conducted separate meta-analyses ($n = 18$); subset for each fitness trait with more than three effect sizes. These models were were intercept only REML models with study and taxon as group-level effects. Further

358 details on model parameters can be found by accessing the “R” code.

359 Thirdly, we measured the impact of environment, sex and their interaction on the effect size (Hedges’ g ,
360 $\ln RR$, $\ln CVR$ and $\ln VR$) associated with the manipulation of sexual selection, by fitting these predictors
361 as moderators in a pair of separate mixed-effects meta-analyses. These meta-analyses were restricted to
362 effect sizes calculated from unambiguous outcomes (*i.e.* those scored as being directly or indirectly related
363 to population fitness), as well as those where we were able to define the environmental conditions as either
364 stressful or benign (Hedges’ g : $n = 330$; $\ln RR$, $\ln CVR$ & $\ln VR$: $n = 269$). We again fit study ID, fitness
365 component, and taxon as random/group level effects. Models investigating other moderators such as number
366 of generations and blinding are presented in [Supplementary Table 9](#).

367 For our meta-analyses investigating the effects of environment and sex on the magnitude and variance of
368 fitness-related traits we provide estimates of heterogeneity present in the dataset. We use the statistic I^2 as an
369 estimate of the proportion of variance in effect size that is due to differences between levels of a random effect
370 (*e.g.* studies)⁹⁸. I^2 is preferred over other statistics as it is independent of sample size, is easily interpretable,
371 and can be partitioned between random effects³⁷. Within ecology and evolution heterogeneity in datasets is
372 often high, with the mean I^2 from 86 studies above 90%³⁸.

373 Meta-analyses fit by REML were implemented in the *metafor* R package⁹⁹, while their Bayesian equivalents
374 used the R package *brms* to run models in Stan¹⁰⁰.

375 Publication bias

376 We tested for publication bias via funnel plots, using Egger’s test to quantify plot asymmetry^{101,102}. Addi-
377 tionally, we tested for time-lag bias⁴¹, in which effect size magnitudes decline over time as more data are
378 collected. Additionally, we assessed a potential source of publication bias through the correlation between
379 effect size and journal impact factor⁴², which can arise if null or countervailing results are more difficult to
380 publish (impact factors were from *InCites Journal Citation Reports*).

381 Code availability

382 The code used to perform this meta-analysis is freely available on Github ([https://justincally.github.io/](https://justincally.github.io/SexualSelection/)
383 [SexualSelection/](https://justincally.github.io/SexualSelection/)). Here the code is available through a neatly compiled html, with the R Markdown file
384 found within the corresponding Github repository: (<https://github.com/JustinCally/SexualSelection>).

385 Data availability

386 All data are freely available on Github (<https://github.com/JustinCally/SexualSelection>). The repository
387 provides a source data file that was used to create all results, tables and figures.

REFERENCES

1. Andersson, M. B. *Sexual selection*. (Princeton University Press, 1994).
2. Darwin, C. *The descent of man and selection in relation to sex*. **1**, (John Murray, 1871).
3. Kimura, M. On the change of population fitness by natural selection. *Heredity* **12**, 145–167 (1958).
4. Lande, R. Natural selection and random genetic drift in phenotypic evolution. *Evolution* **30**, 314–334 (1976).
5. Candolin, U. & Heuschele, J. Is sexual selection beneficial during adaptation to environmental change? *Trends in Ecology & Evolution* **23**, 446–452 (2008).
6. Holman, L. & Kokko, H. The consequences of polyandry for population viability, extinction risk and conservation. *Philosophical Transactions of the Royal Society B-Biological Sciences* **368**, (2013).
7. Kokko, H. & Brooks, R. Sexy to die for? Sexual selection and the risk of extinction. *Annales Zoologici Fennici* **40**, 207–219 (2003).
8. Whitlock, M. C. & Agrawal, A. F. Purging the genome with sexual selection: Reducing mutation load through selection on males. *Evolution* **63**, 569–582 (2009).
9. Rankin, D. J. & Kokko, H. Do males matter? The role of males in population dynamics. *Oikos* **116**, 335–348 (2007).
10. Agrawal, A. F. Sexual selection and the maintenance of sexual reproduction. *Nature* **411**, 692–695 (2001).
11. Siller, S. Sexual selection and the maintenance of sex. *Nature* **411**, 689–692 (2001).
12. Whitlock, M. C. Fixation of new alleles and the extinction of small populations: Drift load, beneficial alleles, and sexual selection. *Evolution* **54**, 1855–1861 (2000).
13. Berger, D. *et al.* Intralocus sexual conflict and environmental stress. *Evolution* **68**, 2184–2196 (2014).
14. Bonduriansky, R. & Chenoweth, S. F. Intralocus sexual conflict. *Trends in Ecology & Evolution* **24**, 280–8 (2009).
15. Pennell, T. M. & Morrow, E. H. Two sexes, one genome: The evolutionary dynamics of intralocus sexual conflict. *Ecology and Evolution* **3**, 1819–1834 (2013).
16. Pischedda, A. & Chippindale, A. K. Intralocus sexual conflict diminishes the benefits of sexual selection. *PLOS Biology* **4**, e356 (2006).
17. Harano, T., Okada, K., Nakayama, S., Miyatake, T. & Hosken, D. J. Intralocus sexual conflict unresolved by sex-limited trait expression. *Current Biology* **20**, 2036–2039 (2010).
18. Plesnar Bielak, A., Skrzynecka, A. M., Miler, K. & Radwan, J. Selection for alternative male reproductive tactics alters intralocus sexual conflict. *Evolution* **68**, 2137–2144 (2014).
19. Berger, D. *et al.* Intralocus sexual conflict and the tragedy of the commons in seed beetles. *The American Naturalist* **188**, E98–E112 (2016).
20. Chenoweth, S. F., Appleton, N. C., Allen, S. L. & Rundle, H. D. Genomic evidence that sexual selection

- impedes adaptation to a novel environment. *Current Biology* **25**, 1860–1866 (2015).
21. Rankin, D. J., Dieckmann, U. & Kokko, H. Sexual conflict and the tragedy of the commons. *The American Naturalist* **177**, 780–791 (2011).
22. Fromhage, L., Elgar, M. A. & Schneider, J. M. Faithful without care: The evolution of monogyny. *Evolution* **59**, 1400–1405 (2007).
23. Kokko, H. & Jennions Michael, D. Parental investment, sexual selection and sex ratios. *Journal of Evolutionary Biology* **21**, 919–948 (2008).
24. Trivers, R. *Parental investment and sexual selection*. **136**, (Biological Laboratories, Harvard University, 1972).
25. Doherty, P. F. *et al.* Sexual selection affects local extinction and turnover in bird communities. *Proceedings of the National Academy of Sciences of the United States of America* **100**, 5858–5862 (2003).
26. Morrow, E. H. & Fricke, C. Sexual selection and the risk of extinction in mammals. *Proceedings of the Royal Society B-Biological Sciences* **271**, 2395–2401 (2004).
27. Morrow, E. H. & Pitcher, T. E. Sexual selection and the risk of extinction in birds. *Proceedings of the Royal Society B-Biological Sciences* **270**, 1793–1799 (2003).
28. Martins, M. J. F., Puckett, T. M., Lockwood, R., Swaddle, J. P. & Hunt, G. High male sexual investment as a driver of extinction in fossil ostracods. *Nature* **556**, 366–369 (2018).
29. Holman, L. & Jacomb, F. The effects of stress and sex on selection, genetic covariance, and the evolutionary response. *Journal of Evolutionary Biology* **30**, 1898–1909 (2017).
30. Lewis, Z., Wedell, N. & Hunt, J. Evidence for strong intralocus sexual conflict in the Indian meal moth, *Plodia interpunctella*. *Evolution* **65**, 2085–2097 (2011).
31. Martinossi-Alilibert, I., Arnqvist, G. & Berger, D. Sex-specific selection under environmental stress in seed beetles. *Journal of Evolutionary Biology* **30**, 161–173 (2017).
32. Martinossi-Alilibert, I. *et al.* The consequences of sexual selection in well-adapted and maladapted populations of bean beetles. *Evolution* **72**, 518–530 (2017).
33. Poissant, J., Wilson Alastair, J. & Coltman David, W. Sex-specific genetic variance and the evolution of sexual dimorphism: A systematic review of cross-sex genetic correlations. *Evolution* **64**, 97–107 (2010).
34. Collet, J. M. *et al.* Rapid evolution of the intersexual genetic correlation for fitness in *Drosophila melanogaster*. *Evolution* **70**, 781–795 (2016).
35. Connallon, T. & Hall, M. D. Genetic correlations and sex-specific adaptation in changing environments. *Evolution* **70**, 2198 (2016).
36. Long, T. A. F., Agrawal, A. F. & Rowe, L. The effect of sexual selection on offspring fitness depends on the nature of genetic variation. *Current Biology* **22**, 204–208 (2012).
37. Nakagawa, S., Noble, D. W. A., Senior, A. M. & Lagisz, M. Meta-evaluation of meta-analysis: Ten appraisal questions for biologists. *BMC Biology* **15**, 18 (2017).
38. Senior, A. M. *et al.* Heterogeneity in ecological and evolutionary meta-analyses: Its magnitude and

458 implications. *Ecology* **97**, 3293–3299 (2016).

459 39. Nakagawa, S. *et al.* Meta-analysis of variation: Ecological and evolutionary applications and beyond.
460 *Methods in Ecology and Evolution* **6**, 143–152 (2015).

461 40. Lau, J., Ioannidis, J. P. A., Terrin, N., Schmid, C. H. & Olkin, I. Evidence based medicine: The case of
462 the misleading funnel plot. *BMJ: British Medical Journal* **333**, 597 (2006).

463 41. Jennions, M. D. & Møller, A. P. Relationships fade with time: A meta-analysis of temporal trends in
464 publication in ecology and evolution. *Proceedings: Biological Sciences* **269**, 43–48 (2002).

465 42. Murtaugh, P. A. Journal quality, effect size, and publication bias in meta-analysis. *Ecology* **83**, 1162–1166
466 (2002).

467 43. Prokop, Z. M., Michalczyk, Ł., Drobnia, S. M., Herdegen, M. & Radwan, J. Meta-analysis suggests
468 choosy females get sexy sons more than ‘good genes’. *Evolution* **66**, 2665–2673 (2012).

469 44. Folstad, I. & Karter, A. J. Parasites, bright males, and the immunocompetence handicap. *The American*
470 *Naturalist* **139**, 603–622 (1992).

471 45. Foo, Y. Z., Nakagawa, S., Rhodes, G. & Simmons, L. W. The effects of sex hormones on immune function:
472 A meta-analysis. *Biological Reviews* **92**, 551–571 (2016).

473 46. Harts, A. M. F., Schwanz, L. E. & Kokko, H. Demography can favour female-advantageous alleles.
474 *Proceedings of the Royal Society B: Biological Sciences* **281**, (2014).

475 47. Crowley, P. H. Sexual dimorphism with female demographic dominance: Age, size, and sex ratio at
476 maturation. *Ecology* **81**, 2592–2605 (2000).

477 48. Lorch, P. D., Proulx, S., Rowe, L. & Day, T. Condition-dependent sexual selection can accelerate
478 adaptation. *Evolutionary Ecology Research* **5**, 867–881 (2003).

479 49. Proulx, S. R. Mating systems and the evolution of niche breadth. *The American Naturalist* **154**, 89–98
480 (1999).

481 50. Proulx, S. R. Niche shifts and expansion due to sexual selection. *Evolutionary Ecology Research* **4**,
482 351–369 (2002).

483 51. Blows, M. W. & Hoffmann, A. A. A reassessment of genetic limits to evolutionary change. *Ecology* **86**,
484 1371–1384 (2005).

485 52. Lande, R. Quantitative genetic-analysis of multivariate evolution, applied to brain - body size allometry.
486 *Evolution* **33**, 402–416 (1979).

487 53. Singh, A. & Punzalan, D. The strength of sex-specific selection in the wild. *Evolution* **72**, 2818–2824
488 (2018).

489 54. Wyman, M. J. & Rowe, L. Male bias in distributions of additive genetic, residual, and phenotypic
490 variances of shared traits. *The American Naturalist* **184**, 326–337 (2014).

491 55. Connallon, T. Genic capture, sex linkage, and the heritability of fitness. *The American Naturalist* **175**,
492 564–576 (2010).

493 56. Hastings, I. M. Manifestations of sexual selection may depend on the genetic basis of sex determination.

494 *Proceedings of the Royal Society of London. Series B: Biological Sciences* **258**, 83 (1994).

495 57. Reeve, H. K. & Pfennig, D. W. Genetic biases for showy males: Are some genetic systems especially
496 conducive to sexual selection? *Proceedings of the National Academy of Sciences* **100**, 1089 (2003).

497 58. Sturgill, D., Zhang, Y., Parisi, M. & Oliver, B. Demasculinization of X chromosomes in the *Drosophila*
498 genus. *Nature* **450**, 238–241 (2007).

499 59. Agrawal, A. F. Are males the more 'sensitive' sex? *Heredity* **107**, 20–1 (2011).

500 60. Mallet, M. A. & Chippindale, A. K. Inbreeding reveals stronger net selection on *Drosophila melanogaster*
501 males: Implications for mutation load and the fitness of sexual females. *Heredity* **106**, 994–1002 (2011).

502 61. Mallet, M. A., Bouchard, J. M., Kimber, C. M. & Chippindale, A. K. Experimental mutation-accumulation
503 on the X chromosome of *Drosophila melanogaster* reveals stronger selection on males than females. *BMC*
504 *Evolutionary Biology* **11**, 156 (2011).

505 62. Sharp, N. P. & Agrawal, A. F. Male-biased fitness effects of spontaneous mutations in *Drosophila*
506 *melanogaster*. *Evolution* **67**, 1189–1195 (2012).

507 63. Li, X.-Y. & Holman, L. Evolution of female choice under intralocus sexual conflict and genotype-
508 by-environment interactions. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**,
509 20170425 (2018).

510 64. Gavrillets, S. Rapid evolution of reproductive barriers driven by sexual conflict. *Nature* **403**, 886 (2000).

511 65. Kokko, H. & Rankin, D. J. Lonely hearts or sex in the city? Density-dependent effects in mating systems.
512 *Philosophical Transactions of the Royal Society B: Biological Sciences* **361**, 319–334 (2006).

513 66. Martin, O. Y. & Hosken, D. J. Costs and benefits of evolving under experimentally enforced polyandry or
514 monogamy. *Evolution* **57**, 2765–2772 (2003).

515 67. Crudgington, H. S., Beckerman, A. P., Brustle, L., Green, K. & Snook, R. R. Experimental removal and
516 elevation of sexual selection: Does sexual selection generate manipulative males and resistant females? *The*
517 *American Naturalist* **165**, S72–S87 (2005).

518 68. Crudgington, H. S., Fellows, S. & Snook, R. R. Increased opportunity for sexual conflict promotes harmful
519 males with elevated courtship frequencies. *Journal of Evolutionary Biology* **23**, 440–446 (2010).

520 69. Crudgington, H. S., Fellows, S., Badcock, N. S. & Snook, R. R. Experimental manipulation of sexual
521 selection promotes greater male mating capacity but does not alter sperm investment. *Evolution* **63**, 926–938
522 (2009).

523 70. Holland, B. & Rice, W. R. Experimental removal of sexual selection reverses intersexual antagonistic
524 coevolution and removes a reproductive load. *Proceedings of the National Academy of Sciences of the United*
525 *States of America* **96**, 5083–5088 (1999).

526 71. Pitnick, S., Miller, G. T., Reagan, J. & Holland, B. Males' evolutionary responses to experimental removal
527 of sexual selection. *Proceedings of the Royal Society B-Biological Sciences* **268**, 1071–1080 (2001).

528 72. Connallon, T. The geography of sex-specific selection, local adaptation, and sexual dimorphism. *Evolution*

529 **69**, 2333–2344 (2015).

530 73. Punzalan, D., Delcourt, M. & Rundle, H. D. Comparing the intersex genetic correlation for fitness across
531 novel environments in the fruit fly, *Drosophila serrata*. *Heredity* **112**, 143 (2013).

532 74. Delcourt, M., Blows, M. W. & Rundle, H. D. Sexually antagonistic genetic variance for fitness in an
533 ancestral and a novel environment. *Proceedings of the Royal Society B: Biological Sciences* **276**, 2009 (2009).

534 75. Kraaijeveld, K., Kraaijeveld-Smit, F. J. L. & Maan, M. E. Sexual selection and speciation: The
535 comparative evidence revisited. *Biological Reviews* **86**, 367–377 (2011).

536 76. Lehtonen, J., Jennions, M. D. & Kokko, H. The many costs of sex. *Trends in Ecology & Evolution* **27**,
537 172–8 (2012).

538 77. Kleiman, M. & Hadany, L. The evolution of obligate sex: The roles of sexual selection and recombination.
539 *Ecology and Evolution* **5**, 2572–83 (2015).

540 78. Charge, R., Teplitsky, C., Sorci, G. & Low, M. Can sexual selection theory inform genetic management of
541 captive populations? A review. *Evolutionary Applications* **7**, 1120–1133 (2014).

542 79. Russell, T. *et al.* MHC diversity and female age underpin reproductive success in an Australian icon; the
543 Tasmanian Devil. *Scientific Reports* **8**, 4175 (2018).

544 80. Cunningham, E. J. A. & Russell, A. F. Egg investment is influenced by male attractiveness in the mallard.
545 *Nature* **404**, 74 (2000).

546 81. Loyau, A. & Lacroix, F. Watching sexy displays improves hatching success and offspring growth through
547 maternal allocation. *Proceedings of the Royal Society B: Biological Sciences* **277**, 3453 (2010).

548 82. Loyau, A., Saint Jalme, M., Mauget, R. & Sorci, G. Male sexual attractiveness affects the investment
549 of maternal resources into the eggs in peafowl (*Pavo cristatus*). *Behavioral Ecology and Sociobiology* **61**,
550 1043–1052 (2007).

551 83. Arellano-Aguilar, O. & Macias Garcia, C. Exposure to pesticides impairs the expression of fish ornaments
552 reducing the availability of attractive males. *Proceedings of the Royal Society B: Biological Sciences* **275**,
553 1343–1351 (2008).

554 84. Botha, L. M., Jones, T. M. & Hopkins, G. R. Effects of lifetime exposure to artificial light at night on
555 cricket (*Teleogryllus commodus*) courtship and mating behaviour. *Animal Behaviour* **129**, 181–188 (2017).

556 85. Candolin, U., Salesto, T. & Evers, M. Changed environmental conditions weaken sexual selection in
557 sticklebacks. *Journal of Evolutionary Biology* **20**, 233–239 (2007).

558 86. Swaddle, J. P. & Page, L. C. High levels of environmental noise erode pair preferences in zebra finches:
559 Implications for noise pollution. *Animal Behaviour* **74**, 363–368 (2007).

560 87. Wong, B. B. M., Candolin, U. & Lindström, K. Environmental deterioration compromises socially enforced
561 signals of male quality in three-spined sticklebacks. *The American Naturalist* **170**, 184–189 (2007).

562 88. Knell, R. J. & Martínez-Ruiz, C. Selective harvest focused on sexual signal traits can lead to extinction
563 under directional environmental change. *Proceedings of the Royal Society B: Biological Sciences* **284**, (2017).

564 89. Rohatgi, A. WebPlotDigitalizer: HTML5 based online tool to extract numerical data from plot images.

- 565 <https://automeris.io/WebPlotDigitizer/> **Version 3.17**, (2011).
- 566 90. Hedges, L. V. & Olkin, I. *Statistical methods for meta-analysis*. (Academic Press, Inc, 1985).
- 567 91. Rosenberg, M. S., Rothstein, H. R. & Gurevitch, J. Effect sizes: Conventional choices and calculations.
568 in *Handbook of meta-analysis in ecology and evolution* (eds. Koricheva, J., Gurevitch, J. & Mengersen, K.) p.
569 61–71 (Princeton University Press, 2013).
- 570 92. Del Re, M. Compute.es: Compute effect sizes. *R Package* **0.2-4**, (2013).
- 571 93. Firman, R. C., Gomendio, M., Roldan, E. R. S. & Simmons, L. W. The coevolution of ova defensiveness
572 with sperm competitiveness in house mice. *The American Naturalist* **183**, 565–572 (2014).
- 573 94. Plesnar-Bielak, A., Skrzynecka, A. M., Prokop, Z. M. & Radwan, J. Mating system affects population
574 performance and extinction risk under environmental challenge. *Proceedings of the Royal Society B-Biological*
575 *Sciences* **279**, 4661–4667 (2012).
- 576 95. Hedges, L. V., Gurevitch, J. & Curtis, P. S. The meta-analysis of response ratios in experimental ecology.
577 *Ecology* **80**, 1150–1156 (1999).
- 578 96. Osenberg, C. W., Sarnelle, O. & Cooper, S. D. Effect size in ecological experiments: The application of
579 biological models in meta-Analysis. *The American Naturalist* **150**, 798–812 (1997).
- 580 97. Senior, A. M., Gosby, A. K., Lu, J., Simpson, S. J. & Raubenheimer, D. Meta-analysis of variance: An
581 illustration comparing the effects of two dietary interventions on variability in weight. *Evolution, Medicine,*
582 *and Public Health* **2016**, 244–255 (2016).
- 583 98. Higgins, J. P. T. & Thompson, S. G. Quantifying heterogeneity in a meta-analysis. *Statistics in Medicine*
584 **21**, 1539–1558 (2002).
- 585 99. Viechtbauer, W. Conducting meta-analyses in R with the metafor package. *Journal of Statistical Software*
586 **36**, 1–48 (2010).
- 587 100. Bürkner, P.-C. Brms: An R package for Bayesian multilevel models using Stan. *Journal of Statistical*
588 *Software* **80**, 1–28 (2016).
- 589 101. Egger, M., G., D. S., Schneider, M. & Minder, C. Bias in meta-analysis detected by a simple, graphical
590 test. *BMJ* **315**, 629–634 (1997).
- 591 102. Sterne, J. A. C. & Egger, M. Regression methods to detect publication and other bias in meta-analysis.
592 in *Publication bias in meta-analysis* (eds. Rothstein, H. R., Sutton, A. J. & Borenstein, M.) **1**, 99–110 (John
593 Wiley & Sons, Ltd, 2005).

594 ACKNOWLEDGEMENTS

595 We are very grateful to Paco Garcia-Gonzalez for sharing a collection of literature that helped to guide our
596 study, and to Tim Connallon for comments on the manuscript.

597 AUTHOR CONTRIBUTIONS

598 All authors designed the study; JGC collected the data; JGC and LH conducted the meta-analyses; JGC
599 wrote the first draft of the manuscript; and all authors contributed substantially to all further revisions.

600 COMPETING INTERESTS

601 The authors declare no competing interests.

TABLES AND FIGURE LEGENDS

Table 1: Multilevel meta-analysis model results

Parameters	Estimate	SE	LCI	UCI	z	p
REML Model						
Intercept	0.19	0.12	-0.05	0.43	1.53	0.13
Both Sexes	0	0.07	-0.14	0.15	0.04	0.97
Female Sex	0.11	0.03	0.05	0.17	3.73	0.00
Stressed Environment	-0.16	0.04	-0.24	-0.07	-3.63	0.00
Both Sexes x Stressed Environment	0.18	0.09	0.01	0.36	2.07	0.04
Female Sex x Stressed Environment	0.26	0.05	0.16	0.37	5.11	0.00
Bayesian Model						
Intercept	0.188	0.18	-0.17	0.52		
Both Sexes	0.003	0.07	-0.14	0.14		
Female Sex	0.113	0.03	0.05	0.17		
Stressed Environment	-0.156	0.04	-0.24	-0.07		
Both Sexes x Stressed Environment	0.182	0.09	0.01	0.35		
Female Sex x Stressed Environment	0.264	0.05	0.16	0.37		

* Moderator variables whose 95% confidence intervals do not cross zero are shown in bold.

† Hedges' g is the response variable for this model

Table 2: Hypothesis tests showing how sex and environmental stress interact to modulate effect size

Condition	Test	Statistical Approach	Estimate	SE	LCI	UCI
In benign environments	Female > Male	REML	0.113	0.03	0.05	0.17
		Bayesian	0.113	0.03	0.05	0.17
In stressful environments		REML	0.377	0.05	0.29	0.47
		Bayesian	0.377	0.05	0.29	0.47
For Females	Stressful > Benign	REML	0.108	0.04	0.04	0.18
		Bayesian	0.109	0.04	0.04	0.18
For Males	Benign > Stressful	REML	0.156	0.04	0.07	0.24
		Bayesian	0.156	0.04	0.07	0.24
For Both	Stressful > Benign	REML	0.028	0.08	-0.13	0.18
		Bayesian	0.026	0.08	-0.13	0.18

* Moderator variables whose 95% confidence intervals do not cross zero are shown in bold.

† Hedges' g is the response variable for this model

Figure 1: The effect of sexual selection on direct, indirect and ambiguous fitness components. The effect sizes used in this meta-analysis ($n = 459$) were grouped into either direct, indirect or ambiguous measures of fitness. Overall, effect sizes were more often positive than negative. Predicted average values are presented as a diamond with 95 % CIs for each fitness-relationship category. The estimates presented here are from REML models with the grand mean across all effect sizes ($\beta = 0.25$) shown as the blue dotted line. Predictions from both Bayesian and REML models can be found in Supplementary Table 6 and source data are provided at <https://github.com/JustinCally/SexualSelection>.

Figure 2: The roles of environmental stress and sex. (a) Sexual selection tends to increase the population mean values of fitness traits, especially for female traits and for populations living under stressful conditions. (b) Under stressful conditions, sexual selection tends to reduce the phenotypic variance in fitness traits, especially for traits measured in females or mixed-sex individuals. The points with error bars show the mean effect sizes and their 95% CIs, determined from a meta-regression fit using restricted maximum likelihood (REML); the point sizes are proportional to the number of effect sizes (see Supplementary Tables 14,18). Results from Bayesian meta-regression are shown as posterior prediction density curves, with vertical lines indicating the median. Source data are provided at <https://github.com/JustinCally/SexualSelection>.

Figure 3: Tests for publication bias in the dataset. Tests of publication bias are mixed and suggest publication bias may be present. (a) Inspection and statistical tests of the funnel plot reveal large amounts of heterogeneity in the dataset with asymmetry from increased low-powered, large effect studies. (b) No significant correlation exists between journal impact factor and effect size, (c) additionally there is no significant correlation between effect size and year of publication when testing for the time-lag bias. For (b) and (c) point size is proportional to the precision of the effect size (*i.e.* the inverse of its variance). For (a), the dashed red lines represent the 95 % CIs of the grand mean estimate for all effect sizes and the black dotted and dashed lines depict the 95 % and 99.8 % CIs for the dataset. The grey envelopes in (b) and (c) represent the 95 % CI of the linear regression. Source data are provided at <https://github.com/JustinCally/SexualSelection>.

Figure 4: PRISMA diagram. Flow of inclusion and exclusion of studies identified during the literature search, presented as a PRISMA diagram with number of published papers in brackets.