



Taylor & Francis  
Taylor & Francis Group



---

## A Simple Proof of Sion's Minimax Theorem

Author(s): Jürgen Kindler

Source: *The American Mathematical Monthly*, Apr., 2005, Vol. 112, No. 4 (Apr., 2005), pp. 356-358

Published by: Taylor & Francis, Ltd. on behalf of the Mathematical Association of America

Stable URL: <https://www.jstor.org/stable/30037472>

### REFERENCES

Linked references are available on JSTOR for this article:

[https://www.jstor.org/stable/30037472?seq=1&cid=pdf-reference#references\\_tab\\_contents](https://www.jstor.org/stable/30037472?seq=1&cid=pdf-reference#references_tab_contents)

You may need to log in to JSTOR to access the linked references.

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



JSTOR

Taylor & Francis, Ltd. and Mathematical Association of America are collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*

only if

$$\begin{vmatrix} \sigma_2 & \sigma_3 & \dots & \dots & \dots & \sigma_{m+1} \\ \sigma_3 & \sigma_4 & \dots & \dots & \dots & \sigma_{m+2} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \sigma_{m+1} & \sigma_{m+2} & \dots & \dots & \dots & \sigma_{2m} \end{vmatrix} = 0 \quad (N = 2m + 1)$$

and

$$\begin{vmatrix} \sigma_3 & \sigma_4 & \dots & \dots & \dots & \sigma_{m+2} \\ \sigma_4 & \sigma_5 & \dots & \dots & \dots & \sigma_{m+3} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \sigma_{m+2} & \sigma_{m+3} & \dots & \dots & \dots & \sigma_{2m+1}, \end{vmatrix} = 0 \quad (N = 2m + 2),$$

where  $\sigma_j$  are the coefficients in the formal expansion of the function

$$f(t) = \sqrt{\left(t + \frac{1}{a^2}\right) \left(t + \frac{1}{b^2}\right) \left(t + 1 - \frac{c^2}{a^2} - \frac{d^2}{b^2}\right) + t \left(\frac{c^2}{a^4} + \frac{d^2}{b^4}\right) + \frac{c^2}{a^4 b^2} + \frac{d^2}{a^2 b^4}}$$

into a power series  $f(t) = \sum_{j=0}^{\infty} \sigma_j t^j$ .

# REFERENCES

1. M. Berger, *Geometry II*, Springer-Verlag, Berlin, 1996.
2. B. Mirman, UB-matrices and conditions for Poncelet polygons to be closed, *Linear Algebra Appl.* **360** (2003) 123–150.

*Department of Mathematics and Computer Science, Suffolk University, Boston, MA 02114-4280*  
*bmirman@rcn.com*

# A Simple Proof of Sion’s Minimax Theorem

Jürgen Kindler

The following theorem due to Sion [3] is fundamental in convex analysis and in the theory of games.

**Theorem.** *Let  $X$  and  $Y$  be nonvoid convex and compact subsets of two linear topological spaces, and let  $f : X \times Y \rightarrow \mathbb{R}$  be a function that is upper semicontinuous and quasi-concave in the first variable and lower semicontinuous and quasi-convex in the second variable. Then*

$$\min_{y \in Y} \max_{x \in X} f(x, y) = \max_{x \in X} \min_{y \in Y} f(x, y).$$

The assumptions on  $f$  mean that for every real  $\alpha$  the upper sets  $U_\alpha(y) := \{x \in X : f(x, y) \geq \alpha\}$  for  $y$  in  $Y$  and the lower sets  $L_\alpha(x) := \{y \in Y : f(x, y) \leq \alpha\}$  for  $x$  in  $X$  are closed and convex. It is well known and easy to see that the minima and maxima in the theorem exist under our topological assumptions.

Sion proved his theorem with the aid of Helly's theorem and the "KKM-Theorem" of Knaster, Kuratowski, and Mazurkiewicz. The special case where  $X$  and  $Y$  are finite-dimensional simplexes and  $f$  is a continuous function was proved by von Neumann [4] by reducing the problem to the 1-dimensional case.

We present a proof that is close in spirit to von Neumann's original proof. It uses only the 1-dimensional KKM-theorem (i.e., every interval in  $\mathbb{R}$  is connected) and the 1-dimensional Helly theorem (i.e., any family of pairwise intersecting compact intervals in  $\mathbb{R}$  has nonvoid intersection).

The following lemma was inspired by Groemer's paper [1] and by [2].

**Lemma.** *In the situation of the theorem, let  $K_1$  and  $K_2$  be nonvoid convex and compact subsets of  $X$  with  $X = K_1 \cup K_2$ . Then*

$$\xi := \max_{i \in \{1,2\}} \min_{y \in Y} \max_{x \in K_i} f(x, y) = \beta := \min_{y \in Y} \max_{x \in X} f(x, y).$$

*Proof.* Choose  $z_i$  in  $Y$  with  $\max_{x \in K_i} f(x, z_i) \leq \xi$  for  $i = 1, 2$ . Then we have  $\xi \geq \max_{x \in X} \min_{y \in I} f(x, y)$ , where  $I = [z_1, z_2]$  is the convex hull of  $\{z_1, z_2\}$ . Suppose that  $\min_{y \in I} \max_{x \in X} f(x, y) > \delta > \xi$  for some real  $\delta$ . Then we get  $\bigcap_{x \in X} L_\delta(x) \cap I = \emptyset$  and  $\bigcap_{y \in I} U_\delta(y) = \emptyset$ . By the 1-dimensional Helly theorem, there exist  $x_1$  and  $x_2$  in  $X$  with

$$L_\delta(x_1) \cap L_\delta(x_2) \cap I = \emptyset \quad (1)$$

and there exist  $y_1$  and  $y_2$  in  $I$  with

$$U_\delta(y_1) \cap U_\delta(y_2) \cap [x_1, x_2] = \emptyset,$$

which implies that

$$L_\delta(x_i) \cap \{y_1, y_2\} \neq \emptyset \quad (i = 1, 2). \quad (2)$$

Since the sets  $L_\delta(x_i)$  are closed and  $[y_1, y_2]$  is connected, relations (1) and (2) ensure the existence of  $y_0$  such that

$$y_0 \in [y_1, y_2] \setminus (L_\delta(x_1) \cup L_\delta(x_2)). \quad (3)$$

A similar argument shows that there exists  $x_0$  such that

$$x_0 \in [x_1, x_2] \setminus (U_\delta(y_1) \cup U_\delta(y_2)). \quad (4)$$

By (3) we have  $\{x_1, x_2\} \subset U_\eta(y_0)$  for some  $\eta > \delta$ . Since  $x_0 \in [x_1, x_2] \subset U_\eta(y_0)$  we infer that  $f(x_0, y_0) > \delta$ . Similarly, (4) implies that  $f(x_0, y_0) < \delta$ , a contradiction. Therefore we have  $\xi \geq \min_{y \in I} \max_{x \in X} f(x, y) \geq \beta (\geq \xi)$ . ■

*Proof of the theorem.* We fix a finite subset  $A$  of  $X$  and set

$$\beta = \min_{y \in Y} \max_{x \in [A]} f(x, y),$$

where  $[A]$  denotes the convex hull of  $A$ . Let  $\mathcal{K}$  denote the system of all nonvoid closed convex subsets  $K$  of  $[A]$  with  $\min_{y \in Y} \max_{x \in K} f(x, y) = \beta$ . Let  $\mathcal{C}$  be a chain in  $\mathcal{K}$ . Then the fact that  $U_\beta(y) \cap C \neq \emptyset$  for all  $C$  in  $\mathcal{C}$  and  $y$  in  $Y$  implies that  $U_\beta(y) \cap \bigcap \mathcal{C} \neq \emptyset$  for each  $y$  in  $Y$  (i.e.,  $\bigcap \mathcal{C} \in \mathcal{K}$ ). Therefore,  $\mathcal{K}$  is inductively ordered by inclusion.

By Zorn's Lemma,  $\mathcal{K}$  has a minimal element  $K_0$ . Suppose that  $K_0$  has proper closed convex subsets  $K_1$  and  $K_2$  with  $K_0 = K_1 \cup K_2$ . Then our lemma tells us that either  $K_1 \in \mathcal{K}$  or  $K_2 \in \mathcal{K}$ , contradicting the minimality of  $K_0$ . Hence,  $K_0$  must be a singleton  $\{x_0\}$ . This implies that

$$\alpha := \max_{x \in X} \min_{y \in Y} f(x, y) \geq \min_{y \in Y} f(x_0, y) = \min_{y \in Y} \max_{x \in [A]} f(x, y) \geq \min_{y \in Y} \max_{x \in A} f(x, y),$$

whence  $\bigcap_{x \in A} L_\alpha(x) \neq \emptyset$  for each finite subset  $A$  of  $X$ . Since  $Y$  is compact and all the sets  $L_\alpha(x)$  are closed, there exists a  $y_0$  in  $\bigcap_{x \in X} L_\alpha(x)$ . Accordingly,

$$\alpha \geq \max_{x \in X} f(x, y_0) \geq \min_{y \in Y} \max_{x \in X} f(x, y) (\geq \alpha). \quad \blacksquare$$

**Remark.** If one does not want to use Zorn's Lemma, one can also proceed in the foregoing proof as in [1] or [2]: if  $K$  in  $\mathcal{K}$  is not a singleton, then it can be split by some hyperplane into two convex compact subsets  $K = K_1 \cup L_1$ , and by our lemma we may assume that  $K_1$  belongs to  $\mathcal{K}$ . Continuing this process, one can arrange the construction of sets  $K_n$  in  $\mathcal{K}$  ( $n = 1, 2, 3, \dots$ ) such that their diameters decrease to 0, hence,  $\bigcap_{n \in \mathbb{N}} K_n$  is a singleton  $\{x_0\}$ . Now  $U_\beta(y) \cap K_n \neq \emptyset$  for each  $n$ , so  $x_0$  lies in  $U_\beta(y)$  for each  $y$  in  $Y$  (i.e.,  $\{x_0\}$  is a member of  $\mathcal{K}$ ).

#### REFERENCES

1. H. Groemer, On the min-max theorem for finite two-person zero-sum games, *Z. Wahrscheinlichkeitstheorie und Verw. Geb.* **9** (1967) 59–61.
2. J. Kindler, Über Spiele auf konvexen Mengen, in *Operations Res. Verfahren*, no. 26, Hain, Meisenheim/Glan, 1977, pp. 695–704.
3. M. Sion, On general minimax theorems, *Pac. J. Math.* **8** (1958) 171–176.
4. J. von Neumann, Zur Theorie der Gesellschaftsspiele, *Math. Ann.* **100** (1928) 295–320.

*Fachbereich Mathematik, Technische Universität Darmstadt, Schlossgartenstr. 7, D-64289 Darmstadt, Germany*  
*kindler@mathematik.tu-darmstadt.de*

---

## A Constructive Approach to Singular Value Decomposition and Symmetric Schur Factorization

---

**John Clifford, David James, Michael Lachance, and Joan Remski**

---

**1. INTRODUCTION.** Fundamental to understanding linear transformations in  $\mathbb{R}^n$  is the fact that the image of a sphere under a matrix mapping  $\mathbf{A}$  is an ellipsoid. This geometrical point is seldom emphasized in sophomore-level linear algebra courses because its demonstration depends on the “Singular Value Decomposition,” the proof of which, in most texts, requires the somewhat intimidating “Symmetric Schur Factorization.” So the opportunity to make a very appealing geometrical point is lost. We believe that this is both unfortunate and unnecessary.