Schistosomiasis

This article is about the disease. For the organism, see <u>Schistosoma</u>.

Schistosomiasis

Other names Bilharzia, snail fever, Katayama fever^{[1][2]}

11-year-old boy with abdominal fluid and portal hypertension due to

schistosomiasis (Agusan del Sur, Philippines)

Pronunciation /\sistasa'mai\text{\text{maides}}\sistassa'-to\text{\text{\text{c}}}\display-\sigma'-so\text{\text{\text{c}}}\display-\frac{[3][4]}{}

Medical specialty Infectious disease

Symptoms

Abdominal pain, diarrhea, bloody stool, blood in the

urine^[5]

Complications
Liver damage, kidney failure, infertility, bladder

cancer^[5]

Causes Schistosomes from freshwater snails^[5]

Finding eggs of the parasite in urine or stool,

Diagnostic method antibodies in

antibodies in blood [5]

Prevention Access to clean water [5]

Medication Praziquantel^[5]

Frequency $252 \text{ million } (2015)^{[6]}$ **Deaths** $4.400-200.000^{[7][8]}$

Schistosomiasis, also known as **snail fever** and **bilharzia**, ^[9] is a <u>disease</u> caused by <u>parasitic flatworms</u> called <u>schistosomes</u>. ^[5] The <u>urinary tract</u> or the <u>intestines</u> may be infected. ^[5]

Symptoms include abdominal pain, diarrhea, bloody stool, or blood in the urine. Those who have been infected for a long time (chronic infection) may experience liver damage, kidney failure, infertility, or bladder cancer. In children, it may cause poor growth and learning difficulty.

Cause

[change | change source]

The disease is caused by contact with <u>fresh water</u> containing the parasites (schistosomes). These parasites are from infected <u>freshwater snails</u>. The disease is very common among children in <u>developing countries</u>, as they are more likely to play in infected water. Other high-risk groups include farmers, fishermen, and people using unclean water. It is a <u>helminth</u>

infections. [10] Eggs of the parasite may be found in urine or stool. [5] Antibodies may also be found in the blood. [5]

Prevention and treatment

[change | change source]

To prevent the disease use clean water and reduce snails. [5] The medicine praziguantel may be given once a year to everybody in some places. [5] This will decrease the number of people infected. [5] Praziguantel is also the treatment recommended by the World Health Organization for those who are known to be infected. [5]

Epidemiology

[change | change source]

Schistosomiasis affected about 252 million people in 2015. [6] An estimated 4,400 to 200,000 people die from it each year. [7][8] The disease is mainly found in Africa, Asia, and South America. [5] About 700 million people, in more than 70 countries, live where the disease is common. [7][11] In tropical countries, schistosomiasis is a great economic problem. [12] Schistosomiasis is listed as a neglected tropical disease. [13]

References

[change | change source]

- 1. ↑ "Schistosomiasis (bilharzia)". NHS Choices. 17 December 2011. Archived from the original on 15 March 2014. Retrieved 15 March
- 2. ↑ "Schistosomiasis". Patient.info. 2 December 2013. Archived from the original on 26 June 2015. Retrieved 11 June 2014.
- 3. ↑ "schistosomiasis definition of schistosomiasis in English from the Oxford dictionary". OxfordDictionaries.com. Archived from the original on July 16, 2014. Retrieved 20 January 2016.
- 4. ↑ "Schistosomiasis". *Merriam-Webster Dictionary*.
 5. ↑ 5.00 5.01 5.02 5.03 5.04 5.05 5.06 5.07 5.08 5.09 5.10 5.11 5.12 5.13 5.14 5.15 5.16 5.17 5.18 5.19 5.20 5.21 <u>"Schistosomiasis Fact sheet N°115"</u>. World Health Organization. 3 February 2014. Archived from the original on 12 March 2014. Retrieved 15 March 2014.
- 6. \uparrow 6.0 6.1 Vos T, Allen C, Arora M, Barber RM, Bhutta ZA, Brown A, et al. (GBD 2015 Mortality and Causes of Death Collaborators) (October 2016). "Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015; a systematic analysis for the Global Burden of Disease Study 2015". Lanc

- *et.* **388** (10053): 1545–1602. doi:10.1016/S0140-6736(16)31678-6. PMC 5055577 . PMID 27733282.
- $^{7.}$ \uparrow $^{7.0}$ $^{7.1}$ $^{7.2}$ Thétiot-Laurent SA, Boissier J, Robert A, Meunier B (July 2013). "Schistosomiasis chemotherapy". *Angewandte Chemie*. **52** (31): 7936–56. doi:10.1002/anie.201208390. PMID 23813602.
- 8. ↑ 8.0 8.1 Vos T, Allen C, Arora M, Barber RM, Bhutta ZA, Brown A, et al. (GBD 2015 Mortality and Causes of Death Collaborators) (October 2016). "Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: a systematic analysis for the Global Burden of Disease Study 2015". Lancet. 388 (10053): 1459-1544. doi:10.1016/s0140-6736(16)31012-1. PMC 5388903 . PMID 27733281.
- 9. ↑ Colley DG, Bustinduy AL, Secor WE, King CH (June 2014). "Human schistosomiasis". *Lancet*. **383** (9936): 2253-64. doi:10.1016/s0140-6736(13)61949-2. PMC 4672382 . PMID 24698483.
- 10. ↑ "Chapter 3 Infectious Diseases Related To Travel". cdc.gov. 1 August 2013. Archived from the original on 2 April 2015. Retrieved 30 November 2014.
- 11. ↑ "Schistosomiasis A major public health problem". World Health Organization. Archived from the original on 5 April 2014. Retrieved 15 March 2014.
- 12. ↑ The Carter Center. "Schistosomiasis Control Program". Archived from the original on 20 July 2008. Retrieved 17 July 2008.
- 13. <u>↑ "Neglected Tropical Diseases"</u>. cdc.gov. 6 June 2011. <u>Archived</u> from the original on 4 December 2014. Retrieved 28 November 2014.

This <u>short article</u> about <u>biology</u> can be made longer. You can help Wikipedia by <u>adding to it</u>.

Retrieved from "https://simple.wikipedia.org/w/index.php? title=Schistosomiasis&oldid=9597060"