arXiv:2609.17087v1 [math.LO] 15 Sep 2026

The Boolean Prime Ideal Theorem and Vitali sets*

Jindřich Zapletal
University of Florida
zapletal@ufl.edu

September 16, 2026

Abstract

If ZFC is consistent then so is the theory ZF+DC+BPI+there is no Vitali set.

1 Introduction

The Boolean Prime Ideal theorem (BPI) is the statement that every Boolean algebra carries an ultrafilter. This is a prominent consequence of the Axiom of Choice (AC); it is equivalent, among other things, to the statement that every consistent first order theory has a consistent completion. It is not equivalent to the Axiom of Choice, as a classical theorem of Halpern and Lévy [2] showed. Later, Blass [1] showed that BPI in permutation models is connected with the Ramsey properties of structures used to generate the models.

Since the methods of [2] and [1] necessarily imply a violation of the Axiom of Dependent Choices (DC), it is natural to ask whether the conjunction of the BPI and DC is equivalent to AC. [9] found a model which satisfies BPI+DC, but the Axiom of Choice for collections of cardinality $\aleph_1$ fails. That claim remains beyond the ability of the present writer to check, but more recently [10] found such a model using a rather standard permutation technique. In all similar models in the current literature, e.g. [5], the reals are well-ordered. To rectify this shortcoming, I show in this paper that BPI+DC does not imply the existence of Vitali sets.

Theorem 1.1. *If ZFC is consistent then so is the theory ZF+DC+BPI+there is no Vitali set.*

The approach to the proof of Theorem 1.1 is quite different from the standard symmetric or permutation model constructions; it uses intersection models of [7] instead. This in fact simplifies the argument considerably and brings it closer to the heart of a traditional forcing practitioner. The following is a brief description of the model.

*2020 AMS subject classification 03E25, 22F05. Keywords: Boolean Prime Ideal Theorem, Axiom of Dependent Choices, intersection model

Definition 1.2. $\mathbb{E}_0$ is the equivalence relation on 2^ω connecting two binary sequences if they agree on all but finitely many positions.

As is customary in descriptive set theory, I will call the $\mathbb{E}_0$ relation the *Vitali equivalence relation*; a *Vitali set* is a subset of 2^ω intersecting each $\mathbb{E}_0$-class in exactly one element.

Definition 1.3. The $\mathbb{E}_0$-forcing is the poset of all Borel subsets of 2^ω modulo the σ-ideal generated by Borel partial $\mathbb{E}_0$-selectors, ordered by inclusion. Its generic object will be viewed as a function in 2^ω.

Definition 1.4. $\mathbb{P}$ is the countable support product of ω_2 many copies of the $\mathbb{E}_0$-forcing. Its generic object will be viewed as a function $c\colon \omega_2 \times \omega \to 2$.

Now, it is possible to describe the model for Theorem 1.1. Start with a model V of ZFC+CH. Let $c\colon \omega_2 \times \omega \to 2$ be a function $\mathbb{P}$-generic over V. For each $n \in \omega$, write $c_n\colon \omega_2 \times \omega \to 2$ for the function defined by $c_n(\alpha, m) = c(\alpha, n+m)$. Write $W = \bigcap_n V[c_n]$; then, the model W satisfies ZF+DC+BPI+there are no Vitali sets.

This paper uses the standard set theoretic notation of [6]. The work contained herein was partly supported by National Science Foundation grant DMS 2348371.

2 General theorems on intersection models

In this section, I gather the relevant general results on intersection models. The key contribution here is Proposition 2.4 which provides a testable criterion for BPI.

Definition 2.1. In a generic extension of V, a sequence $\langle c_n\colon n \in \omega\rangle$ of sets of ordinals is *coherent* if for every $m \in \omega$, $\langle c_n\colon n \geq m\rangle \in V[c_m]$ holds.

Definition 2.2. If $\langle c_n\colon n \in \omega\rangle$ is a coherent sequence, the *associated model* is the intersection $\bigcap_n V[c_n]$.

The intersection model is a model of ZF by [7, Theorem 4.2.9]. In fact, it satisfies DC by an argument in the supplemental electronic materials to [7]. Let me recreate the simple argument here for completeness.

Proposition 2.3. *Let $\langle c_n\colon n \in \omega\rangle$ be a coherent sequence of sets of ordinals generic over V. The associated intersection model satisfies DC.*

Proof. Let $\langle R, \leq\rangle$ be a partial order in the intersection model without minimal elements; I must produce an infinite descending sequence in the intersection model.

Let β be an ordinal such that for all $n \in \omega$, $c_n \subset \beta$ holds. Let α be a sufficiently large ordinal, so that the cumulative hierarchy rank of R is much smaller than α, and each c_n is generic over V using a poset of rank much smaller

than α. Let $\preceq$ be a well-ordering of V_α in the ground model. For each $n \in \omega$, let P_n be the $\leq$-first poset in V adding a subset of β such that c_n is generic over P_n. By recursion on $n \in \omega$ build a descending sequence $\langle r_n : n \in \omega \rangle$ of conditions in R in the following way: $r_0 \in R$ is arbitrary, and for each $n \in \omega$, $r_{n+1} = \tau_{n+1}/c_{n+1}$ where τ_{n+1} is the $\preceq$-least P_{n+1}-name such that $\tau_{n+1}/c_{n+1} < r_n$. Note that for each $m \in \omega$, the sequence $\langle r_n : n \geq m \rangle$ can be recovered by the same recursion process in $V[c_m]$ starting from r_m. It follows that the descending sequence $\langle r_n : n \in \omega \rangle$ is in the intersection model as required. □

There is a surprisingly simple tool for the verification of BPI in the intersection models.

Proposition 2.4. *Let $\langle c_n : n \in \omega \rangle$ be a coherent sequence of sets of ordinals generic over V. Suppose that there is a nonprincipal ultrafilter $U \in V$ on ω such that U generates an ultrafilter in $V[c_0]$. Then BPI holds in the intersection model.*

Note that the proposition says that if the largest generic extension is not far from the ground model, all intermediate intersection models satisfy BPI. This observation together with Proposition 2.3 provides an endless supply of heretofore unknown models of ZF+DC+BPI.

Proof. Let B be a Boolean algebra in the intersection model; I must find an ultrafilter on B in the intersection model. As in Proposition 2.3, I construct a sequence $\langle F_n : n \in \omega \rangle$ of ultrafilters on B such that for every $m \in \omega$, $\langle F_n : n \geq m \rangle \in V[c_n]$ holds.

This is easy to do. Let β be an ordinal such that for all $n \in \omega$, $c_n \subset \beta$ holds. Let α be a sufficiently large ordinal, so that the rank of B is much smaller than α, and each c_n is generic over V using a poset of rank much smaller than α. Let $\leq$ be a well-ordering of V_α in the ground model. For each $n \in \omega$, let P_n be the $\leq$-first poset in V adding a subset of β such that c_n is generic over P_n, and let τ_n be $\leq$-first P_n-name such that τ_n/c_n is an ultrafilter on B. Write $F_n = \tau_n/c_n$ and note that the sequence $\langle F_n : n \in \omega \rangle$ has the required coherence property.

Now, let $U \in V$ be a nonprincipal ultrafilter on ω which generates an ultrafilter in $V[c_0]$; by an abuse of notation, let U also denote the ultrafilter generated by U in $V[c_0]$ or $V[c_n]$ for $n \in \omega$. Let $F = \{b \in B : \{n \in \omega : b \in F_n\} \in U\}$. Clearly, this is an ultrafilter on B; I must prove that for every $m \in \omega$, $F \in V[c_m]$. This is immediate though. F is recovered in $V[c_m]$ as $F = \{b \in B : \{n \geq m : b \in F_n\} \in U\}$, using the fact that both $U \in V[c_n]$ and $\langle F_n : n \geq m \rangle \in V[c_n]$ hold. □

3 General theorems on $\mathbb{E}_0$ forcing

In this section, I recall the basic facts about the generic objects used to form the intersection model for Theorem 1.1. None of them are new.

The $\mathbb{E}_0$-forcing has appeared in the literature many times [11, Section 4.7.1]. In this section I describe its well-known combinatorial presentation and state an ultrafilter preservation theorem needed for the final proof.

Definition 3.1. The $\mathbb{E}_0$-tree is an ever-branching tree $T \subset 2^{<\omega}$ such that for every splitnode $t \in T$ and every $u \in T$ extending t, the node obtaining from u by flipping its $|t|$-th entry is in T as well.

Let I be the σ-ideal on 2^ω σ-generated by Borel partial $\mathbb{E}_0$-selectors. The (proof of the) Glimm-Effros dichotomy of [4] immediately implies that a Borel subset of 2^ω is I-positive if and only if it contains all branches of some $\mathbb{E}_0$-tree. Thus, the poset of Borel sets ordered by inclusion modulo I is in the forcing sense equivalent to the poset of $\mathbb{E}_0$-trees ordered by inclusion, and this is the combinatorial presentation of the poset I use below.

Fact 3.2. *Let U be a Ramsey ultrafilter on ω. Let $\mathbb{P}$ be a countable support product of any number of $\mathbb{E}_0$-forcings. In the $\mathbb{P}$-extension, U still generates an ultrafilter.*

Proof. This is a concatenation of several arguments well-known from pre-existing literature; I will only provide an outline of the proof. Suppose that $p \in \mathbb{P}$ is a condition and $p \Vdash \sigma \subset \omega$ is a set. I must show that there is a condition $r \leq p$ and a set $a \in U$ such that $r \Vdash \check a \subseteq \sigma$ or $r \Vdash \check a \cap \sigma = 0$.

Let I denote the index set for the product $\mathbb{P}$. A standard fusion argument provides a strengthening $q \leq p$ and a countable set $J \subset I$ such that below q, σ is in fact a name in the product $\mathbb{Q}$ of J-many copies of the $\mathbb{E}_0$-forcing. The poset $\mathbb{Q}$ adds no independent reals by [12, Example 1.10]; so the set $A = \{a \subset \omega \colon \exists r \leq q\, r \Vdash \check a \subseteq \sigma \text{ or } r \Vdash \check a \cap \sigma = 0\}$ is dense in the algebra $\mathcal{P}(\omega)$ modulo finite. Use the continuous reading of names to strengthen q if necessary to make sure that the set $A \subset \mathcal{P}(\omega)$ is analytic. Every Ramsey ultrafilter has a nonempty intersection with every analytic set dense in $\mathcal{P}(\omega)$ modulo finite by the arguments of [8]. This provides the desired condition r and the set $a \in U$. □

4 The conclusion of the argument

At this point, everything is ready to conclude the proof of Theorem 1.1. Suppose that the Continuum Hypothesis holds. Then there is a Ramsey ultrafilter on ω, pick one and call it U. Let $\mathbb{P}$ be a countable support product of ω_2-many copies of $\mathbb{E}_0$-forcing. Let $\dot c$ be the $\mathbb{P}$-name for the generic object, viewed as a map from $\omega_2 \times \omega$ to 2.

Starting from a model of ZFC+CH, let $c \colon \omega_2 \times \omega \to 2$ be a $\mathbb{P}$-generic function, and for every $n \in \omega$ write c_n for the function from $\omega_2 \times \omega$ to 2 defined by $c_n(\alpha, m) = c(\alpha, n + m)$. It is immediate that the sequence $\langle c_n \colon n \in \omega \rangle$ is coherent, so the intersection model satisfies ZF+DC by Proposition 2.3. The ultrafilter U generates an ultrafilter in $V[c]$ by Fact 3.2. Proposition 2.4 then shows that the intersection model satisfies BPI.

It remains to show that there is no Vitali set in the intersection model. In fact, observe that the sequence $\langle [c_\alpha]_{\mathbb{E}_0} : \alpha \in \omega_2 \rangle$ belongs to the intersection model; I will show that there is no selector on this sequence of sets. This is a consequence of a rather standard Δ-system argument in the ground model. To state it, I need to be able to move freely between the generic function and its fragments on the coherent sequence. The following definition is designed to facilitate such moves.

Definition 4.1. Suppose that $n \in \omega$ is a number.

1. For a function $c: \omega_2 \times \omega \to 2$, the function $c|n: \omega_2 \times \omega \to 2$ is defined by $(c|n)(\alpha, m) = c(\alpha, n+m)$.
2. For a condition $p \in \mathbb{P}$ and a function $v: \omega \to 2^n$ such that for every $\alpha \in \omega_2$, $v(\alpha) \in p(\alpha)$ holds, write $p|v$ for the condition defined by the demand $w \in (p|v)(\alpha)$ iff $v(\alpha)^\frown w \in p(\alpha)$.
3. For a condition $p \in \mathbb{P}$ and a function $v: \mathrm{supp}(p) \to \alpha$, write $p||v$ for the condition in $\mathbb{P}$ with the same domain defined by the demand $w \in (p||v)(\alpha)$ iff w is an initial segment of $v(\alpha)$ or there is $u \in p(\alpha)$ such that $w = v(\alpha)^u$.

Proposition 4.2. *Let $n \in \omega$ be a number. The poset $\mathbb{P}$ forces the function $\dot{c}|n$ to be $\mathbb{P}$-generic over the ground model.*

Proof. Consider the set D of all conditions p such that for every $\alpha \in \mathrm{supp}(p)$, the stem of $p(\alpha)$ is longer than n. It is obvious that the set $D \subset P$ is dense. Let $\pi: D \to \mathbb{P}$ be the function defined by $\pi(p) = p|v$ for some (any) function $v: \omega_2 \to 2^n$ such that for every $\alpha \in \mathrm{supp}(p)$, $v(\alpha) \in p(\alpha)$. It is not difficult to check that $\mathbb{P}$ forces the π-image of $D \cap \dot{G}$ to be a generic filter on $\mathbb{P}$ which yields the function $\dot{c}|n$. □

I need to show that there is no Vitali set in the intersection model. Suppose towards a contradiction that some condition $p \in \mathbb{P}$ forces $\sigma \subset 2^\omega$ to be a set in the intersection model with singleton intersection with all the sets $[\dot{c}_\alpha]_{\mathbb{E}_0}$ for $\alpha \in \omega_2$. For each ordinal $\alpha \in \omega_2$, there must be a condition $p_\alpha \leq p$ and a number $m_\alpha \in \omega$ such that p_α forces the unique element of σ in the $\mathbb{E}_0$-class of $\dot{c}_\alpha$ to agree with $\dot{c}_\alpha$ past $\check{m}_\alpha$. There must be a number $n_\alpha \in \omega$ such that the tree $p_\alpha(\alpha)$ contains two splitnodes at the level n_α which differ at some entry larger than m_α. By the CH assumption, there must be a cofinal set $A \subset \omega_2$ such that the numbers m_α for all $\alpha \in A$ coincide and are equal to some m, the numbers n_α for all $\alpha \in A$ coincide and are equal to some n, the sets $\mathrm{dom}(p_\alpha)$ for $\alpha \in A$ form a Δ-system with heart $h \subset \omega$, and the conditions $p_\alpha \restriction h$ all coincide and are equal to some $q \in \mathbb{P}$.

Let $r \leq q$ be a condition and τ be a $\mathbb{P}$-name such that $r \Vdash \sigma/\dot{c} = \tau/\dot{c}|\check{n}$. Choose an ordinal $\alpha \in A$ so that $\mathrm{dom}(p_\alpha) \cap \mathrm{dom}(r) = h$; so, p_α is compatible with r. Let r_α be the specific common lower bound of them defined by $\mathrm{supp}(r_\alpha) = \mathrm{supp}(r) \cup \mathrm{supp}(p_\alpha)$, for each $\beta \in \mathrm{supp}(r)$ $r_\alpha(\beta) = r(\beta)$, and for each $\beta \in \mathrm{supp}(p_\alpha) \setminus h$ $r_\alpha(\beta) = p_\alpha(\beta)$. Choose functions $v_0, v_1: \omega_2 \to 2^n$ such that they

differ only on α-th entry, for each $\beta \neq \alpha$ $v_0(\beta) = v_1(\beta) \in r_\alpha(gb)$, and the nodes $v_0(\alpha), v_1(\alpha) \in p_\alpha(\alpha)$ disagree at some entry larger than m. Note that the conditions $r_\alpha|v_0$ and $r_\alpha|v_1$ coincide by the symmetry properties of the $\mathbb{E}_0$-tree $p_\alpha(\alpha)$.

Find a condition $s \leq r_\alpha|v_0$ and a string $w \in 2^n$ such that s forces the unique element of τ which is $\mathbb{E}_0$-equivalent to $(0^n)^\frown \dot{c}_\alpha$ to start with $\check{w}$. By the choice of v_0 and v_1, one of the strings $v_0(\alpha), v_1(\alpha) \in 2^n$ disagrees with w at an entry $k > m$; for definiteness, assume $v_0(\alpha)(k) = 0$ while $w(k) = 1$. Consider the condition $s||v_0 \in \mathbb{P}_0$. It is stronger than p_α, so it forces the unique element of σ in the $\mathbb{E}_0$-class of $\dot{c}_\alpha$ to have 0 at m-th entry. However, by the choice of s it also forces that unique element to start with w. This is a contradiction, showing that no Vitali sets can exist in the intersection model.

References

[1] Andreas Blass. Partitions and permutation groups. In *Model Theoretic Methods in Finite Combinatorics*, Contemporary Mathematics 558, pages 453–466. American Mathematical Society, Providence, 2011.

[2] James D. Halpern and Azriel Lévy. The Boolean prime ideal theorem does not imply the axiom of choice. In *Axiomatic set theory*, volume 13 of *Proceedings of Symposia in Pure Mathematics*, pages 84–134. 1971.

[3] J.D. Halpern and H. Lauchli. A partition theorem. *Transactions of American Mathematical Society*, 124:360–367, 1966.

[4] Leo Harrington, Alexander S. Kechris, and Alain Louveau. A Glimm-Effros dichotomy for Borel equivalence relations. *Journal of the American Mathematical Society*, 3:903–928, 1990.

[5] Peter Holy and Jonathan Schilhan. The ordering principle and higher dependent choice, 2025. https://arxiv.org/abs/2510.14821.

[6] Thomas Jech. *Set Theory*. Springer Verlag, New York, 2002.

[7] Paul Larson and Jindřich Zapletal. *Geometric set theory*. AMS Surveys and Monographs. American Mathematical Society, Providence, 2020.

[8] Adrian R.D. Mathias. Happy families. *Annals of Mathematical Logic*, 12:59–111, 1977.

[9] David Pincus. Adding dependent choice to the prime ideal theorem. Logic colloquium 76, Proc. Conf., Oxford 1976, Stud. Logic Found. Math., Vol. 87, 547-565, 1977.

[10] Brian Ransom. On BPI in symmetric extensions part 1, 2025. https://arxiv.org/abs/2511.21684.

[11] Jindřich Zapletal. *Forcing Idealized*. Cambridge Tracts in Mathematics 174. Cambridge University Press, Cambridge, 2008.

[12] Jindrich Zapletal. Subadditive families of hypergraphs. *Annals of Pure and Applied Logic*, 175, 2024. Paper No. 103322, 9 pp.