An Implementation of DVCM in Water Hammer Event

刘锦坤; 姜逸轩; 黑建聪

Tsinghua University

2024.5.21

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

Introduction of DVCM




Fig. 9. Definition sketch for discrete vapor cavity model (adapted from Tijsseling, 1993, Fig. 4.9).

Sketch of DVCM Model

The sketch of DVCM is shown in the figure above. The DVCM model is a 1D model that can simulate the water hammer event in the pipeline. The model allows the discrete cavitation bubbles to be generated and collapsed at the grid points.

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

The basic equations of MOC(method of characteristics) are farmiliar to us. The equations are shown below:

$$\begin{cases} H_{j}^{t} - H_{j-1}^{t-\Delta t} + \frac{a}{gA}((Q_{u})_{j}^{t} - Q_{j-1}^{t-\Delta t}) + \frac{f\Delta x}{2gDA^{2}}(Q_{u})_{j}^{t}|Q_{j-1}^{t-\Delta t}| = 0\\ H_{j}^{t} - H_{j+1}^{t-\Delta t} - \frac{a}{gA}(Q_{j}^{t} - (Q_{u})_{j+1}^{t-\Delta t}) - \frac{f\Delta x}{2gDA^{2}}Q_{j}^{t}|(Q_{u})_{j+1}^{t-\Delta t}| = 0 \end{cases}$$

$$(1)$$

◆□▶ ◆□▶ ◆臺▶ ◆臺▶ · 臺 · 釣९♂

$$\begin{cases}
H_{j}^{t} - H_{j-1}^{t-\Delta t} + \frac{a}{gA}((Q_{u})_{j}^{t} - Q_{j-1}^{t-\Delta t}) + \frac{f\Delta x}{2gDA^{2}}(Q_{u})_{j}^{t}|Q_{j-1}^{t-\Delta t}| = 0 \\
H_{j}^{t} - H_{j+1}^{t-\Delta t} - \frac{a}{gA}(Q_{j}^{t} - (Q_{u})_{j+1}^{t-\Delta t}) - \frac{f\Delta x}{2gDA^{2}}Q_{j}^{t}|(Q_{u})_{j+1}^{t-\Delta t}| = 0 \\
Q_{u} = Q
\end{cases}$$
(1)

H represents the head of the water, Q represents the downstream flow rate of the water, Q_u represents the upstream flow rate. The footnote represents the number of node, the headnode represents the time.

As we did in the previous assignment, we consider that $Q_u = Q$, and then solve the equations of the head $H_j^{t+\Delta t}$ and flow rate $Q_i^{t+\Delta t}$ these two unknowns.

But in the DVCM model, the cavitation bubbles are considered. Once if the head drops below the vapor pressure H^* , the cavitation happens and 2 equations are attached to the original equations:

$$\begin{cases} \dot{V} = -(Q_u) + Q \\ H = H^* \end{cases}$$

Since when $Q_u \neq Q$ and in which V represents the volumn of the cavitation bubble.

The discrete form of the equations would be more useful as shown in the following:

$$\begin{cases}
V_j^{t+\Delta t} = V_j^t + \psi(-(Q_u)_j^{t+\Delta t} + Q_j^{t+\Delta t})\Delta t + (1-\psi)(-(Q_u)_j^t + Q_j^t)\Delta t \\
H_j^{t+\Delta t} = H^*
\end{cases}$$
(2)

As we have learnt in mathmatical analysis, ψ is the weight factor generated by the mean value theorem of integrals.

Therefore, the unknowns are $(Q_u)_j^{t+\Delta t}$ and $Q_j^{t+\Delta t}$ in (1) and (2) instead of $H_j^{t+\Delta t}$ and $Q_j^{t+\Delta t}$ (equals to $(Q_u)_j^{t+\Delta t}$).

After solving the Q and (Q_u) , we can get the ΔV and then update the V. Once updated V is less than 0, the cavitation bubble collapses and the equations go back to the original form.

4□ > 4□ > 4 = > 4 = > = 90

Sumarry of Details of DVCM

In sumarry, when $H_i^t > H^*$ the normal equations are:

$$\begin{cases} H_{j}^{t} - H_{j-1}^{t-\Delta t} + \frac{a}{gA}((Q_{u})_{j}^{t} - Q_{j-1}^{t-\Delta t}) + \frac{f\Delta x}{2gDA^{2}}(Q_{u})_{j}^{t}|Q_{j-1}^{t-\Delta t}| = 0\\ H_{j}^{t} - H_{j+1}^{t-\Delta t} - \frac{a}{gA}(Q_{j}^{t} - (Q_{u})_{j+1}^{t-\Delta t}) - \frac{f\Delta x}{2gDA^{2}}Q_{j}^{t}|(Q_{u})_{j+1}^{t-\Delta t}| = 0\\ Q_{u} = Q \end{cases}$$

$$(1)$$

And once $H_i^t < H^*$ the equations are added:

$$\begin{cases} V_j^{t+\Delta t} = V_j^t + \psi(-(Q_u)_j^{t+\Delta t} + Q_j^{t+\Delta t})\Delta t + (1-\psi)(-(Q_u)_j^t + Q_j^t)\Delta t \\ H_j^{t+\Delta t} = H^* \end{cases}$$
(2)

And when $V_i^{t+\Delta t} < 0$, the equations go back to (1).

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

TALK IS CHEAP, SHOW ME THE CODE

Some key parts of the code are shown below:

Key parts of the code to implement DVCM

Introduction of DVCM


Details of DVCM

Implementation of DVCM

Results of DVCM


Future Work

Results of DVCM



Results of DVCM

Compared to Normal MOC



Results of MOC

Physics Features to Be Explained

- ▶ The cavitation bubbles are generated and collapsed in the DVCM model so in some periods of time ,the pressure is a line with value H^* .
- ▶ The water head in the DVCM model is lower than the normal MOC model after the first oscillation. This could be explained that the generation and the diminishment of the cavitation bubbles consume the energy of the water. (Seems Good News? Less Pressure, Less Damage? Not Sure Yet!)
- ▶ The water head in the DVCM model is more fluctuated than the normal MOC model due to the turbulence of the cavitation bubbles.

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

Future the Work

Congratulation: We have set the primary step on the moon!

- ▶ What do the experimenters say?
- Considering whether the DVCM can be developed by the conservation of mass and energy.
- ▶ What about other improved 1D models (DGCM discrete gas cavity model)?
- ▶ What about high dimensional models? Seems to be hard, limited desire to try. (QAQ)

Introduction of DVCM

Details of DVCM

Implementation of DVCM

Results of DVCM

Future Work

Acknowledgement

- ► Thanks to the teachers for the guidance.
- ▶ Thanks to the groupmates for their cooperation.
- ▶ Thanks to the audience for their attention.
- ► Thanks to Yuhan Zhang for her help.