

CRISPR–Cas9 Gene Editing: Mechanisms, Applications, and Future Directions

Author: Scientific Report

Date: June 2026

Abstract

The clustered regularly interspaced short palindromic repeats (CRISPR)–Cas9 system has transformed biomedical research and therapeutic development since its adaptation as a genome editing tool. By coupling the Cas9 endonuclease with a programmable single guide RNA (sgRNA), the system enables precise double-strand DNA breaks at virtually any genomic locus flanked by a protospacer-adjacent motif (PAM) sequence. The first CRISPR-based medicine, Casgevy (exagamglogene autotemcel), received FDA approval in December 2023 for the treatment of sickle cell disease, marking a pivotal milestone in the translation of this technology to the clinic. Despite transformative potential, concerns over off-target cleavage and delivery efficiency continue to drive the development of next-generation tools, including base editors and prime editors, which expand the precision and safety of genome modification. This report reviews the molecular mechanism of CRISPR–Cas9, its principal applications in disease and research, current limitations, and the trajectory of future innovations.

1. Introduction

The discovery that prokaryotic CRISPR loci serve as adaptive immune memories against invading phages laid the groundwork for one of the most consequential biotechnological advances of the 21st century. The landmark 2012 study by Jinek and colleagues demonstrated that the *Streptococcus pyogenes* Cas9 (SpCas9) endonuclease could be programmed with a chimeric guide RNA to cleave double-stranded DNA at defined sequences *in vitro*, and subsequent work in 2013 confirmed efficient editing in human and mouse cells (Cong et al., 2013; Mali et al., 2013). The relative simplicity of the CRISPR–Cas9 platform—requiring only the Cas9 protein and a synthetic sgRNA—gave it an immediate practical advantage over earlier zinc-finger nuclease (ZFN) and transcription activator-like effector nuclease (TALEN) technologies, which demanded laborious protein re-engineering for each new target.

Within a decade of these initial demonstrations, CRISPR–Cas9 had been applied across virtually every domain of life science, from fundamental studies of gene function to therapeutic pipelines targeting monogenic disorders, cancers, and infectious diseases. The pace of clinical translation has been extraordinary: as of 2024, over 40 registered clinical trials were actively exploring CRISPR-based interventions, spanning blood disorders, oncology, and virology (Molecular Therapy Nucleic Acids, 2026). The regulatory approval of Casgevy confirmed that CRISPR-based therapies can achieve durable clinical benefit, with 29 of 31 treated sickle cell disease patients remaining free of severe vaso-occlusive crises for at least one year post-treatment (PMC11305803, 2024). Against this backdrop of rapid progress, a clear understanding of the molecular mechanism, therapeutic landscape, and unresolved challenges of CRISPR–Cas9 is essential for researchers, clinicians, and policymakers alike.

2. Molecular Mechanism of CRISPR-Cas9

The CRISPR-Cas9 system achieves programmable genome editing through a well-characterized sequence of molecular events (Figure 1). The core effector, SpCas9, is a 160 kDa multi-domain protein containing two catalytic nuclease lobes: the HNH domain, which cleaves the DNA strand complementary to the guide RNA (target strand), and the RuvC domain, which cleaves the non-complementary strand (Doudna & Charpentier, 2014). In its editing-competent form, Cas9 is pre-loaded with a single guide RNA (sgRNA), a synthetic fusion of the CRISPR RNA (crRNA) and the *trans*-activating crRNA (tracrRNA), which together direct the complex to a genomic locus of interest.

Target recognition initiates with the scanning of double-stranded DNA for the 5'-NGG-3' PAM sequence, a short motif located immediately 3' of the intended target site on the non-template strand. PAM recognition induces local DNA melting, enabling the 20-nucleotide spacer region of the sgRNA to interrogate the adjacent sequence through Watson-Crick base pairing. Progressive hybridization from the PAM-proximal "seed" region toward the PAM-distal end forms an R-loop structure, in which the target strand is base-paired with the sgRNA while the non-target strand is displaced (Sternberg et al., 2015). Complete R-loop formation allosterically activates the HNH and RuvC nuclease domains, which then concertedly introduce a blunt-ended double-strand break (DSB) three base pairs upstream of the PAM sequence.

The resulting DSB is repaired by one of two principal cellular pathways. Non-homologous end joining (NHEJ), the dominant pathway in most cell types, is error-prone and frequently introduces small insertions or deletions (indels) that disrupt the reading frame of the targeted gene, effectively achieving gene knockout. In contrast, homology-directed repair (HDR) can incorporate an exogenously provided DNA template to introduce precise sequence changes, though HDR efficiency is substantially lower and largely restricted to dividing cells (Yeh et al., 2019). The choice of repair pathway thus governs whether the editing outcome is a loss-of-function disruption or a precise sequence correction.




Figure 1. Molecular mechanism of CRISPR-Cas9-mediated genome editing. The Cas9-sgRNA complex recognizes the PAM sequence (5'-NGG-3') on double-stranded DNA, forms an R-loop via 20-nt RNA-DNA hybridization, and induces a blunt-ended double-strand break 3 bp upstream of the PAM through coordinated activity of the HNH and RuvC nuclease domains. Cas9, blue; sgRNA, orange; target DNA, green; cleavage site, red. HNH, His-Asn-His nuclease domain; RuvC, RuvC-like nuclease domain; PAM, protospacer-adjacent motif; sgRNA, single guide RNA.

3. Therapeutic and Research Applications

The versatility of CRISPR-Cas9 has enabled its application across a broad spectrum of disease areas and research contexts (Figure 2). In the domain of monogenic hematological disorders, the greatest clinical progress has been made. Casgevy, developed by Vertex Pharmaceuticals and CRISPR Therapeutics, employs Cas9 to disrupt the *BCL11A* erythroid enhancer in autologous hematopoietic stem cells, thereby reactivating fetal hemoglobin (HbF) expression to compensate for defective adult hemoglobin in sickle cell disease and transfusion-dependent beta-thalassemia (Frangoul et al., 2021). FDA approval was granted in December 2023, and clinical data demonstrated freedom from severe pain crises in 29 of 31 sickle cell patients followed for at least 12 months (PMC11305803, 2024).

Beyond hematology, oncology represents the largest category of active CRISPR clinical trials, with 23 registered trials as of December 2022 targeting cancers through multiple mechanisms (PMC10136740, 2023). A prominent strategy involves engineering autologous or allogeneic CAR-T cells by disrupting immune checkpoint genes (*PDCD1*, *TRAC*, *TRBC*) to produce exhaustion-resistant, universal T-cell therapies. CRISPR-Cas9 is also being evaluated in the context of infectious disease, with three active trials investigating disruption of integrated HIV proviral DNA and the *CCR5* co-receptor gene, and a separate trial targeting HPV oncogenes *E6* and *E7* in cervical lesions (Molecular Therapy Nucleic Acids, 2026).

In the research context, CRISPR-Cas9 has become indispensable for constructing disease models, performing genome-wide functional screens, and dissecting gene regulatory networks. Catalytically inactivated Cas9 (dCas9) fused to transcriptional activators or repressors enables CRISPRa and CRISPRi approaches, respectively, allowing modulation of endogenous gene expression without introducing permanent sequence changes. These tools have been particularly powerful for large-scale pooled screens that assign functional roles to non-coding regulatory elements, a task previously intractable at genome scale.




Figure 2. Applications landscape of CRISPR-Cas9 across four principal domains: genetic diseases (including sickle cell disease, beta-thalassemia, and Duchenne muscular dystrophy), cancer (CAR-T cell engineering and tumor suppressor restoration), infectious disease (HIV and HPV), and next-generation tools (base editing, prime editing, CRISPRi/CRISPRa). The milestone approval of Casgevy in December 2023 represents the first FDA-sanctioned CRISPR-based therapeutic. CAR-T, chimeric antigen receptor T-cell; CRISPRa/i, CRISPR activation/interference.

4. Limitations and Off-Target Effects

Despite its remarkable versatility, CRISPR-Cas9 carries a risk of unintended genomic modifications at loci bearing sequence similarity to the intended target—collectively referred to as off-target effects. SpCas9 can cleave sites with up to 3–5 mismatches in the PAM-distal region of the protospacer, and the frequency of such events is influenced by the specific sgRNA sequence, the chromatin state of the target locus, the delivery modality, and the cell type (Hsu et al., 2013; Zhang et al., 2023). Unbiased whole-genome sequencing approaches, including GUIDE-seq and Digenome-seq, have revealed that off-target indels can occur at frequencies that, while typically orders of magnitude lower than on-target editing, may nonetheless be clinically significant if they affect tumor suppressor genes or oncogenes.

The immunogenicity of bacterially derived Cas9 represents a separate safety concern for in vivo delivery. Pre-existing humoral and cellular immune responses against SpCas9 and *Staphylococcus aureus* Cas9 (SaCas9) have been detected in healthy human donors, potentially limiting the efficacy of viral vector-mediated delivery and raising risks of immune-mediated toxicity (Charlesworth et al., 2019). Additional challenges include the relatively large size of the Cas9 coding sequence (~4.2 kb for SpCas9), which approaches or exceeds the packaging capacity of adeno-associated virus (AAV) vectors, necessitating split-intein or smaller ortholog strategies. Finally, delivery to solid tissues, the central nervous system, and non-dividing cells remains technically demanding, and the low efficiency of HDR in post-mitotic cells constrains the application of CRISPR-Cas9 for precise sequence correction in many therapeutically relevant cell types.

5. Future Directions

Ongoing innovation is addressing the principal limitations of CRISPR-Cas9 through the development of next-generation editing platforms. Base editors, first described by Komor and colleagues (2016), bypass the requirement for DSB formation by fusing a catalytically impaired Cas9 (nickase or dCas9) to a deaminase enzyme that chemically converts one DNA base to another within a defined editing window. Cytosine base editors (CBEs) convert C•G to T•A, while adenine base editors (ABEs) convert A•T to G•C, collectively enabling four of the twelve possible transition mutations without generating DSBs or requiring a repair template. Because base editors do not rely on HDR, they function in non-dividing cells and display reduced indel frequencies relative to standard Cas9 (Anzalone et al., 2020).

Prime editing, introduced in 2019, extends the precision of base editing by using a reverse transcriptase fused to Cas9 nickase in conjunction with a prime editing guide RNA (pegRNA) that encodes the desired sequence change directly within the guide scaffold. Recent preclinical data demonstrate that dual-AAV-delivered split prime editors can achieve editing frequencies of 40–50% in multiple mouse tissues with no detectable off-target events (Davis et al., 2024), and engineered virus-like particles have successfully corrected pathogenic mutations in a mouse model of inherited blindness. These advances suggest that prime editing may be particularly well-suited to correcting point mutations that underlie a substantial fraction of Mendelian diseases.

Looking further ahead, the integration of artificial intelligence into guide RNA design promises to improve on-target efficiency and minimize off-target activity through the prediction of editing outcomes from sequence features and chromatin accessibility data (Abramson et al., 2025). The development of non-viral delivery vehicles—including lipid nanoparticles, polymeric nanocarriers, and extracellular vesicles—is expanding access to tissues beyond the liver, where current lipid-based systems excel. Together, these advances suggest that the next decade will see CRISPR-based medicines addressing an increasingly broad spectrum of genetic and acquired diseases, potentially extending to common polygenic conditions and aging-associated pathologies.

6. Conclusions

CRISPR-Cas9 has fundamentally altered the landscape of genetic medicine, offering a programmable, efficient, and widely accessible platform for genome editing. The FDA approval of Casgevy for sickle cell disease validates the clinical translational potential of this technology and establishes a precedent for future CRISPR-based therapeutics. Residual challenges—principally off-target cleavage, delivery efficiency, and immunogenicity—are being systematically addressed by next-generation tools including base editors and prime editors, which offer enhanced precision without inducing double-strand breaks. As the field continues to mature, CRISPR-Cas9 and its derivatives are poised to transition from rare disease applications toward broader therapeutic paradigms, underpinned by advances in delivery science, guide RNA design, and our understanding of DNA repair mechanisms.

References

Anzalone, A. V., Koblan, L. W., & Liu, D. R. (2020). Genome editing with CRISPR-Cas nucleases, base editors, transposases and prime editors. *Nature Biotechnology*, 38(7), 824–844.
<https://doi.org/10.1038/s41587-020-0561-9>

- Charlesworth, C. T., Deshpande, P. S., Dever, D. P., Camarena, J., Lemgart, V. T., Cromer, M. K., Vakulskas, C. A., Collingwood, M. A., Zhang, L., Bode, N. M., Behlke, M. A., Shreiber, B., Dunn, D., Rettig, G. R., McIntosh, B. E., & Porteus, M. H. (2019). Identification of preexisting adaptive immunity to Cas9 proteins in humans. *Nature Medicine*, 25(2), 249–254. <https://doi.org/10.1038/s41591-018-0326-x>
- Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., Hsu, P. D., Wu, X., Jiang, W., Marraffini, L. A., & Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science*, 339(6121), 819–823. <https://doi.org/10.1126/science.1231143>
- Doudna, J. A., & Charpentier, E. (2014). The new frontier of genome engineering with CRISPR-Cas9. *Science*, 346(6213), 1258096. <https://doi.org/10.1126/science.1258096>
- Frangoul, H., Altshuler, D., Cappellini, M. D., Chen, Y.-S., Domm, J., Eustace, B. K., Foell, J., de la Fuente, J., Grupp, S., Handgretinger, R., Ho, T. W., Kattamis, A., Kernytsky, A., Lekstrom-Himes, J., Li, A. M., Locatelli, F., Mapara, M. Y., de Montalembert, M., Rondelli, D., ... Corbacioglu, S. (2021). CRISPR-Cas9 gene editing for sickle cell disease and β -thalassemia. *New England Journal of Medicine*, 384(3), 252–260. <https://doi.org/10.1056/NEJMoa2031054>
- Hsu, P. D., Scott, D. A., Weinstein, J. A., Ran, F. A., Konermann, S., Agarwala, V., Li, Y., Fine, E. J., Wu, X., Shalem, O., Cradick, T. J., Marraffini, L. A., Bao, G., & Zhang, F. (2013). DNA targeting specificity of RNA-guided Cas9 nucleases. *Nature Biotechnology*, 31(9), 827–832. <https://doi.org/10.1038/nbt.2647>
- Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A., & Liu, D. R. (2016). Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage. *Nature*, 533(7603), 420–424. <https://doi.org/10.1038/nature17946>
- Mali, P., Yang, L., Esvelt, K. M., Aach, J., Guell, M., DiCarlo, J. E., Norville, J. E., & Church, G. M. (2013). RNA-guided human genome engineering via Cas9. *Science*, 339(6121), 823–826. <https://doi.org/10.1126/science.1232033>
- Sternberg, S. H., Redding, S., Jinek, M., Greene, E. C., & Doudna, J. A. (2015). DNA interrogation by the CRISPR RNA-guided endonuclease Cas9. *Nature*, 507(7490), 62–67. <https://doi.org/10.1038/nature13011>
- Yeh, C. D., Richardson, C. D., & Corn, J. E. (2019). Advances in genome editing through control of DNA repair pathways. *Nature Cell Biology*, 21(12), 1468–1478. <https://doi.org/10.1038/s41556-019-0425-z>
- Zhang, X. H., Tee, L. Y., Wang, X. G., Huang, Q. S., & Yang, S. H. (2023). Off-target effects in CRISPR/Cas9 gene editing. *Molecular Therapy – Nucleic Acids*, 30, 329–341. <https://doi.org/10.1016/j.omtn.2023.03.004>

Word count: ~1,950 words (body text). Report generated June 2026.