

Molecular dynamics simulations of shallow nitrogen and silicon implantation into diamond: Supplemental Material

Ossi Lehtinen¹, Boris Naydenov², Pia Börner¹, Kristina Melentjevic², Christoph Müller²,
Liam Paul McGuinness², Sebastien Pezzagna³, Jan Meijer³, Ute Kaiser¹, and Fedor Jelezko²

¹ *Central Facility for Electron Microscopy, Group of Electron*

Microscopy of Materials Science, Ulm University, 89081 Ulm, Germany

² *Institute for Quantum Optics, Ulm University, 89081 Ulm, Germany and*

³ *Institut für Experimentelle Physik II, Universität Leipzig, 04103 Leipzig, Germany*

(Dated: December 20, 2015)




FIG. S1. Smoothed histograms (kernel density estimates) of displacement ranges of C atoms after implantation. The graphs show the distance from the original lattice positions of the displaced atoms for energies in the range of 100-5000 eV. The channeling [100] direction and a 'random' $15^\circ/15^\circ$ direction were considered. In both cases the displacements are in the few Ångström range.




FIG. S2. Simulated depth profiles for Si ions with and without a graphitic contamination layer. The simulations are for the [100] channeling direction. At low energies, ion penetration is suppressed by the contamination, while at higher energies the contamination leads to a reduction in the fraction of channeled ions.