

INDIAN INSTITUTE OF TECHNOLOGY, BOMBAY
DEPARTMENT OF CHEMISTRY



MANUAL

CH-117 L

CH117L: Activities

Sr. No.	Date	Activity
1	13 th March 2023- 6 th May 2023	Regular lab experiments (Expt.1- Expt.7)
2	8 th May 2023- 12 th May 2023	Compensation Lab
3	18 th May 2023- 30 th May 2023	CH117L Practical Examination
4	7 th June 2023	CH 117L Quiz
5	12 th June 2023– 19 th June 2023	End-sem examination
6	21 st June 2023	Crib session
7	25 th June 2023	Last Date for Uploading Grades

For any queries contact below:

Email Id: sushantc@chem.iitb.ac.in

(please mention “CH117L” in the subject, while sending emails).

Lab intercom number: 4183

General rules and regulations to be observed in Chemistry Ist Year U.G. laboratory.

1. **Full sleeves white lab coat** (preferably cotton), **safety goggles**, and **shoes** are mandatory while working in Chemistry Laboratory.
2. Only formal dresses (full) covering from neck to toe are allowed in the Chemistry laboratory. Students with shorts, three-fourths etc., will not be allowed.
3. **Punctuality:** Punctuality & discipline is a must in the Chemistry laboratory.
4. You should reach 5 mins before the scheduled time.
5. Penalties for late coming are as follows:
 - a) 5 minutes late, 1-mark penalty
 - b) 10 minutes late, 2-mark penalty
 - c) 15 minutes late, 5-mark penalty
 - d) 30 minutes late, experiment not allowed.
6. **Chemicals & apparatus:** Safety should be practiced while performing the experiments. You may come across with some new instruments, chemicals, acids, etc..., please handle them carefully and contact the instructor for any special precautions. Handle the glass wares gently.
7. **Lab report:** You are supposed to submit the lab report in a flat cardboard file, on a regular basis. You can purchase this file from the IITB consumer society shop located at the basement of the Main bldg. Also, buy a bunch of loose sheets (for write-up) and 5-6 graph papers along with the file. The file will remain in the laboratory until all experiments are completed.
8. Do not write anything in the file before coming to the lab.
9. TA will explain to you, what you should write in the journal [Objective, theory, principle, calculation, result, etc.] Please do not write procedures in the journal (file).
10. **Viva:** A viva session for each experiment will be conducted by the respective TA during laboratory hours. You should read and prepare for the experiment before coming to the laboratory to have a proper understanding of that experiment.
11. **Crib session:** In case of discrepancy a student can clarify the marks from the TA before the next lab i.e. within 7 days only.
12. **Compensation:**
 - a. Only one compensation lab will be conducted towards the end of the semester i.e. after all experiments are completed. If any student fails to attend the lab due to medical reasons and has a pink slip from IITB hospital, he/she should inform the staff in charge about the same in the next scheduled lab and apply (for compensatory lab) in hard copy along with all relevant documents.
 - b. Compensation will not be provided for any personal reasons such as family functions, brother's/sister's marriage or festival celebrations etc.
 - c. Compensation for more than one experiment (if, on medical grounds) will require approval from the course In-Charge.
 - d. Apart from medical reasons, the compensatory lab can be offered to a student who participates in an Academic activity (from IIT Bombay) that is scheduled on the day of his/her regular lab. The student should provide a Dean's recommendation letter and get prior permission from the Instructor in-charge.
13. Change of practical slot is not permitted.
14. **Mobile phones:** Mobile phones should be switched OFF and kept aside in the bag while working in the laboratory.
15. For calculations, please use calculators instead of mobile phones.
16. **Bags:** You can keep your bags or belongings in the drawers of the laboratory table.

17. **Feedback:** Feedback will be collected from you at the time of your last experiment for the improvisation of the course structure.
18. Please carry this Lab manual every time you come to the lab.

Marks distribution

1. The Lab course contains seven experiments.
2. Semester marks are out of 100.
3. 70 marks for regular experiment and 30 marks for final exam + written quiz.
4. 70 marks are further distributed in seven experiments with 10 marks for every experiment.
5. These 10 marks are split into further 3 parts as below:
 - a) **Performance** carries 5 marks, this includes your performance, punctuality, conduct, discipline, etc.
 - b) **Viva** carries 2 marks. The TA will ask questions based on the experiment (not confined to the manual), marks will be allotted based on your answers.
 - c) **Journal writing** (3 marks) will be explained by the respective TA at the time of the experiment.
6. Remaining 30 marks will be for the practical examination (20Marks) + quiz (10 marks).

CH 117L Exam: Instructions for students

Dear Students,

You are requested to carefully read the following instructions and understand the modalities

The practical examination is for **one and a half hours only**. You are requested to come 10 mins before the commencement of the examination. The timetable is available on Moodle.

- 1) Students should wear Lab-coat, shoes, and safety glasses for the entire duration.
- 2) You should bring only a pen, pencil, eraser, scale, calculator, and ID card for the examination. You should produce your ID card on demand, by the invigilator.
- 3) Mobile phones and smartwatches are not allowed during the examination.
- 4) **If we find that the student is having a mobile/smartwatch with him/her during the exam an FR grade will be awarded (even if not using it).**
- 5) You should not talk with your friend or neighbor during the examination till the time you are inside the laboratory.
- 6) **If you are found talking or communicating with other students by any means, it will be treated as academic misconduct and FR Grade will be awarded.**
- 7) If you are facing any difficulty or if you wish to talk about something, then you should contact the **TA/Staff only**. He/she will resolve the problem.
- 8) A question slip is attached to the answer sheet, do not detach the question slip from the answer sheet.
- 9) A number is printed on the question slip, kindly ensure that the number on the question slip and the given solutions are the same. You can copy the same number to your answer sheet as well.
- 10) The readings which you observe during the experiment should be written in the answer sheet and initialized by the TA assigned to your table.

- 11) If you have two sets of experiments then you should take at least one signature in each set.
- 12) Extra solutions/ apparatus will not be provided during the examination.
- 13) For experiments like redox titration, Zinc powder & water baths are provided on the side-table platforms.
- 14) You should write observation, observation table, calculation, and results only in the given answer sheet. Please do not write any theory, principle, or procedure for the experiment.
- 15) Viva session will not be there for the examination.
- 16) Examination marks are based on performance/ conduct during the exam (5 Marks), Readings (10 Marks), and Calculations/ Results (5 Marks).
- 17) In case during the experiment you forget and are not able to perform the experiment, the **Staff** will provide the hint for the experiment by deducting 5 marks. If you need more help then more marks will be deducted.
- 18) No help will be provided in the calculation part.
- 19) Graph paper if required will be provided by the TA.
- 20) Bio-breaks are not allowed.
- 21) You should ensure that your workplace and all apparatus are cleaned before you leave the lab.

Instructions for Quiz

- 1) The quiz will be conducted on a Question-cum-Answer sheet.
- 2) All are requested to bring their Id cards and produce them on demand by the invigilator.
- 3) Mobile phones and smartwatches are not allowed in examination hall.
- 4) **If we find that the student is having a mobile/smartwatch with him/her during the exam an FR grade will be awarded (even if not using it).**
- 5) You should not talk with your friend or neighbor during the examination till the time you are inside the laboratory. **If you are found talking or communicating with other students by any means, it will be treated as academic misconduct and FR Grade will be awarded.**
- 6) If you are facing any difficulty or if you wish to talk about something, then you should contact the **TA/Staff only**. He/she will resolve the problem.
- 7) You can carry a water bottle for drinking purposes in the examination hall.
- 8) You can bring a pen, pencil, eraser, scale and calculator along with you for writing the exam.
- 9) Bio-breaks are not allowed.
- 10) The seating plan for the exam will be shared with you later.

All the best!

Instructors CH-117L

[Please refer to the last page for a detailed schedule of the CH117L lab course.]

CH-117L 2022-2023: Spring semester

Details of Faculty members assigned for the respective slots of CH117L are as below:

Day/Div	Time(hrs)	Name of Faculty	Contact No.	Email ID
MON (P ₂₁)	0930-1230	Prof. Sanjog Nagarkar	7190	nagarkarss@iitb.ac.in
MON (P ₁₅)	1400-1655	Prof. Gopal Rajaraman	7183	rajaraman@iitb.ac.in
TUE (P ₁₉)	0830-1130	Prof. Ramaswamy Murugavel	7163	muks@iitb.ac.in
TUE (P ₁₃)	1400-1655	Prof. Rodney Fernandes	7174	rfernand@iitb.ac.in
THURS (P ₂₂)	0930-1230	Prof. Arindam Chowdhury	7154	achowdhury@iitb.ac.in
THURS (P ₁₆)	1400-1655	Prof. Sandip Kar	7193	sandipkar@iitb.ac.in
Fri (P ₂₀)	0930-1230	Prof. Anindya Datta	7149	adutta@iitb.ac.in
FRI (P ₁₄)	1400-1655	Prof. Sambasivarao Kotha	7160	srk@iitb.ac.in

(Prof. Sanjog Nagarkar)

Course In-charge

CONTENTS

Experiment No.	Title of the Experiment	Page No.
1	Electrochemical Cell	8
2	Removal efficiency for a Pharma Pollutant in water using MAC	13
3	Estimation of Iron	18
4	Oscillatory Chemical Reactions	20
5	Chemical kinetics	25
6	Complexometric Titration	30
7	Thin Layer Chromatography	33

1. Electrochemical Cells

Objectives:

- (A) To measure the standard electrode potential of $\text{Zn}^{2+} / \text{Zn}$ couple.
- (B) To determine the concentration of Fe^{2+} by potentiometric titration.

Introduction:

Electrochemical techniques can be broadly classified as equilibrium or dynamic techniques. In the present exercise we shall be using equilibrium techniques. These techniques are based on the Nernst equation:

$$E = E^\circ + 2.303 (RT/nF) \log a_{\text{ox}}/a_{\text{red}}$$


Which defines the electrochemical system at equilibrium. E is the reversible electrode potential measured against a suitable reference electrode. E° is the standard electrode potential measured against the same reference electrode is the number of electrons transferred in the red-ox processes; a_{ox} and a_{red} are the activities of the oxidized and reduced forms respectively. Thus, by measuring the electrode potential (against a suitable reference) one can determine E° .

In part (I) of this exercise, we will determine the standard electrode potential E° for red-ox couple Zn^{2+}/Zn




In part (II) we will make use of the Nernst equation to determine the concentration of Fe^{2+} in a solution of ferrous ammonium sulfate. The inert electrode or indicator electrode (Pt in this case) is used to sense the red-ox potential of the solution as it is being titrated against an oxidizing agent ($0.5\text{N K}_2\text{Cr}_2\text{O}_7$ in this experiment).


Procedure:

- (A) To determine the standard reduction potential of the reaction



Take sufficient amount of 0.1 M zinc sulphate solution in a beaker to partially immerse the zinc foil in this solution. This forms the Zinc half-cell. Connect this half-cell by means of a salt bridge (as shown in figure) with the saturated calomel half-cell to form the following galvanic cell:





Calculations:

(A) Determination of $E^{\circ}\text{Zn}^{++}/\text{Zn}$:

Cell set up: $(-) \text{Zn}/\text{Zn}^{++} (0.1 \text{ M}) \parallel \text{KCl (satd.)} \mid \text{Hg}_2\text{Cl}_2(\text{s}) \mid \text{Hg}$

The cell potential $E^{\circ}\text{cell} = E^{\circ}\text{R} - E^{\circ}\text{L}$

$E^{\circ}\text{R}$ = Reversible potential of SCE (= 0.2420 V), for Ag/AgCl (= 0.22249V).

$E^{\circ}\text{L}$ = Reversible potential $E_{\text{Zn}/\text{Zn}^{++}}$

$$E_{\text{Zn}/\text{Zn}^{++}} = E^{\circ}\text{Zn}/\text{Zn}^{++} - \frac{RT}{2F} \ln (a_{\text{Zn}^{++}}/a_{\text{Zn}})$$

Since, $a_{\text{Zn}} = \text{unity}$ and $a_{\text{Zn}^{++}} = 0.1 \times 0.15$ (activity coefficient = 0.15), we get

$$E_{\text{Zn}/\text{Zn}^{++}} = E^{\circ}\text{Zn}/\text{Zn}^{++} - (0.0592/2) \log (0.015.)$$


$E_{\text{cell}} = E^{\circ}\text{Zn}/\text{Zn}^{++} - 0.0296 * \log 0.015 + 0.2420$ volts, and therefore,

$$E^{\circ}\text{Zn}/\text{Zn}^{++} = E_{\text{cell}} + 0.0296 * \log 0.015 - 0.2420 \text{ volts}$$

$$E^{\circ}\text{Zn}^{++}/\text{Zn} = - E^{\circ}\text{Zn}/\text{Zn}^{++}$$


(B) To determine the normality of the ferrous ammonium sulphate solution by titrating potentiometrically with 0.5 N $\text{K}_2\text{Cr}_2\text{O}_7$ solution.

Pipette out 25mL of ferrous solution into a 100mL beaker. Dip a platinum electrode into the solution. Connect this with the saturated calomel half-cell through the salt bridge to form the following galvanic cell:



Connect the +ve and -ve half cells to +ve and -ve ends of a digital voltmeter. Measure the EMF of this cell as a function of $[\text{Fe}^{2+}]$ (i.e. vol. of $\text{K}_2\text{Cr}_2\text{O}_7$) directly using the digital voltmeter in the circuit.

Add by means of a burette, small quantities of 0.5N $\text{K}_2\text{Cr}_2\text{O}_7$ solution. Stir well and note the reading on the voltmeter after each addition. Record your results as per the format of Table I. Plot the e.m.f. against mL of $\text{K}_2\text{Cr}_2\text{O}_7$ solution added. It may be noted that after the equivalence point, the galvanic cell will be



Note the point (mL of $\text{K}_2\text{Cr}_2\text{O}_7$ solution, say V mL) where there is a relatively large change of potential. That will be the equivalence point and it has to be obtained as shown in figure. On another graph, plot the first derivative of e.m.f. $(\Delta E)/(\Delta V)$ against V_{avg} of $\text{K}_2\text{Cr}_2\text{O}_7$ added by recording your data as per the format of Table II. The inflexion point corresponding to peak of the curve gives a more reliable equivalence point.

The normality of the ferrous solution may be evaluated, using the relation:

Normality of the Ferrous solution = $0.5 \times V/25$ (N)

The red-ox reaction is:



Two model tables for recording your results follow. The actual end point will vary and therefore, the range in which titration should be done with smaller volume change has to be determined individually. The format of these tables is only to serve as an example.

Table I:

Volume of 0.5N $\text{K}_2\text{Cr}_2\text{O}_7$ (mL)	E.M.F (volts)	Volume of 0.5N $\text{K}_2\text{Cr}_2\text{O}_7$ (mL)	E.M.F (volts)
1.0		4.6	
1.5		4.8	
2.0		5.0	
2.5		5.2	
3.0		5.4	
3.5		5.6	
4.0		6.0	
4.2		7.0	
4.4		8.0	

Table II:

Volume of $\text{K}_2\text{Cr}_2\text{O}_7$ (V)	E.M.F (volts)	ΔE	ΔV	$\Delta E/\Delta V$	V_{avg}
1 mL	0.32				
2 mL	0.40	0.08	1.0	0.08	1.5
3 mL	0.53	0.13	1.0	0.13	2.5

Suggested Reading:

1. Physical Chemistry, P.W. Atkins, 5th Edition (ELBS/OUP) 1994, Pages: 324-348.
2. Vogel's Textbook of Quantitative Analysis revised by G. H. Jeffery, J. Basset J. Mendham and R. C. Denny, 5th Edition, Pages: 573-582.

Hazards and Preventions:

- As with all chemistry laboratories, lab coats, goggles or safety glasses and closed-toe shoes are essential.
- Make sure connections of wires are proper, otherwise there are chances of short circuit.
- Sulphuric acid is very **hazardous**. Read MSDS for the use of Sulphuric Acid properly. It is provided here <https://www.finarchemicals.com/msds/Sulphuric%20acid.pdf>
- Make sure no water or solvent is spilled on the batteries.
- No discarding of chemicals in the sink. Ask Lab Staff/supervisor for the procedure for proper disposal.
- Report any incident with promptness to the lab supervisor/staff.

Additional questions related to Electrochemistry:

1. What do you mean by an electrochemical interface?
2. What is an Electric double layer?
3. What is a Zeta Potential?
4. Does the sign of Zeta potential differ from the sign of the actual surface charge?
5. In your opinion, why do you think electrochemistry is important for the modern science?
6. What is an overpotential?
7. What is the difference between Power Density and Energy Density of a Battery?
8. In electrochemical cell, why is a salt bridge required? And how does it work?
9. If the Zn/Zn^{2+} is the cathode and Mg/Mg^{2+} is anode in the setup of an electrochemical assembly, and the E^0_{cell} is +0.65 V. The reactions occurring at both the electrodes are:
10. Standard Electrode potential for the half reactions, Fe^{2+}/Fe is -0.44 V and Cu^{2+}/Cu is 0.34 V. E^0_{cell} is given by _____ V
11. If you are given the value of the Standard Electrode potential of a cell, E^0_{cell} , the reaction occurring under standard conditions will be spontaneous is E^0_{cell} is _____.
12. Choose the correct option:
13. For a reaction having Cu/Cu^{2+} ($E^{\text{Cu}/\text{Cu}^{2+}} = -0.34$ V) and Zn/Zn^{2+} ($E^{\text{Zn}/\text{Zn}^{2+}} = 0.76$ V) as half cells, Cu^{2+} will be product/reactant.
14. State whether True or False

Salt bridge is a glass tube filled with agar jelly containing electrolytes such as KCl and KNO_3 , their role is to ensure that the solutions in the cell remain electrically neutral and electrons flow from one half cell to the other.

15. State whether True or False
 - (a) In a potentiometric titration, the emf of the cell is equal to zero at equivalence point.
 - (b) Potentiometric titrations have an added advantage over volumetric titrations, in the fact that they can be easily performed for turbid or colored solutions.
16. The Nernst equation for reaction of Ferrous ammonium sulphate with Potassium Dichromate, $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$ is:

$$E = E^0_{\text{cell}} - \frac{RT}{nF} \frac{[\text{Cr}^{3+}]^2 [\text{Fe}^{3+}]}{[\text{Cr}_2\text{O}_7^{2-}] [\text{Fe}^{2+}]}$$

The value of n is _____.

17. State whether True or False

For the reaction: $Mg^{2+}(aq) + 2e^{-} \rightarrow Mg(s)$, $E^{\circ} = -2.37\text{ V}$, The electrode potential varies linearly with the activity/concentration of Mg^{2+} .

18. State whether True or False

In a Galvanic cell, electrons flow in the external circuit from anode to cathode while releasing electrical energy.

19. In the potentiometric titration of Ferrous Ammonium sulphate with Potassium Dichromate, after the equivalence point, addition of Potassium dichromate increases the EMF. This increase is due:

Excess of $Cr_2O_7^{2-}$ ions

Reduction of Fe^{3+} to Fe^{2+} ions

Oxidation of Cr^{3+} ions

Additional Learning Sources:


1. Li-ion batteries: <https://pubs.acs.org/doi/10.1021/acsenergylett.0c02584>
<https://iopscience.iop.org/article/10.1149/2.0251701jes/pdf>
2. Initiatives by GoI on Renewable Energy: <https://mnre.gov.in/new-technologies/hydrogen-energy> (Not all of these are related to Electrochemistry, though)
3. Fuel cell Technologies and progress in India: https://www.nasa.gov/centers/glenn/technology/fuel_cells.html#:~:text=Glenn%20is%20investigating%20three%20types,were%20used%20through%20the%201990s and <https://www.fchea.org/in-transition/2020/6/9/fuel-cell-and-hydrogen-development-in-india>
4. Future of Electrochemistry: <https://link.springer.com/content/pdf/10.1007/s10008-020-04585-3.pdf>
5. Modern Batteries: Metal-Air Electrochemical Cell

2. Spectroscopic evaluation of removal efficiency for a Pharma Pollutant in water using MAC

Aim: To evaluate removal efficiency for a Pharmaceutical Pollutant in Water Using a Magnetite-Activated Carbon Nanocomposite by Colorimetry

Overview: In the last decade, traces of pharmaceuticals, typically in the parts per billion (ppb) or parts per trillion (ppt) range, have been reported in the water cycle, including surface waters, wastewater, ground water and drinking water. Their presence in water, especially in drinking water, even of these very low concentrations, has raised concerns regarding the potential risks to human health. Conventional drinking-water treatment processes have been shown, in many cases, largely ineffective in removing pharmaceuticals. Advanced water treatments such as activated carbon (AC) are shown to be able to achieve very high removal rates for targeted pharmaceutical compounds. The effectiveness of AC is due to its extended surface area, complex porous structure, high adsorption capacity and high degree of surface reactivity.

In this lab, you will modify the AC by embedding magnetic magnetite (Fe_3O_4) nanoparticles (MNPs) onto the AC matrix using a simple one-step room temperature synthetic process. The MNPs are produced by mixing Fe(II) and Fe(III) salts together in a basic solution as shown in equation 1:



The resultant product, referred to as MAC, represents an example of an exciting class of advanced materials, known as nanocomposites. A nanocomposite is a multiphase solid material where one of the phases has one, two, or three dimensions of less than 100 nm. Examples of familiar natural nanocomposites include hard and tough bones, teeth, molluscan shells and sea urchin spine. They are built of alternating nanolayers of hard inorganic minerals and soft stress dissipating organic polymers. Nanocomposites offer unique properties that can find valuable technological applications and that cannot be achieved through the use of conventional materials

In this lab, you will show that MAC nanocomposite is also effective at removing pharmaceutical pollutants containing benzene rings from water using a derivative of aspirin, salicylate dianion, which serves as a metabolite of aspirin. Aspirin, also known as acetylsalicylic acid, is one of the most widely used medicinal agents. One of the major metabolites of aspirin is salicylic acid, which takes the forms of salicylates in neutral or basic solutions. Figure 2 shows the molecular structures of aspirin, salicylic acid, and salicylate dianion, all contain one benzene ring.






Figure 2. The molecular structures of a) acetylsalicylic acid or aspirin, b) salicylic acid, and c) salicylate dianion.

One of the spectrophotometric methods (using Beer-Lambert's law) for the determination of the amount of aspirin present in water is based on the highly colored complex formed between the aspirin derivative, salicylate dianion, and iron(III) ion. The complex has a maximum absorption at 530 nm. Complexation of aspirin is produced by a two-stage reaction. First, aspirin is heated in a solution of sodium hydroxide, forming the salicylate dianion as shown in Equation 2:



Then, the addition of acidified iron(III) ion produces the violet Fe^{3+} -salicylate complex or tetraaquaosalicylatoiron (III) complex as shown in Equation 3:



Safety:

Gloves and goggles must be worn at all times. Aqueous ammonia, FeCl_2 , FeCl_3 , HCl and NaOH are corrosive. FeCl_2 is also a mutagen. Handle all of these materials with care and wash immediately with water in case of skin contact. The magnets strongly attract each other. Avoid placing two magnets in close proximity to avoid pinching. Rare earth (Neodymium) magnets are fragile and can break easily. They have a strong magnetic field that can damage electronic equipment (mobile phones, watches, etc.) and magnetic data storage media.

Chemicals & Apparatus:

- 0.1g of activated carbon
- 4.0 mL of 1M FeCl_3 solution
- 1 mL of 2M FeCl_2 solution
- 50 mL NH_3 solution
- 15 mL of the “Pharm. Pollutant” or salicylate dianion stock solution, equivalent to 1600 ppm of aspirin
- 25 mL of acidified 0.15 M iron (III) chloride solution
- a 5-mL or 10-mL graduated pipette/ micropipette
- a stirring rod
- 100/ 150 mL beakers
- a burette
- a neodymium magnet

- test tubes
- a 0.50-mL graduated pipette/ micropipette
- 10-mL volumetric flasks
- 2-3 Disposal droppers
- a Colorimeter and 2 cuvettes

Procedure:

A. Preparation of Magnetite-Activated Carbon (MAC) Nanocomposite

1. Add 0.1 g of activated carbon, 4.0 mL of 1M FeCl_3 solution and 1.0 mL of 2M FeCl_2 solution to a 150-mL beaker and mix the content with a stirring rod for about 2 minutes.
2. Fill a burette with 1.4 M NH_3 solution. Add a total of 50 mL NH_3 solution dropwise to the content of the beaker in 5 minutes. Throughout the addition, continuously and vigorously mix the content in the beaker with the stirring rod. CAUTION: Although 1.4 M NH_3 is fairly dilute, NH_3 is corrosive.
3. Place the beaker on a magnet to allow the MAC nanocomposite particles to settle.
4. Decant (pour off) and discard the clear liquid without losing a substantial quantity of solid. This works best if you keep a magnet under the container.
5. Rinse with 5 mL deionized water from a wash bottle and then decant the rinse with the magnet under the container as before.

B. Removal of Pharmaceutical Water Pollutant- an Aspirin “Metabolite”




Figure 5: Graphical representation of pharmaceutical pollutant removal lab procedure.

1. Obtain 15 mL of the “Pharm. Pollutant” stock solution in a clean beaker, labeled as “**A: Control**”.
2. Add 10 mL of the “Pharm. Pollutant” - Control” solution in the 150-mL beaker containing the MAC nanocomposite prepared in Part A. Stir the water vigorously for 5 minutes using a stirring rod.
3. Place the beaker as in Step 3 on a magnet to allow the MAC nanocomposite particles to settle.
4. The upper liquid is the “**treated Pharm. Pollutant**” water. Transfer about 5 mL of the treated “Pharm. Pollutant” water using a disposal plastic dropper to a clean and dry test tube labeled as “**B: Treated**”.

C. Spectrophotometric Analysis

1. Complexing Salicylate Dianion with acidified iron(III) ions:

- a) Obtain 25 mL of acidified 0.15 M iron (III) chloride solution in a clear and dry beaker. Label the beaker.
- b) Prepare a 0.50-mL graduated pipette by first rinsing it with deionized water twice and then the “Pharm. Pollutant” stock solution twice (in the beaker labeled “A: Control”). Using the 0.50-mL graduated pipette, carefully transfer 0.40 mL of the “Pharm. Pollutant” stock solution into a clean and dry 10-mL volumetric flask. Using a clean disposable dropper (can be reused for the same solution in Step C1(c)), carefully add acidified 0.15 M iron (III) chloride solution to the mark. Mix the content and label the volumetric flask as **“A: Fe³⁺-salicylate complex – Control”**
- c) Repeat Steps C1(b) with the treated “Pharm. Pollutant” water in the test tube labeled as “B: Treated”. Label the volumetric flask as **“B: Fe³⁺-salicylate complex – Treated”**

2. Measurements with Spectrophotometer:

Standard Curve Construction

- a. Measure and record the absorbances of a set of Fe³⁺-salicylate complex standards (in cuvettes) with a Colorimeter set at 530 nm. Use the pale-yellow iron (III) chloride solution as a blank to set the instrument calibration.
- Table for Sets of Fe³⁺-salicylate Complex Standard Solutions for Beer’s Law Construction:

Standard Solution	Vol. of standard salicylate dianion solution (mL)	Dilute these solutions with acidified 0.15M iron (III)chloride solution up to 10 mL mark in given volumetric flasks
Standard 1	0.5	
Standard 2	0.4	
Standard 3	0.3	
Standard 4	0.2	
Standard 5	0.1	

- b. Plot a graph of Absorbance vs Concentration in ppm (Beer’s Law) for the Fe³⁺-salicylate complex system.

“Pharm. Pollutant” Concentration Determination

- c. Measure and record the absorbance of the solution transferred into a cuvette from the volumetric flask labeled as “A: Fe³⁺-salicylate complex – Control” using the spectrophotometer set at 530 nm.
- d. Repeat Step 2(c) for the solution from the volumetric flask labeled as “B: Fe³⁺-salicylate complex – Treated”.

Calculations:

- a. Use your data obtained in 2(c) and 2(d), determined the mg/L (or ppm) of equivalent aspirin in the Fe³⁺-salicylate complex samples from the calibration curve and best-linear-fit equation in the form of $A = k \cdot C + b$ reported in 2(c).
- b. Determined the salicylate dianion concentrations in mg/L (or ppm) of Aspirin in the “control” and “treated” “Pharm. Pollutant” (stock) samples before complexation. Complexation was a dilution process for salicylate dianion. $[C \cdot V = C_{\text{diluted}} \cdot V_{\text{diluted}}; V = 0.4 \text{ mL}; V_{\text{diluted}} = 10.0 \text{ mL}]$
- c. Determine MAC’s Removal Rate using Equation 6.

$$\text{Removal Rate} = \frac{100 \times (C_0 - C_t)}{C_0}$$

d. Determine MAC's Capacity using the Equation 7

$$\text{Removal Capacity} = \frac{(C_0 - C_t)V}{m}$$

e. where $V = 10.0 \text{ mL}$ = volume of "Pharm. Pollutant"-water treated. Use the theoretical value of 0.10 g AC in MAC for the calculation.

f. Determine the percent error of the Method for determining the concentration of the "control"- "Pharm. Pollutant" or salicylate dianion stock sample. The expected concentration of this "Pharm. Pollutant" solution is 1600 ppm Aspirin.

Chemical Disposal:

All wastes containing the magnetite nanoparticles should be placed in a waste container. Liquid waste without nanoparticles can go down the drain with excess water. Solid waste can be disposed of in the trash.

Names: _____ Section: _____

DATA SHEET – Spectrophotometric Analysis

Concentration, ppm Aspirin	Absorbance at 530 nm
16	
32	
48	
64	
80	
Fe ³⁺ -salicylate Complex Control, A ₀	
Fe ³⁺ -salicylate Complex Treated, A	

Beer's Law Equation: _____

R² value: _____

Fe³⁺-salicylate Complex Control, C₀, diluted: _____ ppm Aspirin

Fe³⁺-salicylate Complex Treated, C_{diluted}: _____ ppm Aspirin

"Pharm. Pollutant" or Salicylate Dianion Control (stock), C₀: _____ ppm Aspirin

"Pharm. Pollutant" or Salicylate Dianion Treated (stock), C: _____ ppm Aspirin

MAC's Removal Rate: _____

MAC's Removal Capacity: _____

"Pharm. Pollutant" or Salicylate Dianion Control (stock) C₀, expected = _____

"Pharm. Pollutant" of Salicylate Dianion Control (stock) C₀, determined = _____

% Error = _____

References:

- *J.Chem. Educ.* 2019, 96, 4, 751-755

3. Determination of Iron

Objective: To estimate the amount of ferrous and ferric ion in a solution containing both.

Introduction:

Iron belongs to the 3d transition series of metals with electronic configuration $3d^6 4s^2$ and forms several oxides with different oxidation states and crystal structures. Of these oxidation states, +2 and +3 states are commonly encountered, the oxides being abundant in nature. Common examples are magnetite (Fe_3O_4) and hematite (Fe_2O_3).

Iron oxide is a raw material for producing iron and steel and a catalyst for many industrial processes (eg., manufacture of NH_3). The general importance of iron oxides in other disciplines of technology besides metallurgy lies in the fact that these oxides serve as raw materials for synthesis of magnetic materials. For instance, γFe_2O_3 is used as a magnetic material in recording tapes. Iron oxides mixed with other metallic oxides are used in magnetic materials such as ferrites, perovskites etc., which are useful in semiconductor technology, microwave devices, ferroelectrics etc.

The properties of these iron containing materials depend on the purity and the required stoichiometry. The presence of impurities and also deviation from the stoichiometry adversely affect the properties and hence their technological importance. It is therefore obvious that strict quality control is of importance which involves the estimation of the constituent elements. Further, these elements (eg., Fe_3O_4) invariably contain iron in different oxidation states namely +2 (Ferrous) and +3 (Ferric) and their accurate quantitative determination is of considerable importance.

The quantitative estimation of ferrous and ferric either separately or in a mixture containing both, can be carried out by conventional volumetric and gravimetric methods or using instrumental methods such as spectrophotometric methods. The choice of method depends on the percentage of iron present. The present exercise deals with the volumetric estimation of iron in +2 and +3 oxidation states as the required titrations are relatively simple and less time consuming.

Principle:

The quantitative estimation of Fe^{2+} either separately or in a mixture containing Fe^{3+} can be carried out by volumetric titration involving an oxidising agent (either $KMnO_4$ or $K_2Cr_2O_7$). The estimation of Ferric generally involves the quantitative reduction of ferric to ferrous state and subsequent estimation of Ferrous by titration with a standard oxidizing agent such as $KMnO_4$ or $K_2Cr_2O_7$. The reduction of Fe^{3+} to Fe^{2+} can be brought about by one of the following reagents (a) Stannous chloride (b) Amalgamated Zinc (c) Sulphurous acid (d) Titanous salts (e) Hydrogen Sulfide and (f) Zn and dil H_2SO_4 .

Procedure:

Make up the solution supplied in 100 mL standard flask by adding small amounts of distilled water very carefully to bring it upto the 100 mL mark (lower meniscus coinciding with the mark) using distilled water. When nearing the 100 mL mark add additional distilled water, a drop at a time. Shake thoroughly to make the solution homogenous. Record your flask No.

(a) Estimation of Ferrous: Pipette out 10 mL of diluted solution into a clean (need not be dry) conical flask. Add 1 test tube full of 2N H₂SO₄. Titrate with KMnO₄ taken in the burette. Do not add more than 0.5 mL of KMnO₄ at a time. Continue the titration till the solution changes from colourless to light pink. Repeat the titration till two successive burette readings do not differ by more than 0.1 mL.

(b) Estimation of Ferric: Pipette out 10 mL of the diluted solution into a clean conical flask. Add about 150 mg of Zn powder. Heat the solution on a water bath with continuous swirling till the solution turns almost colourless. Add 1 test tube full of 2N H₂SO₄ and allow it to stand for some time till all excess Zn has reacted. (If some Zn is left, add some more H₂SO₄). Cool the conical flask and titrate with KMnO₄ as in (a).

Calculations:

$$1000 \text{ mL} = 1\text{N KMnO}_4 = 56 \text{ gm. Fe}^{2+}$$

$$(a) V_{\text{solution}} = 10 \text{ mL.} \quad N_{\text{KMnO}_4} = a = 0.1\text{N,} \quad V_{\text{KMnO}_4} = b \text{ mL.}$$

$$\text{Amount of Fe}^{2+} \text{ in the given solution} = \frac{10 \times a \times b \times 56}{1000} = x \text{ g}$$

$$(b) V_{\text{solution}} = 10 \text{ mL.} \quad N_{\text{KMnO}_4} = a = 0.1\text{N,} \quad V_{\text{KMnO}_4} = c \text{ mL.}$$

$$\text{Total Fe}^{2+} \text{ in the given solution} = \frac{10 \times a \times c \times 56}{1000} = y \text{ g}$$

$$\text{Amount of Ferric Iron} = (y - x) \text{ g}$$

Suggested Reading:

Vogel's Textbook of Quantitative Analysis Revised by G.H.Jeffery, J.Basset, J.Mendham and R.C.Denny, 5th edition, Pages: 261, 368-372, 409-416

4. Oscillatory Chemical Reaction

Objectives:

- a) To introduce the students to a fascinating example of an oscillating chemical reaction.
- b) To introduce the concept of non-equilibrium thermodynamics.
- c) To speculate on the possible causes and applications of oscillating chemical phenomena.

Introduction:

The heartbeat, the tick-tock of a grandpa clock and the motion of waves at a beach are fascinating, and with history repeating itself, periodic phenomena have a certain appeal. At first these reactions were treated with some scepticism because it was felt that true chemical oscillation would be contra-thermodynamic. But now there are numerous indisputable experimental examples and a theory of non-equilibrium thermodynamics. The theory was developed by Lars Onsager of Yale University and Ilya Prigogine of the Free University, Brussels. The theory showed that although chemical oscillation about equilibrium, analogous to the motion of a pendulum, was indeed an impossibility, oscillation far from equilibrium is perfectly consistent with physical laws.

Some well-known oscillating chemical reactions

Bray-Liebhafsky (BL) Reaction:

The first example of a homogenous oscillating chemical reaction was discovered in 1921 by William C. Bray at the University of California, Berkeley and later studied exhaustively by Herman Liebhafsky at Texas A & M University. The reaction is the iodate catalyzed decomposition of hydrogen peroxide. The rate of oxygen evolution and intensity of the iodine color vary in a periodic fashion.

Belusov-Zhabotinskii (BZ) Reaction:

This reaction was discovered by B. P. Belusov at the Institute of Biophysics in the USSR in 1951. The discovery was initially ignored but later taken up by A M Zhabotinskii at Moscow State University. The reaction mixture consists of malonic acid, acidified bromate and a ceric salt and shows oscillations in bromine concentration, going from yellow to colorless and back.

Briggs-Rauscher (BR) Reaction:

Two San Francisco high school teachers, T.S. Briggs and W. C. Rauscher, developed a hybrid of the BZ and BL reactions that provides a visually striking lecture demonstration that oscillates for 20 minutes or more before settling down to thermodynamic equilibrium. The reaction mixture consists of KIO_3 , H_2O_2 , HClO_4 , malonic acid, MnSO_4 and starch to visualize the oscillations in I_2 concentration.

Mechanisms for chemical oscillators:

Mechanisms for oscillatory reactions are rarely, if ever, simple. They generally involve 10 or more stoichiometrically independent species and perhaps twice as many elementary steps. Although they are less elaborate than the mechanisms that describe combustion and atmospheric processes, they are nonetheless too complex to be grasped intuitively. Development of a mechanism for an oscillating reaction usually requires both extensive computer simulation and a variety of kinetic experiments aimed at establishing the rate laws and rate constants of as many elementary steps as possible. The remainder are estimated either by comparison with related reactions or by fitting the calculated and experimental data.

The earliest model of oscillatory chemical behaviour proposed by A. J. Lotka in 1920 consists of three reactions, two of them autocatalytic. The precursor A is taken to be in large excess so that its concentration has a constant value, say a.

Representing the concentrations of the autocatalytic species X and Y by x and y, respectively, it can be shown that for any constant C the curve given by

$$k_2(x+y) - k_3 \ln x - k_1 \ln y = C$$

is a solution of the three rate equations. The solution equation represents a closed curve, almost an ellipse, in x-y space. (Try plotting it). Therefore, as the system follows the curve, the concentrations of X and Y vary in a periodic fashion.

Application of chemical oscillators:

Does chemical oscillation have any practical applications, or is it merely a source of attractive, but frivolous lecture demonstrations? Many of the potential practical applications of the knowledge gleaned from the study of chemical oscillators lie in biology. It has been suggested that certain processes might operate more efficiently in an oscillatory than a steady-state fashion. Model calculations show that glycolysis may have evolved as an oscillatory chemical process because this mode of operation is a few percent more efficient than the steady-state alternative. By an examination of what happens when chemical oscillators are perturbed at different points in their cycles, a plausible explanation of the onset.

There may be important analogies between dynamical processes in chemistry and geology. Theories used in explaining periodic Liesegang rings (the periodic precipitation phenomena that occur when ions like silver and chromate, that from a sparingly soluble salt, diffuse towards each other from opposite ends of a gel filled tube) have been adapted to the study of the origin of layering patterns in certain rock formations. Although the time and length scales are quite different, the fundamental physical principles remain the same.

In today's exercise we are going to perform a variation of the Briggs-Rauscher reaction.

Safety:

1. Wear goggles at all times. Use gloves when handling the chemicals.
2. Hydrogen peroxide solutions (3.579 M) are prepared from 30% hydrogen peroxide, which can cause skin and eye irritation on contact, and abdominal pain if ingested. The 0.2 M sulfuric acid may cause harm to the eyes or skin and is harmful to the digestive system if ingested. Dilute product solutions of iodide/iodine may stain the skin.
3. Check with the TA for disposal of the waste liquid and materials. DO NOT DUMP ALL CHEMICALS IN THE SINK.

Procedure:

Part1: Observation of patterns in a diffusion controlled oscillating reactions

For certain reactions, the reactants (and intermediates) diffuse slowly through the solution. Interesting patterns emerge when such a reaction is also oscillatory. We will observe and play with such a reaction:

1. In a 50 mL beaker, mix the 6 solutions in the following order: MnSO_4 (0.005 M, 5 mL), H_2SO_4 (0.2M, 5 mL), malonic acid (0.03 M, 5 mL), KIO_3 (0.15M, 5 mL), H_2O_2 (3.579 M, 10 mL), and 2 mL 0.5% starch indicator for visualizing the patterns.
2. Transfer the above solution quickly to the petri dish.
3. Observe any changes in colour and any pattern formation.
Start with step 2, while the patterns are emerging.

Part 2: Observation and measurement of an oscillatory reaction

Now we will look at an example where the entire solution changes colour periodically and measure the timings.

1. Mix the following solutions in the order given below so that in the mixture each component has the concentration mentioned:

KIO ₃	0.07 M
H ₂ SO ₄	0.05 M
Malonic acid	0.05 M
MnSO ₄	0.007 M
Starch	0.01%
H ₂ O ₂	0.06 M

Watch the fun. Report your observations carefully.

2. Note the timings of the oscillations using the provided stop-watch.
3. Consider the mechanism provided above. Write the rate law corresponding to [X] and [Y].
If initially, [X]=[y]=0, qualitatively sketch [X] and [Y] as a function of time.
4. Change concentrations of the different solutions and record the timings of oscillations.
5. Interpret any changes in oscillation timescales based on the above mechanism.

Suggested reading:

1. The above write-up has been adapted from, “Patterns in Time and Space - Generated by Chemistry”, I. R. Epstein, C and E News, March 1987.
2. Part 1 of the procedure taken from
<https://pubs.acs.org/doi/abs/10.1021/acs.jchemed.0c00892>
3. “An Oscillating Iodine Clock”, T. S. Brigg and W.C. Rauis cher, Journal of chemical Education., Vol no. 50, Issue no 7,Page no 496,year 1973
4. “Oscillating Chemical Reactions”,I.R. Epstein, K. Kustin, P. DeKepper and M. Orban, Scientific American, Vol no.248, Page no.112,year 1983
5. More on oscillatory reactions:
[https://chem.libretexts.org/Bookshelves/Physical_and_Theoretical_Chemistry_Textbook_Maps/Map%3A_Physical_Chemistry_for_the_Biosciences_\(Chang\)/09%3A_Chemical_Kinetics/9.11%3A_Oscillating_Reactions](https://chem.libretexts.org/Bookshelves/Physical_and_Theoretical_Chemistry_Textbook_Maps/Map%3A_Physical_Chemistry_for_the_Biosciences_(Chang)/09%3A_Chemical_Kinetics/9.11%3A_Oscillating_Reactions)
6. Video of pattern formation in oscillating reaction:
<https://www.youtube.com/watch?v=jRQAndvF4sM>

Moodle questions:

1. For the elementary reaction $A + X \rightarrow 2X$, propose the rate law with respect to concentration of A.
 - a. $d[X]/dt = k[A][X]$
 - b. $-d[X]/dt = k[A][X]$
 - c. $\frac{1}{2} d[X]/dt = k[A][X]$
 - d. $-d[X]/dt = \frac{1}{2} d[X]/dt k[A][X]$
2. Which of the following curves will best represent the concentration of X with time?
3. Which of the following best represents the mechanism for an oscillatory reaction:
 - a. The products keep converting back to reactants, leading to oscillations.
 - b. The rate-determining step keeps changing with time depending on concentrations of different species.

- c. Excess of reactant concentration leads to oscillations in the concentration of intermediates.
 - d. The autocatalytic step in the mechanism leads to oscillations as this step can go both in the forward as well as the backward direction.
4. What safety features are correct for this experiment (select all that apply):
 - a. Dispose of chemicals in the sink when used
 - b. Concentrated H_2O_2 can cause skin and eye irritation and must be used with care.
 - c. Dilute sulfuric acid can harm the eyes and skin if come in direct contact.
 - d. Iodine solutions can stain the skin.
 5. How does Briggs-Rauscher work?
 6. What is origin of amber color in Briggs-Rauscher reaction?
 7. What will be the effect on oscillation time in Briggs-Rauscher reaction if acetone is used as a solvent?
 8. Is Briggs-Rauscher reaction reversible?
 9. What is the role H_2O_2 in the oscillatory reaction?
 10. When does Briggs-Rauscher reaction stop?
 11. What are the main elements of Belousov-Zhabotinsky reaction?
 12. What is the role of malonic acid in Belousov-Zhabotinsky reaction?
 13. Do oscillating reactions go on forever?
 14. Does the frequency of oscillation remain same throughout the process?
 15. How does temperature effect the oscillation?


Sr. No.	Vol. of KIO ₃ (mL)	Vol. of H ₂ SO ₄ (mL)	Vol. of Malonic acid (mL)	Vol. of MnSO ₄ (mL)	Starch	Vol. of H ₂ O ₂ (mL)	Time for first appearance of blue colour	No. of Oscillations	Time period of oscillation	Observations
1	3	3	3	3	6 drops	3				
2	3	4*	3	3	6 drops	3				
3	3	1*	3	3	6 drops	3				
4	3	3	0*	3	6 drops	3				
5	3	3	3	0	6 drops	3				

Interpretation of observation marked with*

5. Investigating chemical kinetics using a self-constructed polarimeter

Objective: i) To construct a working polarimeter using the parts given, and ii) To determine the rate constant for the inversion of sucrose using the self-constructed polarimeter.

Principle: The reaction for the inversion of sucrose is as follows:



This reaction is catalysed by H_3O^+ ions.

The rate of this reaction depends on the concentrations of sucrose, water and H_3O^+ . Since a large excess of water is needed, its concentration may be considered to be a constant and thus making the rate independent of water concentration. Therefore, for a given concentration of H_3O^+ , the rate of the reaction depends only on the concentration of sucrose at a given temperature. If the reaction is of first order, the differential rate expression is

$$\frac{-d[\text{sucrose}]}{dt} = k[\text{sucrose}]$$

Where k is the specific rate constant (or simply the rate constant). In the integral form, the equation becomes:

$$\ln [\text{Sucrose}] = kt + \text{Constant.}$$

With the limits $[\text{Sucrose}]$ at $t = 0$ and $[\text{Sucrose}]_t$ at time t , the equation is

$$k = \frac{2.303 \log_{10}}{t} \frac{[\text{sucrose}]_0}{[\text{sucrose}]_t}$$

The rate of a chemical reaction can be followed by measuring the concentration of either the reactant or product as a function of time. This can be done by either chemical analysis or by choosing a physical property which is a linear function of concentration. Physical methods are often more rapid than chemical analysis and one can obtain a large amount of data within a given (short) time. One such physical property is optical activity.

Typically, optical activity is measured by an instrument called polarimeter. In such an experiment, a plane polarized light of known polarization direction is passed through a solution under study and the angle by which the plane polarized is rotated by the solution is measured. The extent of rotation of the plane polarized light is a measure of the optical activity of the solution.


Reagents required: HCl (2M), sugar solution (20%)

Apparatus required: Appropriately cut linear polarizing film (1), Lego brick base (1), protractor mounted on a cardboard, a plastic dial with an attached polarizer, LEDs (yellow and red), crocodile clips, resistors (10 – 60 Ohms), battery (3V), digital multimeter, Cuvette, 10 mL pipette, beaker.


Procedure:

Construction/assembly of a polarimeter:

Front view of self-constructed polarimeter



Rear view of self-constructed polarimeter:



- 1) Using crocodile clips connect the battery to the source LED (yellow) via a suitable resistor (10 – 60 Ohms)

- 2) Insert the dial with an attached polarizer onto the dial, as shown in the figure. The direction of polarization of this polarizer is orthogonal to the base
- 3) Insert the appropriately cut polarizer with its polarization axis perpendicular* to the polarizer mounted on the dial
- 4) Connect the back LED to the multimeter via crocodile clips

*In principle, the polarization of both the polarizers can be parallel. This only affects the way in which the readings are recorded (i.e., rotating the dial to obtain maximum voltage in multimeter, rather the minimum, as explained below), but not the actual readings.


For more details about the set-up check out the supporting information of the following paper *J. Chem. Educ.* **2020**, *97*, 2196.

Measuring optical rotation using the above constructed polarimeter:

Recording calibration reading: Note the voltage at zero degree on the dial. Move the dial in steps of 0.5° on either side of zero degree on the dial while observing the voltage on the multimeter attached to the detector LED. The lowest reading (mV) in the multimeter represents the point of minimum light transmittance. Check the point of minimum light transmittance by moving the dial on either side to obtain a stable zero transmittance point. Note the degrees on dial for minimum transmittance (θ_0). This is the calibrated zero. At this point both the polarizers are very close orthogonal with respect to each other.

Measuring optical activity with the self-constructed polarimeter: Placing a cuvette with an optical active solution in the instrument will alter the optical zero of the system due to the rotation of the plane polarizer light by the optically active solution. To estimate the extent of rotation, move the dial slowly to each side of zero to determine if the rotation is dextro- or levorotatory (see the below picture for the convention), and to locate the new optical zero as described above. Move the dial in smaller steps around the optical zero to obtain accurate results of rotation. Note the degrees on the dial for minimum light transmittance (θ_x). The magnitude of net rotation = $\theta_x - \theta_0$. The direction of rotation can be ascertained based on the below convention.

Cross check the magnitude and direction of rotation with a commercial polarimeter available in the laboratory.



To measure A_{∞} (optical rotation at $t = \infty$).

Fill the cuvette with given sucrose solution (at $t = \infty$) taking care to avoid air bubbles. Place the cuvette in the appropriate position of the self-constructed polarimeter and measure the angle of rotation, as described above.

To measure A_0 (optical rotation at $t = 0$).

Pipette out 20 mL of 20% sucrose (supplied) into a clean beaker. To this add 20 mL distilled water (by means of pipette). Make the resulting solution homogeneous. Fill the cuvette with the solution and measure the angle of rotation.

To measure A_t (optical rotation at different time intervals).


Pipette out 20 mL of 20% sucrose solution in a beaker. To this, add 20 mL (by a pipette) of 2M HCl. Note the time when HCl is added to the sucrose solution. Fill the cuvette with this solution and measure the angle of rotation at different time intervals (see table).

Time (t) minutes	Optical Rotation (A_t)	($A_t - A_{\infty}$)	Log ($\frac{A_0 - A_{\infty}}{A_t - A_{\infty}}$)	k
5				
10				
20				
30				
40				
60				

Precaution: Wipe the outer surface of the cuvette free of sugar solution with tissue paper before placing inside the polarimeter.

Additional questions:

1. In a polarimeter experiment, the rotation of the plane polarized light by the sample is due to – absorption, scattering, reflection of light or any other process taking place? Explain your answer.
2. Why is it preferred to use a sodium D-line as the source of light for polarimeter experiment? Can other sources of light be used?
3. Although optical rotation is taken as a characteristic signature of a chiral system or molecule, the below shown molecule does not show optical activity in spite of it being chiral (non-racemic). Explain why?



4. Why a circular cuvette is not used for hosting the sucrose solution and measuring the inversion process?
5. Suggest two possible sources of error in – i) construction of the polarimeter, and ii) measurement of the optical activity
6. If the time scale of inversion process in a carbohydrate/molecule is similar to the time scale of the polarimeter measurement, should polarimeter be used to understand such a process? Justify your answer.
7. Does the optical activity of a given molecule change if any of the following conditions are changed – i) solvent in which the measurement is carried out, ii) temperature, and iii) concentration. Justify your answer.

Additional resources:

1. A you tube demonstration of the building a polarimeter,
<https://www.youtube.com/watch?v=3v1nmc7gu7k&t=391s> and
https://www.youtube.com/watch?v=AZL7lgWG6DI&t=2s&ab_channel=CuriositasScience
2. Polarization of light (unpolarized, linearly polarized and circularly polarized light)
 - a. https://www.youtube.com/watch?v=8YkfEft4p-w&t=186s&ab_channel=PhysicsVideosbyEugeneKhutoryansky,
 - b. https://www.youtube.com/watch?v=PJHCADY-Bio&ab_channel=EricMickelsenEricMickelsen
 - c. https://www.youtube.com/watch?v=mjwQTL6G8Fs&ab_channel=MITOpenCourseWare

References:

1. Kvittingen, L.; Sjursnes, B. J., *J. Chem. Educ.* **2020**, 97, 2196.
2. Organic Chemistry, Morrison and Boyd, 6th Edition, Pages: 128-138 and 1191.

6. Complexometric Titration


Objective: Determination of total hardness of water using complexometric titration with Ethylenediaminetetraacetic Acid (EDTA).

Introduction:

Hard water contains high dissolved mineral content—usually, the higher the water hardness, the lower the toxicity of heavy metals to aquatic life. However, hard water is unsuitable for most industrial purposes. Laundries require soft water of zero hardness to avoid precipitation of soaps on textiles. Hard water removes soap from solution as an insoluble precipitate and is not available for detergent action until a sufficient amount is consumed for rendering the water soft. The presence of iron in hard water may lead to staining of fabrics. Boiler feed water should be softened in order to avoid scale formation on boiler walls and tubes leading to pitting corrosion and poor heat transfer. The dyeing industry needs water free from iron which can react with dyes to form insoluble lakes of undesirable shades. A number of chemical methods are available for softening water by precipitation as insoluble carbonates and hydroxides but these do not remove all mineral content since they introduce Na^+ and Cl^- ions. Demineralization can be achieved by distillation. Extent of hardness can be measured in terms of CaCO_3 concentration through titration with EDTA as detailed below.

Principle:

EDTA forms stable complexes with bivalent metal ions in basic or slightly acidic medium and it is widely used as a complexing agent for metal ions. If we represent EDTA by the formula H_4Y , its disodium salt will be $\text{Na}_2\text{H}_2\text{Y}$. The latter yields the complex-forming anion H_2Y^{2-} in aqueous solution. The reaction of EDTA with a bivalent cation M^{++} may therefore be written as:



The structural formula of H_2Y^{2-} is depicted below (Figure 1):




Figure 1. Structure of EDTA (H_2Y^{2-})

In titrating the metal ion solution with EDTA, the indicator usually chosen is Eriochrome Black T (EBT) (Figure 2). The EBT indicator forms a wine-red complex with the metal ion which breaks up giving way to the metal - EDTA complex. At the end point, plus a slight excess of EDTA makes the wine-red color disappear completely, leaving the metal-EDTA complex in solution and the blue color of free indicator appears. The color change at the end point is from wine red to blue-violet.




Figure 2. Structure of EBT

The color changes in the solution containing M^{++} ion and EBT in the presence of EDTA can be followed to comprehend the formation different complexes during the titration. An optical spectroscopy experiment will be performed (mentioned as part B in the following) to monitor the serial changes in the solution. The EBT molecule contains a highly conjugated azo ($-N=N-$) group that exhibits an absorbance feature in the visible region. This band shifts significantly upon the formation of M^{++} -EBT complex. The EDTA is a better chelating ligand (hexadentate) compared to EBT (tridentate). Hence, Addition of EDTA strips the M^{++} ion from the EBT to regenerate the free EBT ligand. The final step will revert the optical spectra back to the original condition.

Procedure:

Experiment A.

I) Standardization of EDTA.

Pipette out 10 mL of 0.02 M $MgSO_4$ soln. in a clean conical flask. Add $\frac{1}{2}$ test tube of buffer (pH = 10) and 2 drops of the EBT indicator. Titrate with EDTA in a burette till the color changes from wine red to blue. Repeat the titration till two successive readings do not differ by more than 0.1 mL. Let the volume of EDTA required be 'a' mL.

Calculations:

Standardization:

$$\text{Molarity of } MgSO_4 = 0.02 \text{ M}$$

$$\text{Molarity of EDTA} = \frac{10 \times 0.02}{a} = M1$$

II) Estimation of total hardness of water.

Make up the given solution to 100 mL in the standard flask by carefully diluting up to the mark using distilled water. Pipette out 10 mL of this solution and then follow the same procedure as in Part- I.

Estimation:

$$\text{Molarity of EDTA} = M1$$

$$\text{Volume of EDTA} = b \text{ mL}$$

Hardness of water in the given solution in terms of

$$\text{CaCO}_3 = \frac{M1 \times b}{10} \times \frac{100}{10} \text{ g} = M1 \times b \text{ grams.} = X \text{ grams.}$$

Results:

1. Molarity of EDTA :M1.....M.
2. Amount of CaCO₃ in the given hard water sample.....Xgm.

Experiment B.

Procedure:

1. Take the EBT solution prepared in deionized water (supplied by the TA), and split it into three portions in the 50 mL beakers.
2. Leave the first beaker as it is.
3. Add three drops of Mg(II) salt solution to the second beaker. Notice the color change.
4. In the third beaker, add Mg(II) salt solution (three drops) followed by five drops of EDTA solution. Notice the sequential color change.
5. Record the final colors of the solutions present in the three beakers.
6. The TA will provide the optical spectra recorded for these three solutions.

Analysis

1. Explain the color change of the EBT solution following the periodic addition of Mg(II) and EDTA. Use the supplied optical spectra to support your hypothesis.

Suggested Reading:

Vogel's Textbook of Quantitative Analysis Revised by G.H.Jeffery, J.Basset, J.Mendham and R.C.Denny, 5th edition, Pages: 309-333

Questions:

1. Why metal-EDTA complex is pale coloured compared to metal-EBT? (Discussion on the conjugation of the ligand on metal complex absorbance properties).
2. Why the metal binding is preferred to EDTA vs EBT? (Discussion on the chelation effect, ligand denticity, and the thermodynamic stability).
3. What is the role of the azo group in EBT? (The role of azo group indifferent varieties of regular dyes can be discussed).

7. Preparation of A Fluorescent Dye And Its Analysis By Thin Layer Chromatography (TLC)

Objectives:

- (A) To prepare a fluorescent dye in microscale using a one pot sequential amide formation – nucleophilic aromatic substitution reactions.
- (B) To understand the principles and application of Thin Layer Chromatography.
- (C) Analysis of TLC using ‘ImageJ’.
- (D) Determination of molecular weight by using TLC-MS.

Introduction:


Chromatography is a technique which is widely used in chemistry and biology for the separation of components of a mixture of compounds. The TLC technique consists of applying the mixture to be separated onto stationary phase (such as a thin layer of silica gel coated on a glass plate or preloaded silica gel on aluminium plate) and allowing a solvent (mobile phase) to flow over the adsorbent.

Once the components are separated, they are detected and visualized using colour reactions or UV source. The detection using colour reactions normally involves spraying the TLC plates with an appropriate reagent to obtain characteristic colours. For example, non-selective visualization can be effected by exposure to iodine vapors, which are absorbed by the organic compounds resulting in brown coloured spots marking the positions of the individual components. Visualization under UV illumination is another common method whereby some compounds can be distinguished by characteristic fluorescent spots.

The most important criterion in qualitative identification is the R_f value, which is defined as


$$R_f = \frac{\text{Distance travelled by the solute}}{\text{Distance travelled by the solvent front}}$$

Fluorophores are the molecules, which can be excited and relax efficiently in a radiative pathway. On absorption of a photon, a molecule in the singlet ground state (S_0) gets excited to the first excited state (S_1). It can relax back to the ground state (S_0) by emission of a photon of lower energy than the incident one (Fluorescence) as depicted below.



In this experiment, we are going to prepare the highly fluorescent dye 4-hexylamino-N-hexyl-1,8-naphthalimide from 4-chloro-1,8-naphthalic anhydride. The reaction can be monitored using TLC with visualization using UV lamp. You would also observe the variation of R_f with the nature of the eluting solvent.

Mass spectrometry



A schematic of the atmospheric solids analysis probe (ASAP).


Advion expression compact mass spectrometer (CMS) and a direct analysis probe, known as the ASAP (atmospheric solids analysis probe). The probe uses the instrument's atmospheric pressure chemical ionization (APCI), eliminating the need for students to carry out any sample preparation before analysis. The APCI ion source with an ASAP sampling probe allows for direct insertion of liquids and solids, making for very quick analyses. Each analysis typically takes 30 s or less.

Preparation of fluorescent dye:

In a reaction vial (8 ml), weigh 5-10 mg of 4-chloro-1,8-naphthalic anhydride and add 1 ml of toluene. The mixture is then stirred with a spin bar on a magnetic stirrer. To this, add 1-hexylamine (500 μ L) by means of a syringe/micropipette. The reaction mixture is then heated and allowed to stir for further 45 minutes under reflux or until all the solid has dissolved. At the end of the heating period, the reaction mixture is allowed to cool. It is then analyzed by using two solvent systems as follows:

1. 20% ethyl acetate in hexane
2. 100% toluene.

The reaction progresses by several readily observable changes during the formation of the final product. The chemical reactions are shown as below:



Following observations can be made during the experiment:

A transient intense orange colour is observed on addition of the amine to the stirred suspension of the anhydride in toluene and the reaction mixture becomes noticeably warmer. With fading of the transient colour, the pale tan of the anhydride gets replaced with the grayish off-white of the acid amide. On heating the reaction mixture a gradual cyclization of the acid amide to the imide takes place to give a yellow solution. When pure, the imide is a yellow solid which exhibits a moderately intense yellow fluorescence. Once the imide is formed, the activating influence of both carbonyl groups is brought to bear on the chlorine. In the presence of the large excess of hexylamine, displacement of the halogen occurs just slightly more slowly than closure of the imide ring to give the bright yellow, intensely fluorescent 4-hexylamine-N-hexyl-1,8-naphthalimide.

On performing a TLC with eluting solvent 20% ethyl acetate in hexane you will observe four to six spots evenly spaced and easily visible, with R_f values falling between 0.8 and 0.3. Using toluene as eluting solvent you will observe four bands with R_f values between 0.5 and 0.05. All the R_f values for the four universally observed spots are collected in observation table.

TLC Procedure:

With a pencil gently mark a base line from the bottom edge of the preloaded TLC plate. The sample should be applied as tiny dots on the base line and well-spaced from each other. Draw a "finish line" a couple of centimetres from the top edge of the silica layer.

The trick in doing a good TLC is the spotting. For good separation (i.e. no streaking or mixing) loading of the sample onto the TLC plate must be neat and compact. Draw a small amount of the sample solution into a capillary, say 1 or 2 cm and lightly touch the capillary to the silica surface so that the sample is soaked up by the silica and forms a tiny spot. It is important to keep the spot small, apply a little at a time. Make about three applications so that there is sufficient amount of the sample loaded to give a distinctly visible coloration in the visualization step. Do not reuse the capillary for another sample to avoid contamination.

After completing the spotting of all the samples, the plate is transferred to a glass chamber containing the solvent to be used as the mobile phase. The solvent level must be a few mm below the starting line. The inside walls of the chamber are lined with filter paper, leaving a gap for viewing the plate. The filter paper is allowed to saturate with the solvent so as to enable the atmosphere in the jar to become saturated with solvent vapour. Allow the solvent to rise along the silica surface and as soon as the solvent front reaches the finish line, remove the plate from the solvent chamber and allow the solvent to dry out. Now capture the image using your smartphone/digital camera as a jpeg file for loading in ImageJ software. Send the pictures you took of the TLC plate to your e-mail or plug in the camera and load the pictures onto the

desktop. Instructions on how to use ImageJ are provided in text at the end of this handout. Calculate relative % of each spot using the following formula:

$$\text{Composition (\%)} \text{ of a particular spot} = \frac{\text{Integrated value (area) of the particular spot}}{\sum \text{Integrated values of all the spots}} \times 100$$

Observation Table:

Solvent	Colour of the spot	R _f	Composition (%) of each spots
20% Ethyl acetate in Hexane			
Toluene			

Procedure for ImageJ software:

ImageJ (free download from <http://imagej.nih.gov/ij/>) is a program that can be used to compare the density of bands or spots. It is most commonly used for biochemistry techniques. This handout assumes that your picture from your TLC plate is a jpeg file.

1. Open the ImageJ program.
2. Open the image file using File > Open in ImageJ. Your TLC plate should be vertical in your image. Rotate it if necessary using Image > Transform.
3. Make sure the rectangle selections tool is clicked.



4. Draw a rectangle around the first lane.
5. After drawing the rectangle around your first lane, press 1. This will leave lane 1 in place, while creating a second rectangle directly on top of it (Figure 1).
6. Use your arrow keys to move the second rectangle over the second lane.
7. Press 3. This will automatically bring up a window that has integrated the lanes for density. This window represents the relative density of the contents of the rectangle over each lane. The rectangles are arranged top to bottom on the profile plot (Figure 2).
A copy of the plate used for ImageJ should be included in your report (Save the image using File > Save As. The image format should be Jpeg).
8. Use the Line tool to draw a line across the baseline for each peak in the TLC lanes. Images will have some background signal, so the peaks likely won't reach down to the baseline of the profile plot. This may make the peak appear to float above the baseline. It is necessary to make sure to close off the peak in order to measure its size.



9. Use the Wand tool to click inside of each of the peaks in the TLC lanes. As you click on a peak, it will be highlighted in yellow. There will also be a window that pops up to give area

	Area
1	20562.865
2	3130.054
3	2370.276
4	17625.137

Figure 2.

The TLC/CMS analysis of the reaction mixture was performed on the Advion **expression** CMS and Plate Express. A solvent composed of 0.1% formic in methanol was used for the elution of the analytes from the TLC plate. The eluted analytes were directed to the CMS for aquisition of the corresponding mass spectra for the reactants and products.

REFERENCES

- (1) Calimente, S.; Strand, S. M.; Chang, S-C.; Lewis, D. E. J. Chem. Ed. 1999, 76, 82-83.
(2) Wagner, A.J.; Miller, S.M.; Naguyen, S.; Lee, G. Y.; Rychnovsky, S.; Link, R.D. J. Chem. Ed. 2014, 91, 716-721.

37

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₂₁**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0691	22b0703	22b0715	22b0727	22b0739	22b0750	22b0761
22b0692	22b0704	22b0716	22b0728	22b0740	22b0751	22b0762
22b0693	22b0705	22b0717	22b0729	22b0741	22b0752	22b0763
22b0694	22b0706	22b0718	22b0730	22b0742	22b0753	22b0764
22b0695	22b0707	22b0719	22b0731	22b0743	22b0754	22b0765
22b0696	22b0708	22b0720	22b0732	22b0744	22b0755	22b0766
22b0697	22b0709	22b0721	22b0733	22b0745	22b0756	22b0767
22b0698	22b0710	22b0722	22b0734	22b0746	22b0757	22b0768
22b0699	22b0711	22b0723	22b0735	22b0747	22b0758	22b0769
22b0700	22b0712	22b0724	22b0736	22b0748	22b0759	22b0770
22b0701	22b0713	22b0725	22b0737	22b0749	22b0760	22b0771
22b0702	22b0714	22b0726	22b0738	-	-	-
-	-	-	-	-	-	-

SCHEDULE FOR DIVISION P₂₁**MONDAY 0930-1230hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
13.03.23	A	B	C	D	E	F	G
20.03.23	G	A	B	C	D	E	F
27.03.23	F	G	A	B	C	D	E
03.04.23	E	F	G	A	B	C	D
10.04.23	D	E	F	G	A	B	C
17.04.23	C	D	E	F	G	A	B
01.05.23	B	C	D	E	F	G	A
08.05.23	Compensation						
22.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₁₅**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0901	22b0914	22b0927	22b0940	22b0953	22b0966	22b0979
22b0902	22b0915	22b0928	22b0941	22b0954	22b0967	22b0980
22b0903	22b0916	22b0929	22b0942	22b0955	22b0968	22b0981
22b0904	22b0917	22b0930	22b0943	22b0956	22b0969	22b0982
22b0905	22b0918	22b0931	22b0944	22b0957	22b0970	22b0983
22b0906	22b0919	22b0932	22b0945	22b0958	22b0971	22b0984
22b0907	22b0920	22b0933	22b0946	22b0959	22b0972	22b0985
22b0908	22b0921	22b0934	22b0947	22b0960	22b0973	22b0986
22b0909	22b0922	22b0935	22b0948	22b0961	22b0974	22b3323
22b0910	22b0923	22b0936	22b0949	22b0962	22b0975	22b3324
22b0911	22b0924	22b0937	22b0950	22b0963	22b0976	22b3325
22b0912	22b0925	22b0938	22b0951	22b0964	22b0977	22b3326
22b0913	22b0926	22b0939	22b0952	22b0965	22b0978	-

SCHEDULE FOR DIVISION P₁₅**MONDAY 1400-1700hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
13.03.23	A	B	C	D	E	F	G
20.03.23	G	A	B	C	D	E	F
27.03.23	F	G	A	B	C	D	E
03.04.23	E	F	G	A	B	C	D
10.04.23	D	E	F	G	A	B	C
17.04.23	C	D	E	F	G	A	B
01.05.23	B	C	D	E	F	G	A
08.05.23	Compensation						
29.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₁₉**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0001	22b0013	22b0024	22b0035	22b0046	22b0057	22b0068
22b0002	22b0014	22b0025	22b0036	22b0047	22b0058	22b0069
22b0003	22b0015	22b0026	22b0037	22b0048	22b0059	22b0070
22b0004	22b0016	22b0027	22b0038	22b0049	22b0060	22b0071
22b0005	22b0017	22b0028	22b0039	22b0050	22b0061	22b0072
22b0006	22b0018	22b0029	22b0040	22b0051	22b0062	22b0073
22b0007	22b0019	22b0030	22b0041	22b0052	22b0063	22b0074
22b0008	22b0020	22b0031	22b0042	22b0053	22b0064	22b0075
22b0009	22b0021	22b0032	22b0043	22b0054	22b0065	22b0076
22b0010	22b0022	22b0033	22b0044	22b0055	22b0066	22b0077
22b0011	22b0023	22b0034	22b0045	22b0056	22b0067	22b0078
22b0012	-	-	-	-	-	-
-	-	-	-	-	-	-

SCHEDULE FOR DIVISION P₁₉**TUESDAY 0830-1130hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
14.03.23	A	B	C	D	E	F	G
21.03.23	G	A	B	C	D	E	F
28.03.23	F	G	A	B	C	D	E
01.04.23	E	F	G	A	B	C	D
11.04.23	D	E	F	G	A	B	C
18.04.23	C	D	E	F	G	A	B
02.05.23	B	C	D	E	F	G	A
09.05.23	Compensation						
23.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₁₃**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0301	22b0314	22b0327	22b0340	22b0353	22b0366	22b0379
22b0302	22b0315	22b0328	22b0341	22b0354	22b0367	22b0380
22b0303	22b0316	22b0329	22b0342	22b0355	22b0368	22b0381
22b0304	22b0317	22b0330	22b0343	22b0356	22b0369	22b0382
22b0305	22b0318	22b0331	22b0344	22b0357	22b0370	22b0383
22b0306	22b0319	22b0332	22b0345	22b0358	22b0371	22b0384
22b0307	22b0320	22b0333	22b0346	22b0359	22b0372	22b0385
22b0308	22b0321	22b0334	22b0347	22b0360	22b0373	22b0386
22b0309	22b0322	22b0335	22b0348	22b0361	22b0374	22b0387
22b0310	22b0323	22b0336	22b0349	22b0362	22b0375	22b0388
22b0311	22b0324	22b0337	22b0350	22b0363	22b0376	22b0389
22b0312	22b0325	22b0338	22b0351	22b0364	22b0377	22b0390
22b0313	22b0326	22b0339	22b0352	22b0365	22b0378	-

SCHEDULE FOR DIVISION P₁₃**TUESDAY 1400-1700hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
14.03.23	A	B	C	D	E	F	G
21.03.23	G	A	B	C	D	E	F
28.03.23	F	G	A	B	C	D	E
01.04.23	E	F	G	A	B	C	D
11.04.23	D	E	F	G	A	B	C
18.04.23	C	D	E	F	G	A	B
02.05.23	B	C	D	E	F	G	A
09.05.23	Compensation						
30.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₂₂**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b1501	22b1513	22b1526	22b1539	22b2705	22b2718	22b2731
22b1502	22b1514	22b1527	22b1540	22b2706	22b2719	22b2732
22b1503	22b1515	22b1528	22b1541	22b2707	22b2720	22b2733
22b1504	22b1516	22b1529	22b1542	22b2708	22b2721	22b2734
22b1505	22b1517	22b1530	22b1543	22b2709	22b2722	22b2735
22b1506	22b1518	22b1531	22b1544	22b2710	22b2723	22b2736
22b1507	22b1519	22b1532	22b1545	22b2711	22b2724	22b2737
22b1508	22b1520	22b1533	22b1546	22b2712	22b2725	22b2738
22b1509	22b1521	22b1534	22b1547	22b2713	22b2726	22b2739
22b1510	22b1522	22b1535	22b2701	22b2714	22b2727	22b2740
22b1511	22b1523	22b1536	22b2702	22b2715	22b2728	22b2741
22b1512	22b1524	22b1537	22b2703	22b2716	22b2729	22b2742
22b1501	22b1525	22b1538	22b2704	22b2717	22b2730	-

SCHEDULE FOR DIVISION P₂₂**THURSDAY 0930-1230hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
16.03.23	A	B	C	D	E	F	G
23.03.23	G	A	B	C	D	E	F
30.03.23	F	G	A	B	C	D	E
06.04.23	E	F	G	A	B	C	D
13.04.23	D	E	F	G	A	B	C
27.04.23	C	D	E	F	G	A	B
04.05.23	B	C	D	E	F	G	A
11.05.23	Compensation						
18.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₁₆**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0987	22b1000	22b1000	22b1026	22b1039	22b1051	22b1063
22b0988	22b1001	22b1001	22b1027	22b1040	22b1052	22b1064
22b0989	22b1002	22b1002	22b1028	22b1041	22b1053	22b1065
22b0990	22b1003	22b1003	22b1029	22b1042	22b1054	22b1066
22b0991	22b1004	22b1004	22b1030	22b1043	22b1055	22b1067
22b0992	22b1005	22b1005	22b1031	22b1044	22b1056	22b1068
22b0993	22b1006	22b1006	22b1032	22b1045	22b1057	22b1069
22b0994	22b1007	22b1007	22b1033	22b1046	22b1058	22b1070
22b0995	22b1008	22b1008	22b1034	22b1047	22b1059	22b1071
22b0996	22b1009	22b1009	22b1035	22b1048	22b1060	22b1072
22b0997	22b1010	22b1010	22b1036	22b1049	22b1061	22b1073
22b0998	22b1011	22b1011	22b1037	22b1050	22b1062	22b1074
22b0999	22b1012	22b1012	22b1038	-	-	-

SCHEDULE FOR DIVISION P₁₆**THURSDAY 1400-1700hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
16.03.23	A	B	C	D	E	F	G
23.03.23	G	A	B	C	D	E	F
30.03.23	F	G	A	B	C	D	E
06.04.23	E	F	G	A	B	C	D
13.04.23	D	E	F	G	A	B	C
27.04.23	C	D	E	F	G	A	B
04.05.23	B	C	D	E	F	G	A
11.05.23	Compensation						
25.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₂₀**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0601	22b0614	22b0627	22b0640	22b0653	22b0666	22b0679
22b0602	22b0615	22b0628	22b0641	22b0654	22b0667	22b0680
22b0603	22b0616	22b0629	22b0642	22b0655	22b0668	22b0681
22b0604	22b0617	22b0630	22b0643	22b0656	22b0669	22b0682
22b0605	22b0618	22b0631	22b0644	22b0657	22b0670	22b0683
22b0606	22b0619	22b0632	22b0645	22b0658	22b0671	22b0684
22b0607	22b0620	22b0633	22b0646	22b0659	22b0672	22b0685
22b0608	22b0621	22b0634	22b0647	22b0660	22b0673	22b0686
22b0609	22b0622	22b0635	22b0648	22b0661	22b0674	22b0687
22b0610	22b0623	22b0636	22b0649	22b0662	22b0675	22b0688
22b0611	22b0624	22b0637	22b0650	22b0663	22b0676	22b0689
22b0612	22b0625	22b0638	22b0651	22b0664	22b0677	22b0690
22b0613	22b0626	22b0639	22b0652	22b0665	22b0678	-

SCHEDULE FOR DIVISION P₂₀**FRIDAY 0930-1230hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
17.03.23	A	B	C	D	E	F	G
24.03.23	G	A	B	C	D	E	F
31.03.23	F	G	A	B	C	D	E
14.04.23	E	F	G	A	B	C	D
15.04.23	D	E	F	G	A	B	C
28.04.23	C	D	E	F	G	A	B
06.05.23	B	C	D	E	F	G	A
12.05.23	Compensation						
19.05.23	Examination						
07.06.23	Quiz						

DEPARTMENT OF CHEMISTRY**CH-117L Spring 2022-2023****DIVISION P₁₄**

Batch A	Batch B	Batch C	Batch D	Batch E	Batch F	Batch G
22b0391	22b0404	22b0417	22b0430	22b0443	22b0456	22b3311
22b0392	22b0405	22b0418	22b0431	22b0444	22b0457	22b3312
22b0393	22b0406	22b0419	22b0432	22b0445	22b0458	22b3313
22b0394	22b0407	22b0420	22b0433	22b0446	22b3301	22b3314
22b0395	22b0408	22b0421	22b0434	22b0447	22b3302	22b3315
22b0396	22b0409	22b0422	22b0435	22b0448	22b3303	22b3316
22b0397	22b0410	22b0423	22b0436	22b0449	22b3304	22b3317
22b0398	22b0411	22b0424	22b0437	22b0450	22b3305	22b3318
22b0399	22b0412	22b0425	22b0438	22b0451	22b3306	22b3319
22b0400	22b0413	22b0426	22b0439	22b0452	22b3307	22b3320
22b0401	22b0414	22b0427	22b0440	22b0453	22b3308	22b3321
22b0402	22b0415	22b0428	22b0441	22b0454	22b3309	22b3322
22b0403	22b0416	22b0429	22b0442	22b0455	22b3310	-

SCHEDULE FOR DIVISION P₁₄**FRIDAY 1400-1700hrs**

Date	Expt. 1	Expt. 2	Expt. 3	Expt. 4	Expt. 5	Expt. 6	Expt. 7
17.03.23	A	B	C	D	E	F	G
24.03.23	G	A	B	C	D	E	F
31.03.23	F	G	A	B	C	D	E
14.04.23	E	F	G	A	B	C	D
15.04.23	D	E	F	G	A	B	C
28.04.23	C	D	E	F	G	A	B
06.05.23	B	C	D	E	F	G	A
12.05.23	Compensation						
26.05.23	Examination						
07.06.23	Quiz						

CH-117L 2022-2023: Spring semester

SCHEDULE

Day/Div	Time(hrs)	March	April	May	Comp. Lab (May)	Lab Exam	Quiz
MON (P ₂₁)	0930-1230	13,20,27	3,10,17	1	8	22-05-23	07-06-23 Venue: LA Halls Time: 8:30 am to 9:15 am
MON (P ₁₅)	1400-1655	13,20,27	3,10,17	1	8	29-05-23	
TUE (P ₁₉)	0830-1130	14,21,28	1*,11,18	2	9	23-05-23	
TUE (P ₁₃)	1400-1655	14,21,28	1*,11,18	2	9	30-05-23	
THURS (P ₂₂)	0930-1230	16,23,30	6,13,27	4	11	18-05-23	
THURS (P ₁₆)	1400-1655	16,23,30	6,13,27	4	11	25-05-23	
Fri (P ₂₀)	0930-1230	17,24,31	14,15*,28	6*	12	19-05-23	
FRI (P ₁₄)	1400-1655	17,24,31	14,15*,28	6*	12	26-05-23	

* Dates marked in asterisks (*) are Saturdays.

The crib session will be held on 21-06-2023.

(Prof. Sanjog Nagarkar)

Course In-charge