

[Dashbo...](#) / [My cour...](#) / [CS23331-DAA-2023-...](#) / [Competitive Program...](#) / [6-Pair with Difference -O\(n\) Time Complexity,O\(1\) Space Com...](#)

Started on	Tuesday, 19 November 2024, 10:16 PM
State	Finished
Completed on	Tuesday, 19 November 2024, 10:19 PM
Time taken	3 mins 38 secs
Marks	1.00/1.00
Grade	4.00 out of 4.00 (100%)

Question 1

Correct

Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2
3  int find_pair_with_difference(int arr[], int n, int k) {
4      int i = 0, j = 1;
5
6      while (i < n && j < n) {
7          int diff = arr[j] - arr[i];
8
9          if (diff == k && i != j) {
10             return 1;
11         } else if (diff < k) {
12             j++;
13         } else {
14             i++;
15         }
16     }
17     return 0;
18 }
19
20 int main() {
21     int n, k;
22     scanf("%d", &n);
23     int arr[n];
24     for (int i = 0; i < n; i++) {
25         scanf("%d", &arr[i]);
26     }
27     scanf("%d", &k);
28     printf("%d\n", find_pair_with_difference(arr, n, k));
29
30     return 0;
31 }
32

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

◀ 5-Pair with Difference-O(n^2)Time Complexity,O(1) Space Complexity

Jump to...

