

Supplemental Material: Movies for “The Elastic Perimeter Disk Model”

Research Note
(Dated: April 28, 2026)

This supplemental material provides six movies of two-disk squeeze simulations at fixed $b = 0.2$, $N = 32$, $\varepsilon_{\text{ref}} = 0.08$, illustrating the squishiness axis as q is varied from area-compressible to nearly incompressible. In all movies: the two EPD disks (blue/orange perimeter nodes) are squeezed between rigid horizontal walls (gray); the color of each node encodes the local contact force magnitude; the title shows current per-particle strain ε_p and instantaneous ν_{meas} .

All movies are provided as animated GIFs in the `movies/` subdirectory.




FIG. 2. **Movie S2** (`movies/S2_q0p1_area_compressible.gif`): Single squeeze simulation at $q = 0.1$ ($q \cdot b = 0.02$, $\nu_{\text{meas}} \approx 0.21$, $\Delta A \approx 5.9\%$). The area penalty is weak; the disk compresses largely in place with little lateral expansion. The flattened top and bottom contact zones are clearly visible; the sides remain nearly straight. Still shown: maximum compression ($\varepsilon_p \approx 0.14$).

MOVIE S1 — FULL SQUISHINESS SWEEP (COMPARISON)




FIG. 1. **Movie S1** (`movies/S1_comparison_q_sweep.gif`): Side-by-side comparison of five q values ($q \in \{0.1, 0.5, 2, 10, 50\}$, left to right) at fixed $b = 0.2$, all compressed to the same wall displacement. The visual transition from strongly flattened area-compressible shapes (left) to laterally bulging nearly-incompressible shapes (right) illustrates the full squishiness axis. Still shown: initial configuration ($\varepsilon_p = 0$).




FIG. 3. **Movie S3** (`movies/S3_q0p5_compressible.gif`): Single squeeze at $q = 0.5$ ($q \cdot b = 0.10$, $\nu_{\text{meas}} \approx 0.42$, $\Delta A \approx 4.3\%$). Moderate area loss; the disk begins to bulge laterally. The transition from predominantly area-compressible to mixed behavior is visible in the rounding of the equatorial cross-section. Still shown: maximum compression.




FIG. 4. **Movie S4** (movies/S4_q2p0_intermediate.gif): Single squeeze at $q = 2.0$ ($q \cdot b = 0.40$, $\nu_{\text{meas}} \approx 0.69$, $\Delta A \approx 2.1\%$). The disk conserves area moderately well; significant lateral bulging is visible. This regime is intermediate between area-compressible and shape-deformable. Still shown: maximum compression.




FIG. 5. **Movie S5** (movies/S5_q10p0_shape_deformable.gif): Single squeeze at $q = 10$ ($q \cdot b = 2.0$, $\nu_{\text{meas}} \approx 0.88$, $\Delta A \approx 0.57\%$). The area is well conserved; the deformation is almost purely shape change. The disk develops a pronounced “peanut” profile under strong compression, with wide flat contact zones and equatorial bulge. Still shown: maximum compression.

MOVIE S2 — AREA-COMPRESSIBLE DISK ($q = 0.1$, $\nu_{\text{meas}} \approx 0.21$)

MOVIE S3 — COMPRESSIBLE DISK ($q = 0.5$, $\nu_{\text{meas}} \approx 0.42$)

MOVIE S4 — INTERMEDIATE DISK ($q = 2.0$, $\nu_{\text{meas}} \approx 0.69$)

MOVIE S5 — SHAPE-DEFORMABLE DISK ($q = 10$, $\nu_{\text{meas}} \approx 0.88$)

MOVIE S6 — NEARLY INCOMPRESSIBLE DISK ($q = 50$, $\nu_{\text{meas}} \approx 0.94$)

MOVIES S7A–S7C — THREE-BODY COLLISIONS: MOMENTUM CONSERVATION

Movies S7a–S7c show three-body disk collision events using `step_rb()`. Two identical EPD disks approach a stationary target from opposite angles; per-frame $\Delta p_x/p_0$




FIG. 6. **Movie S6** (movies/S6_q50p0_nearly_incompressible.gif): Single squeeze at $q = 50$ ($q \cdot b = 10.0$, $\nu_{\text{meas}} \approx 0.94$, $\Delta A \approx 0.12\%$). Near the incompressible limit; the disk conserves area to within 0.1% and deforms entirely by shape change. The lateral bulge is maximal; the contact zone spans a large fraction of the perimeter. This is the “rubber-like” or “jello-like” end of the squishiness axis. Still shown: maximum compression.

and $\Delta p_y/p_0$ are overlaid as text. Three configurations are shown: elastic ($\alpha_d = 0$, $q = 1$), damped ($\alpha_d = 2$, $q = 1$), and rubber-like ($\alpha_d = 0$, $q = 10$). Total linear momentum is conserved to $|\Delta p|/p_0 < 2 \times 10^{-15}$ and angular momentum to $|\Delta L|/L_0 < 1.4 \times 10^{-14}$ in all cases. See `src/validation/rb_extension_benchmarks.py`.

MOVIES S8A–S8D — ELLIPSEPARTICLE: SHAPE GENERALIZATION

Movies S8a–S8d show free-flight collisions of `EllipseParticle` objects (semi-axes $a = 1.5$, $b = 0.8$, $N = 32$ equal-arc-length nodes). Four configurations are covered: E1 (equal ellipses, head-on, no spin), E2 (equal ellipses, oblique impact with initial spin $\omega_0 = \pm 0.8$), E3 (different-size oblique collision with spin), and E4 (same moment of inertia, different mass/density, with spin). Angular momentum is conserved to $|\Delta L|/L_0 \leq 8 \times 10^{-13}$ in all four cases. See `src/validation/ellipse_collision_benchmark.py`.

MOVIE S9 — EMULSION CAPILLARY-WAVE RINGDOWN

Movie S9 shows a single emulsion droplet ($N = 120$, $K_{\text{area}} = 50$, $\kappa = 0.02$, $\alpha = 5.0$) released from a 10% elliptical deformation and allowed to relax. The $n = 2$ capillary-wave mode rings down over ≈ 3 oscillation periods; the exponential amplitude decay gives the measured damping rate $\alpha_{\text{meas}} \approx 2.50 \tau_0^{-1}$, consistent with the near-critical damping target $\alpha_{\text{crit}} = 2\omega_2 \approx 5.46 \tau_0^{-1}$. See `src/validation/emulsion_paper_figures.py`, Benchmark C.

MOVIE S10 — EMULSION T1 TOPOLOGICAL REARRANGEMENT

Movie S10 shows the T1-event benchmark: a center droplet squeezed vertically between two pinned neighbor droplets and two horizontal walls ($d = 1.5 R_0$ gap, $N = 120$, $K_{\text{area}} = 50$). The center droplet elongates vertically, passes through the gap, and snaps horizontally to rest between the two neighbors — the elementary topological rearrangement of an emulsion. Area change during passage is $\Delta A < 2\%$; lateral symmetry is maintained to within $< 1\%$ of R_0 throughout. See `src/validation/emulsion_paper_figures.py`, Benchmark D.

MOVIE S11 — EMULSION DROPLET FALLING UNDER GRAVITY ($\text{Bo} = 0.10$)

Movie S11 shows a single droplet falling freely under gravity at $\text{Bo} = 0.10$ ($N = 120$, $K_{\text{area}} = 50$). The droplet accelerates from rest, reaches terminal-like behavior, and develops a visible oblate deformation: the leading (bottom) edge flattens while the trailing (top) edge rounds, giving a sag ratio $(R_{\text{top}} - R_{\text{bot}})/R_0 \approx 0.065$. Trajectory error relative to ballistic free-fall is $< 3 \times 10^{-4}$ in units of R_0 . See `src/validation/emulsion_paper_figures.py`, Benchmark E.

MOVIE S12 — TERMINAL VELOCITY WITH STOKES DRAG ($\text{Oh} = 0.25$, $g = 0.05$)

Movie S12 shows a single emulsion droplet ($N = 36$, $\kappa = 0.02$, $\text{Oh} = 0.25$, $g = 0.05$) released from rest near the top of a tall box ($L_x = 20$, $L_y = 60$) and falling under gravity with Stokes drag. The right panel shows a real-time speed gauge comparing the measured $|v_{\text{cm},y}|$ to the theoretical terminal velocity $v_t = g/(2\text{Oh}) = 0.10$. The droplet reaches v_t within the first third of the trajectory; subsequent evolution confirms steady terminal-velocity descent. See `papers/summary_of_methods/scripts/test_E_movies.py`, Benchmark E1.

MOVIE S13 — SHEAR FLOW: DRIFT AND ROTATION ($\Gamma = 0.05$, $\text{Oh} = 0.5$)

Movie S13 shows a single emulsion droplet ($N = 36$, $\kappa = 0.02$, $\text{Oh} = 0.5$) in a simple shear background flow $U_x = \Gamma y$, $U_y = 0$ ($\Gamma = 0.05$). The droplet drifts to the right at rate $\dot{x}_{\text{cm}} = \Gamma y_{\text{cm}}$ while rotating clockwise at the predicted Jeffery rate $\omega = -\Gamma/2 = -0.025$. Background shear arrows indicate the local flow direction and magnitude; the right panel shows the instantaneous rotation rate $\omega(t)$ relative to the theoretical value ω_{theory} . See `papers/summary_of_methods/scripts/test_E_movies.py`, Benchmark E3.

