

# The Spatial and Temporal Domains of Modern Ecology

## OR Ecology's Spatio-temporal Domains

### OR Space, Time, and Ecology

Lyndon Estes<sup>\*1,2</sup>, Labeeb Ahmed<sup>3</sup>, Kelly Caylor<sup>2</sup>, Jason Chang<sup>3</sup>,  
Jonathan Choi<sup>4</sup>, Erle Ellis<sup>3</sup>, Paul Elsen<sup>4</sup>, and Tim Treur<sup>4</sup>

<sup>1</sup>Woodrow Wilson School, Princeton University, Princeton, NJ 08544, USA

<sup>2</sup>Civil and Environmental Engineering, Princeton University, Princeton, NJ 08544, USA

<sup>3</sup>Geography and Environmental Systems, University of Maryland Baltimore County,  
Baltimore, MD 21250, USA

<sup>4</sup>Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544, USA

\*To whom correspondence should be addressed; E-mail: [lestes@princeton.edu](mailto:lestes@princeton.edu).

#### An abstract

1 The scales at which ecosystems are observed plays a critical role in shaping our understand-  
2 ing of how they are structured and function (1, 2). Ecological patterns emerge within temporal  
3 and spatial domains that may be coarser or finer than the processes that shape them, which  
4 means that investigation across multiple scales is the *sine qua non* for understanding ecological  
5 phenomena (1). Awareness of the importance of scale has grown rapidly since the 1980s, accel-  
6 erated by the need to understand how changes in the global climate, ocean, and land systems are  
7 affecting everything from individual populations (e.g. cite) to entire biomes (e.g. cite), while  
8 technological advances in areas such as remote sensing and genetics are making it ever-easier  
9 to quantify ecological features across a broad range of scales (2, 3).

Given the importance of multi-scale studies for providing ecological understanding, and the growing ability to undertake them, it is important to rigorously assess whether ecology is becoming a multi-scale discipline. One approach to answering this question is to quantify the spatial and temporal domains within which observations in ecological studies are collected. Observations provide the necessary means for developing and testing the models that explain why ecological patterns vary in time and space (1, 4), thus it stands to reason that the temporal and spatial range of ecological observations, and their density within different portions of those ranges, will shed light on modern ecology's progress towards a holistic, predictive understanding of ecosystems (1, 2). In this study, we quantified the spatio-temporal domain of current ecological studies, using a representative sample of papers published between 2004-2014 in the top 30 ecological journals (by 2014 impact factor) to measure two key dimensions of spatial observation, resolution (grain) and total spatial extent, and their temporal corollaries, sampling interval and total temporal duration. We collected this information from 367 ecological observations (defined here as data collected from non-experimentally manipulated, or "natural" (4), systems) reported within a 148 paper subset of 299 randomly selected articles (1.4% of all papers).

In terms of spatial resolution, here defined as the two-dimensional space in which all measurable features of a natural object were recorded (as opposed to sub-sampled), the majority 63% were collected in plots having resolutions  $<1 \text{ m}^2$ , while 25% were collected within plots of  $1 \text{ m}^2$  up to 1 ha, and the remaining 12% in plots of  $\geq 1 \text{ ha}$  (Fig. 1A). The total spatial extent covered (the number of sampled sites multiplied by the spatial resolution) by 85% of observations was  $<10 \text{ ha}$ , while 31% covered less than  $1 \text{ m}^2$  (Fig. 1B). Only 7% covered an extent  $\geq 10 \text{ ha}$ , with just 1.1% spanning areas  $\geq 10 \text{ million ha}$ .

In the temporal dimensions, 30% of the assessed observations were "once-offs" that were not repeated (Fig. 1C). High frequency observations (ranging between as or more frequent






Figure 2:

## 1. What we did

- (a) We recorded the scales that were actually covered by the observations, rather than potential scales that the collected data might represent—this would require information on autocorrelation length, which is not provided.

Limitation for ocean studies, we did not study third dimension, volume (might be more relevant).

## 2. What we found

(a) Of these, the vast majority (X%) were field-based studies, while Y% involved automated collection of data via instrumentation, and just Z% made use of remote sensing.

### 3. Implications

(a) Although limited sample size, insight into the scales at which science of ecology is making bulk of observations

(b) Despite recognized importance of making observation across scales (*I*), this does not appear to be happening.

(c) Technical capabilities for making observations are not achieving adoption, despite potential of remote sensing to provide multi-scale ecological data (5, 6)

Notes: Wheatley and Johnson (2009) citing arbitrariness of scales in wildlife studies (cited by Chave).

## References and Notes

1. S. A. Levin, *Ecology* **73**, 1943 (1992).

2. J. Chave, *Ecology Letters* **16**, 4 (2013).

3. D. C. Schneider, *BioScience* **51**, 545 (2001).

4. D. Tilman, *Long-term studies in ecology* (Springer, 1989), pp. 136–157.

5. L. D. Estes, A. G. Mwangi, P. R. Reillo, H. H. Shugart, *Animal Conservation* **14**, 521 (2011).

6. L. D. Estes, G. Okin, A. Mwangi, H. H. Shugart, *Remote Sensing of Environment* **112**, 2033 (2008).