

The Spatial and Temporal Domains of Modern Ecology

Lyndon Estes^{*1,2,3}, Paul R. Elsen^{1,4}, Tim Treuer⁵, Labeeb Ahmed⁶,
Kelly Caylor^{1,7}, Jason Chang⁶, Jonathan J. Choi⁵, and Erle Ellis⁶

¹Woodrow Wilson School, Princeton University, Princeton, NJ 08544, USA

²Civil and Environmental Engineering, Princeton University, Princeton, NJ 08544, USA

³Graduate School of Geography, Clark University, Worcester MA 01610, USA

⁴Department of Environmental Science, Policy, and Management,
University of California Berkeley, Berkeley, CA 94720, USA

⁵Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544, USA

⁶Geography and Environmental Systems, University of Maryland Baltimore County,
Baltimore, MD 21250, USA

⁷Earth Research Institute, University of California Santa Barbara, Santa Barbara, CA 93106, USA

*To whom correspondence should be addressed; E-mail: LEstes@clarku.edu.

Abstract

In order to properly understand ecological phenomena, it is necessary to quantify their behavior over the range of spatial and temporal scales at which they manifest. Ecology has been concerned with this need since the early 1990s, and the ability to collect multi-scaled ecological observations has grown rapidly since then. Characterizing the spatial and temporal domains of modern ecological observations can therefore provide important insight into the field's progress in understanding towards a more comprehensive understanding of ecosystem behaviour. To characterize these domains, we conducted a meta-analysis of recent (2004-2014) ecological studies, in which we quantified four primary dimensions of their

reported observations: plot resolution, sampling interval, effective duration (time between start and end of temporal replicates), and effective extent (area enclosed by spatial replicates). We also estimated the *actual* extent and duration, which respectively represent the summed area and time covered by spatial and temporal replicates. **Replace this text with more specific summary of observed scales: Here we show that ecology remains a largely field-based discipline that makes observations within generally narrow spatial and temporal domains, despite the well-established literature on the importance of scale (1–3).**

1 The scales at which ecosystems are observed plays a critical role in shaping our understand-
2 ing of their structure and function (1–3). Ecological patterns emerge from temporal and spatial
3 domains that may be coarser or finer than the processes that shape them, which means that inves-
4 tigation across multiple scales is the *sine qua non* for understanding ecological phenomena (1).
5 This awareness has grown rapidly since the 1980s, accelerated by the need to understand how
6 changes in the global climate, ocean, and land systems are affecting everything from individual
7 populations (4) to entire biomes (5), while technological advances in areas such as remote sens-
8 ing and genetics are making it ever-easier to quantify ecological features across a broad range of
9 scales (2, 6).

10 Given this awareness of the centrality of scale to understanding ecology, and the growing ability
11 to study ecological phenomena across a broad range of spatial and temporal scales, it is important
12 to assess the scales at which current ecological research is conducted. To gain insight into this
13 question, we quantified the spatial and temporal domains of empirical observations that were re-
14 ported in a representative sample of studies published between 2004–2014 in the top 30 ecological
15 journals (by impact factor). Empirical observations provide the necessary means for developing
16 and testing the models that explain why ecological patterns vary in time and space (1, 7), thus
17 it stands to reason that the temporal and spatial distributions of ecological observations may be

indicative of modern ecology’s progress towards achieving a holistic, predictive understanding of ecosystems (*1, 2*).

We characterized observational domains along two key spatial dimensions, resolution (grain) and extent, and their temporal corollaries, interval and duration (Table 1). Here resolution is the area of an individual spatial replicate, or the two-dimensional space in which all measurable features of a natural object(s) were recorded (as opposed to sub-sampled), while extent is the area enclosed by the outer-most spatial replicates, or, if the system or habitat being sampled was distinct from its surrounding matrix (e.g. forest patches in grassland habitats), the summed area of sampled patches (see SI for full definition). Interval refers to the average time elapsed between individual temporal replicates, and duration is the time elapsed between the first and last temporal replicates, or, in the case of temporally unreplicated observations, the estimated time spent collecting the observation (SI). We also calculated two additional metrics, the integrated area of spatial replicates (i.e. resolution multiplied by number of replicates) and the summed observational time of all temporal replicates. We estimated these additional dimensions to evaluate the degree to which the actual scales of ecological observations differ from those they ostensibly represent, and therefore refer to them as the *actual* extent and *actual* duration.

Table 1: The dimensions of ecological observations estimated in this meta-analysis.

Dimension	Description
Resolution	Area (m ²) of an individual spatial replicate (plot)
Extent	Area (ha) encompassed by all spatial replicates
Actual extent	Summed area (ha) of all spatial replicates
Interval	Time elapsed (days) between successive temporal replicates
Duration	Time elapsed (days) between first and last temporal replicates
Actual duration	Summed observational time (days) of all temporal replicates

We calculated these dimensions from 379 discrete observations reported within a 134 paper subset of 348 randomly selected articles (from 42,918 total). An additional 62 papers that were

cited as the source of observations in the selected papers were also reviewed. We confined our analysis to observations made of “natural” (7), or non-experimentally manipulated, systems, given that the inclusion of experiments could have skewed our assessments towards the relatively fine scales at which such studies typically focus.

To account for uncertainty in the estimation of observational dimensions due to 1) unclear methodological description in the reviewed papers, and 2) observer interpretation, we conducted a resampling analysis ($n=1000$) in which scale values were randomly perturbed within the bounds of estimated inter-observer variation (SI). We constructed histograms for each dimension from the mean of the perturbed ensembles, and estimated 95% confidence intervals for each histogram bin (Fig. 1). We constructed kernel density estimates from the full resampled ensemble in order to assess observational distributions within different juxtapositions of the four space-time dimensions (Fig. 2).

In terms of resolution, the majority (67%) of observations were collected in plots of $<1 \text{ m}^2$ resolution, 24% were collected within plots of 1 m^2 up to 1 ha, and the remaining 9% in plots of $\geq 1 \text{ ha}$ (Fig. 1A). The extent of 19% of observations was $<10 \text{ ha}$, 23% covered 10-1,000 ha, 42% 1000-1,000,000 ha, and 16% $>1,000,000 \text{ ha}$ (Fig. 1B).

In the temporal dimensions, 37% of observations were not repeated (Fig. 1C), 17% were repeated at short intervals (sub-second to daily), 20% at daily to monthly intervals, 18% at monthly to yearly intervals, and 8% at yearly to decadal intervals. Duration was one day or less for 31% of sampled observations, while 10% covered one day to one month, 23% lasted one month to one year, 27% covered 1-10 years, and 10% spanned a decade or more (including several paleoecological studies covering centuries to millennia; Fig. 1D).

Juxtaposing these observational dimensions provides further insight into the spatio-temporal distribution of ecological observations and the domains in which they are concentrated (Fig. 2). Contrasting resolution with interval reveals that the majority of temporally replicated observations




Figure 1: Histograms of the resolution (A), extent (B), interval (C), and duration (D) of observations collected from the surveyed ecological studies. Bars represent the average percentages for each bin realized after 1000 perturbed resamples, while grey bars indicate the 95% confidence interval.

(the 37% that were unreplicated were excluded because they lack interval values) had resolutions of 10 cm^2 - 1 m^2 and were revisited at daily to yearly intervals (Fig. 2A). A less dense, oblong concentration of observations bounded on the lower right by monthly to yearly observations at 100 m^2 and on the upper left by near-daily to monthly observations with 1-10 ha resolution is also evident. This lower right to upper left orientation reflects the tradeoff between resolution and interval that is typical of satellite imaging (8), and stands in contrast to the upper right to lower left line that stretches between this concentration and the high frequency (minute-hour intervals), high spatial resolution (0.1 - 100 cm^2) observations. This line demonstrates the opposite tradeoff that occurs with field-based observations, where larger plot sizes demand greater effort that in turn reduces sampling frequency (9).

Contrasting duration and extent (for all observations) reveals two primary domains of observational concentration. The first consists of observations spanning one month to one decade in time and 10-1000 ha in space, while the second is defined by observations of one year to several decades that cover 10,000 to 1,000,000 ha (Fig. 2B). Three other notable, but lesser areas of concentration are also evident, including small area observations (0.1 -1 ha) covering one month to decade, and short duration, temporally unreplicated observations (<1 day) of either 1-10 ha or 10,000-1,000,000 ha.

Comparing the two spatial dimensions against one another (for all observations) shows a primary concentration of observations with 10 cm^2 to 100 m^2 resolution that have extents ranging between slightly over 1,000 to nearly 1,000,000 ha (Fig. 2C). The second-most prominent concentration consists of higher resolution (1 cm^2 - 1 m^2), smaller extent (10-1000 ha) observations, beneath which lies a third and fainter concentration of 1 - $1,000 \text{ cm}^2$ resolution, 1000 m^2 to <10 ha. These three concentrations suggest a tendency for observational extent to increase with resolution, which is a relationship that becomes more pronounced in the (less densely observed) portion of the domain where resolutions $\geq 100 \text{ m}^2$.




Figure 2: Two-dimensional kernel density estimates of observational densities within the domains defined by A) interval and resolution (of temporally replicated observations only), B) duration and extent, C) resolution and extent, and D) interval and duration (of temporally replicated observations). Density estimates were applied to the log-transformed values of each observational dimension, and density estimates are rescaled to represent percentages.

86 A similar tendency for duration to increase with interval was also evident amongst temporally
 87 replicated observations (Fig. 2D), where the majority of observations were repeated at daily to
 88 decadal intervals and spanned ≥ 1 month to <100 years. The orientation of this concentration

shows that observations lasting one year to one decade tend to have corresponding intervals, while those lasting one month to one year have daily to monthly intervals. This suggests that most long-duration observations have just a single temporal replicate. The low densities of observations having sub-daily intervals shows that relatively few high frequency, long duration ecological measurements are undertaken.

To provide further insight into the spatio-temporal domains of ecological observations, we also evaluated the differences in scale between extent and actual extent and between duration and actual duration. To make these comparisons, we calculated the magnitudes of difference (decades) between each dimension and its actual equivalent, and examined how these varied as a function of the scale of the actual measurement (Fig. 3). The majority (80%) of the assessed observations had actual extents of ≤ 1 ha, which on average was 4 to nearly 8 orders of magnitude smaller than the quantified extent (Fig 3.A). Actual extent converged with extent above 100,000 ha, but this applies to just 5% of observations. The actual duration of 65% of observations is < 1 day, which is on average 3 to nearly 9 orders of magnitude shorter than the time span covered by temporal replicates (Fig. 3B). The two duration measures were effectively the same for the 16% of observations exceeding one month of actual duration.

Observational methods

We classified the method used to collect each observation into several broad categories, which were field methods (manual *in situ* data collection), automated (*in situ*) sensing, remote sensing, other geographic data, and paleo-reconstruction approaches. Field methods were used for 80% of observations, automated sensing for 12.4%, remote sensing for 6.3%, and paleo-reconstruction and other geographic data each for less than 1%. Using linear regression (weighted by the number of observations per publication year) to assess whether the relative frequency of observing methods changed during the 10 year study period, the use of remote sensing appeared to increase by 1.3% per year from 2004-2014 ($R^2 = 0.25$, $p < 0.12$), and field methods declined by the same percentage




Figure 3: The difference between *actual* extent (the summed area of spatial replicates) and extent (A) and *actual* duration (the summed sampling duration across temporal replicates) and duration (B). Differences are expressed as decades, or how many orders of magnitude greater the extent or duration is then the actual extent or duration, and are summarized (as box plots, with circle in box representing the mean and line the median) in bins representing different levels of actual extent/duration. The percentages of observations falling within each bin are indicated by the color of the inter-quartile and the numeric value above the upper whisker.

($R^2 = 0.1$, $p < 0.18$), although both slopes failed to meet the customary threshold for statistical significance. Automated sensing methods showed no trend (SI).

Potential biases and uncertainties in quantifying observational scales

There were several potential methodological aspects that could have influenced our assessment of ecology's spatial and temporal domains. The first stems from our finding that many studies did not precisely report observational scales, which meant that we had to estimate, rather than simply record, these values for most observations (specifically, in 63, 60, and 67% of cases for

122 resolution, extent, and actual extent, and 36%, 64%, and 83% of cases for interval, duration,
123 and effective duration). The inevitable errors caused by estimation may have biased our overall
124 findings. However, we attempted to quantify this error by assessing inter-rater disagreement and
125 resampling techniques. The resulting confidence intervals (Fig. 1) suggest that it was unlikely that
126 estimation errors unduly influenced our findings.

127 Related to estimation errors are the rules we used to assess observational scales as well as our
128 selection criteria....

129 Finally, because our review did not include papers beyond 2014, the omission of studies from
130 the most recent years could have introduced bias into our domain estimates. Indeed, if the trend
131 towards increasing use of remote sensing between 2004-2014 was not spurious, we can project that
132 a repeated study applied to papers published between 2004-2017 would find that remote sensing
133 was used for 7.7% of observations (a 22% increase), which would increase the mean effective ex-
134 tent by 17.4% (95% CI = -1.3-67%), or 0.07 orders of magnitude, above that observed here (see SI
135 for details of calculation). Further support for such trend can be found by averaging dimensions by
136 publication year (SI), which reveals that observation extent increased by 0.253 orders of magnitude
137 per year between 2004-2014 ($R^2 = 0.25$, $p < 0.07$). This somewhat clearer trend also suggests that
138 including more recent studies would show observational extent to be somewhat larger, although in
139 this case by a more modest 5.5% (0.02 orders of magnitude).

140 *Insights into the scales of modern ecology*

141
142 the continued dominance of field-based research, and the limited use of methods that allow
143 larger areas to be comprehensively observed, such as remote sensing.

144 satellite observations typically have lower information content than field measurements for
145 a given location, and in many cases only provide proxy measures for the ecological features of
146 interest, such as forest understorey structure (26), thereby making ecologists less inclined to use

the technology (21).

Adapt and tone down, retain these points: Furthermore, the unclear documenting of observational scales implies that scale is not a primary concern in much ecological research (2, 18). (Scale is not a concern that cuts across the entire discipline of ecology). Narrowness and poor documentation (a tendency that is also evident in the geographical sciences (19)) Ecological understanding drawn from many of these observations may have limited generalizability (3, 18, 19), a concern that has been previously noted due to ecology's geographical bias towards anthropogenically undisturbed and temperate ecosystems (20).

Our results provide valuable insight into the spatial and temporal domains being addressed by modern ecological research. Our results show that most observations are collected at small spatial scales, are either unrepeated or relatively infrequent (≥ 1 month interval), and in aggregate cover relatively narrow periods of time (≤ 1 month). Very little research is conducted at high spatial and temporal resolutions over large areas or for long time periods, indicating that, despite the well-established understanding of the importance of multi-scale assessments for understanding ecological patterns and processes (1, 3), efforts focused on larger scales are still relatively sparse within the discipline (1, 3). Furthermore, the unclear documenting of observational scales implies that scale is not a primary concern in much ecological research (2, 18). Taken together, this narrowness and poor documentation (a tendency that is also evident in the geographical sciences (19)) suggests that the ecological understanding drawn from many of these observations may have limited generalizability (3, 18, 19), a concern that has been previously noted due to ecology's geographical bias towards anthropogenically undisturbed and temperate ecosystems (20).

The generally small spatial scales of observation is a consequence of the continued dominance of field-based research, and the limited use of methods that allow larger areas to be comprehensively observed, such as remote sensing. Despite early and repeated calls for ecologists to use remote sensing because it provides a synoptic view that field measurements cannot (21–23), and

subsequent demonstration of its importance for multi-scale studies (24, 25), our results indicate this method has not yet been widely adopted in ecological research. Two reasons lie behind this slow uptake. First, remote sensing can be a challenging method for ecologists to learn, many of whom may not have access to appropriate training (23). Second, satellite observations typically have lower information content than field measurements for a given location, and in many cases only provide proxy measures for the ecological features of interest, such as forest understorey structure (26), thereby making ecologists less inclined to use the technology (21).

In contrast to remote sensing, ecologists have made broader use of technologies that increase the temporal resolution of observations. Automated sensors were used to record 12% of all observations, and accounted for most very high frequency measurements (intervals ≤ 1 hour). As with spatial data, finely resolved temporal data can be aggregated to facilitate multi-scale analyses (and many of the reviewed studies that used automated sensing aggregated the resulting observations before analyzing them), whereas longer-interval data (e.g. annual biomass accumulation) cannot be disaggregated into shorter interval measurements (e.g. weekly biomass accumulation) without interpolation, which turns data into modeled, rather than direct, observations.

In the coming years, rapid technological advances should increase the concentration of ecological observations in currently under-represented domains. The growing numbers of high-resolution satellite sensors, together with new analytical platforms that provide free access to large volumes of pre-processed data and computational power (27), will lower technical barriers that have so far prevented ecologists from adopting this observational technology (23). Similarly, the advent of unmanned systems offers the ability to measure ecological features at high spatial and temporal frequencies over large areas (28), which were scales that were previously impractical to access. The ever-falling cost of sensor technology and the ubiquity of cell phones also means that ecologists, together with a growing army of citizen-scientists, have the unprecedented ability to make spatially dense, high frequency observations over large areas (29–32). A greater attentiveness to

scale in general, including more meticulous documentation of observed dimensions, may help to facilitate the spread of ecological research to sparsely studied scales, while improving transferability of knowledge within the discipline.

References

1. Levin, S. A. The problem of pattern and scale in ecology. *Ecology* **73**, 1943–1967 (1992).
2. Chave, J. The problem of pattern and scale in ecology: What have we learned in 20 years? *Ecology Letters* **16**, 4–16 (2013).
3. Wiens, J. A. Spatial scaling in ecology. *Functional Ecology* **3**, 385–397 (1989). PT: J.
4. Tingley, M. W., Koo, M. S., Moritz, C., Rush, A. C. & Beissinger, S. R. The push and pull of climate change causes heterogeneous shifts in avian elevational ranges. *Global Change Biology* **18**, 3279–3290 (2012).
5. Xiao, J. F. & Moody, A. Photosynthetic activity of US biomes: Responses to the spatial variability and seasonality of precipitation and temperature. *Global Change Biology* **10**, 437–451 (2004).
6. Schneider, D. C. The Rise of the Concept of Scale in Ecology The concept of scale is evolving from verbal expression to quantitative expression. *BioScience* **51**, 545–553 (2001).
7. Tilman, D. Ecological experimentation: Strengths and conceptual problems. In *Long-Term Studies in Ecology*, 136–157 (Springer, 1989).
8. Estes, L. D. *et al.* A platform for crowdsourcing the creation of representative, accurate land-cover maps. *Environmental Modelling & Software* **80**, 41–53 (2016).

- 217 9. Kareiva, P. & Andersen, M. Spatial aspects of species interactions: The wedding of models
218 and experiments. In *Community Ecology*, 35–50 (Springer, 1988).
- 219 10. Underwood, A. J. *Experiments in Ecology: Their Logical Design and Interpretation Using*
220 *Analysis of Variance* (Cambridge University Press, 1997).
- 221 11. Palmer, M. W. & White, P. S. Scale dependence and the species-area relationship. *American*
222 *Naturalist* 717–740 (1994).
- 223 12. Cao, Y., Williams, D. D. & Larsen, D. P. Comparison of Ecological Communities: The
224 Problem of Sample Representativeness. *Ecological Monographs* **72**, 41–56 (2002).
- 225 13. Legendre, P. Spatial autocorrelation - trouble or new paradigm? *Ecology* **74**, 1659–1673
226 (1993).
- 227 14. Hansen, M. C. *et al.* High-resolution global maps of 21st-century forest cover change. *Science*
228 **342**, 850–853 (2013).
- 229 15. Roy, D., Jin, Y., Lewis, P. & Justice, C. Prototyping a global algorithm for systematic fire-
230 affected area mapping using MODIS time series data. *Remote Sensing of Environment* **97**,
231 137–162 (2005).
- 232 16. Jones, B. M. *et al.* Fire Behavior, Weather, and Burn Severity of the 2007 Anaktuvuk River
233 Tundra Fire, North Slope, Alaska. *Arctic, Antarctic, and Alpine Research* **41**, 309–316 (2009).
- 234 17. Clements, C. B. *et al.* Observing the dynamics of wildland grass fires: FireFlux-a field vali-
235 dation experiment. *Bulletin of the American Meteorological Society* **39**, 1369–1382 (2007).
- 236 18. Wheatley, M. & Johnson, C. Factors limiting our understanding of ecological scale. *Ecological*
237 *Complexity* **6**, 150–159 (2009).

- 238 19. Margulies, J. D., Magliocca, N. R., Schmill, M. D. & Ellis, E. C. Ambiguous Geographies:
239 Connecting Case Study Knowledge with Global Change Science. *Annals of the American*
240 *Association of Geographers* **106**, 572–596 (2016).
- 241 20. Martin, L. J., Blossey, B. & Ellis, E. Mapping where ecologists work: Biases in the global
242 distribution of terrestrial ecological observations. *Frontiers in Ecology and the Environment*
243 **10**, 195–201 (2012).
- 244 21. Turner, W. *et al.* Remote sensing for biodiversity science and conservation. *Trends in Ecology*
245 *& Evolution* **18**, 306–314 (2003).
- 246 22. Kerr, J. T. & Ostrovsky, M. From space to species: Ecological applications for remote sensing.
247 *Trends in Ecology & Evolution* **18**, 299–305 (2003). TY - JOUR.
- 248 23. Pettorelli, N. *et al.* Satellite remote sensing for applied ecologists: Opportunities and chal-
249 lenges. *Journal of Applied Ecology* **51**, 839–848 (2014).
- 250 24. Estes, L. D., Okin, G., Mwangi, A. & Shugart, H. H. Habitat selection by a rare forest antelope:
251 A multi-scale approach combining field data and imagery from three sensors. *Remote Sensing*
252 *of Environment* **112**, 2033–2050 (2008).
- 253 25. Estes, L. D., Mwangi, A. G., Reillo, P. R. & Shugart, H. H. Predictive distribution modeling
254 with enhanced remote sensing and multiple validation techniques to support mountain bongo
255 antelope recovery. *Animal Conservation* **14**, 521–532 (2011).
- 256 26. Estes, L., Reillo, P., Mwangi, A., Okin, G. & Shugart, H. Remote sensing of structural com-
257 plexity indices for habitat and species distribution modeling. *Remote Sensing of Environment*
258 **114**, 792–804 (2010).

- 259 27. Google Earth Engine Team. Google Earth Engine: A planetary-scale geo-spatial analysis
260 platform (2015).
- 261 28. Anderson, K. & Gaston, K. J. Lightweight unmanned aerial vehicles will revolutionize spatial
262 ecology. *Frontiers in Ecology and the Environment* **11**, 138–146 (2013).
- 263 29. Wolf, A., Falusi, J., Caylor, K., Sheffield, J. & Wood, E. A GSM-based surface meteorology
264 network in service of improved African hydrological data assimilation and drought forecasting
265 (San Francisco, 2012).
- 266 30. Collins, S. L. *et al.* New opportunities in ecological sensing using wireless sensor networks.
267 *Frontiers in Ecology and the Environment* **4**, 402–407 (2006).
- 268 31. Porter, J. *et al.* Wireless Sensor Networks for Ecology. *BioScience* **55**, 561–572 (2005).
- 269 32. Dickinson, J. L. *et al.* The current state of citizen science as a tool for ecological research and
270 public engagement. *Frontiers in Ecology and the Environment* **10**, 291–297 (2012).

271 **Acknowledgements** This work was supported by funds from the Princeton Environmental Institute
272 Grand Challenges program and the NASA New Investigator Program (NNX15AC64G). Erle Ellis,
273 Jason Chang, and Labeeb Ahmed of the GLOBE Project (<http://globe.umbc.edu>) were supported
274 by the U.S. National Science Foundation (1125210).