

A Degenerate Mixture of $^3\text{He}^*$ and $^4\text{He}^*$ with 3D single particle resolution

K. F. Thomas, S. Li, Y. Wang, B. M. Henson, S. S. Hodgman, A. G. Truscott




Australian
National
University


Department of Quantum Science and Technology, Research School of Physics,
Australian National University, Canberra, Australia

Ultracold $^3\text{He}^* - ^4\text{He}^*$


Cooling and trapping transition



Apparatus



Detection




• Apply a field gradient (along z-axis) to separate isotopes on detector (in arrival time)

• Below critical temperature ($\sim 1\text{ mK}$)

^3He = Degenerate Fermi Gas (DFG): all energy states below Fermi energy have an occupancy of one, and zero otherwise.

^4He = Bose-Einstein Condensate (BEC): all atoms in coherent ground state (equivalent of a laser)



Bose-Fermi Collision

Defintion A collision halo is spherical shell of correlated atom pairs generated by colliding BECs. We can create a collision halo between ^3He and ^4He , and hence generate a mass entangled state.

Generation

1. Bragg diffraction splits BEC into different momentum modes
2. As the BEC and DFG separate Spontaneous s-wave collisions between atom pairs
3. Once the BEC and DFG are fully separated we have a spherical shell of entangled pairs of atoms with equal and opposite momenta.
4. The exact state of two $|\Psi\rangle = \sqrt{1-\mu^2} \sum_{n=0}^{\infty} \mu^n |n\rangle_{\mathbf{k}} |n\rangle_{-\mathbf{k}}$ diametrically opposed modes is a $|\Psi\rangle \approx \frac{1}{\sqrt{2}} |^4\text{He}, ^3\text{He}\rangle + |^3\text{He}, ^4\text{He}\rangle$ two-mode squeezed state.

Correlations

- (a) fermion-boson nonlocal correlation
 - (b) fermion-fermion local antibunching (Pauli exclusion principle)
 - (c) boson-boson local antibunching due to the correlation with fermionic partner.
- Fig. 4. Diagram of various correlation measurements between representative fermionic (F, squares) and bosonic (B, circles) atoms on the s-wave scattering sphere.

Possible uses

- Investigate nonequilibrium properties of bose-fermi mixtures
- Measure correlations between fermion-boson scattering events
- Observe novel phenomena including: quantum phase transitions; and pairing of fermions and formation of composite particles.

Fermionic Anti-Bunching

Information about manybody wavefunction is encoded in the second order correlation function

$$g^{(2)}(\tau) = \frac{\langle N(t) \times N(t+\tau) \rangle}{\langle N(t) \rangle \langle N(t+\tau) \rangle} \text{ averaged over all } t$$

and more generally $g^{(n)}(\tau_1, \dots, \tau_{n-1}) = \frac{\langle N(t) \times N(t+\tau_1) \times \dots \times N(t+\tau_{n-1}) \rangle}{\langle N(t) \rangle \langle N(t+\tau_1) \rangle \dots \langle N(t+\tau_{n-1}) \rangle}$

Bosons bunch and thus have a $g^{(2)}$ peak above 1

Fermions anti-bunch and thus have a $g^{(2)}$ peak below 1, i.e. anti-correlated.

