

King Fahad Central Hospital

STROKE UNIT

POLICY AND PROCEDURE MANUAL

*Aligned with American Heart Association / American Stroke Association (AHA/ASA)
Scientific Guidelines and The Joint Commission Stroke Certification Standards*

Manual Number: SU-MANUAL-01

Effective Date: [Effective Date] Review Date: [Review Date]

Approved By: Medical Director / Stroke Program Medical Director

Applies To: Hospital Wide

Table of Contents

Table of Contents.....	2
Manual Introduction and Scope	6
Manual Contents	6
SU-001 Stroke Program Organization, Governance and Staffing	7
1 .Purpose.....	7
2 .Scope	7
3 .Definitions and Abbreviations	7
4 .Policy	7
5 .Multidisciplinary Stroke Team.....	8
6 .Staffing and Competency Requirements.....	8
7 .Committee Structure and Performance Review	8
8 .References.....	8
SU-002 Stroke Code Activation — Pre-Hospital and In-Hospital	10
1 .Purpose.....	10
2 .Definitions and Abbreviations	10
3 .Policy	10
4 .Procedure — In-Hospital Stroke Code Activation	10
5 .Procedure — Pre-Hospital Notification	11
6 .Documentation.....	11
7 .References.....	11
8 .Attachments	12
SU-003 Acute Ischemic Stroke Management — IV Thrombolysis and Mechanical Thrombectomy	13
1 .Purpose.....	13
2 .Definitions and Abbreviations	13
3 .Policy	13
4 .Procedure — Initial Assessment (0–6 Hours from Onset/LKW).....	13
5 .Procedure — IV Thrombolytic Administration	14
6 .Procedure — Mechanical Thrombectomy	14
7 .Procedure — Extended Window Evaluation (6–24 Hours)	14
8 .General Supportive Care (All Acute Ischemic Stroke Patients)	15
9 .Admission Criteria	15
10 .References.....	15
11 .Attachments	16
SU-004 Spontaneous Intracerebral and Subarachnoid Hemorrhage Management.....	17
1 .Purpose.....	17
2 .Definitions	17
3 .Policy	17
4 .Procedure — Intracerebral Hemorrhage	17
5 .Procedure — Subarachnoid Hemorrhage	18
6 .Admission and Escalation Criteria	18
7 .References.....	18
8 .Attachments	19
SU-005 Transient Ischemic Attack (TIA) Evaluation and Management	20
1 .Purpose.....	20
2 .Definitions	20
3 .Policy	20
4 .Procedure	20
5 .Admission Criteria	21
6 .References.....	21
SU-006 NIHSS Assessment, Neurological Monitoring and Dysphagia/Swallow Screening.....	22
1 .Purpose.....	22

2 .Policy — NIHSS	22
3 .Policy — Dysphagia / Swallow Screening	22
4 .Documentation.....	22
5 .References.....	23
6 .Attachments.....	23
SU-007 Stroke Rehabilitation and Early Mobilization.....	24
1 .Purpose.....	24
2 .Policy	24
3 .Procedure — Rehabilitation Assessment	24
4 .Procedure — Therapeutic Interventions.....	24
5 .Discharge Rehabilitation Planning.....	24
6 .References.....	25
SU-008 Discharge Planning, Secondary Prevention and Patient/Family Education	26
1 .Purpose.....	26
2 .Policy	26
3 .Required Patient/Family Education Topics	26
4 .Discharge Criteria from the Stroke Unit	26
5 .Documentation.....	27
6 .References.....	27
7 .Attachments.....	27
SU-009 Stroke Performance Improvement, Data Registry and Peer Review	28
1 .Purpose.....	28
2 .Policy	28
3 .Procedure	28
4 .References.....	29
5 .Attachments.....	29
SU-010 Code Stroke Team — Roles, Responsibilities and Activation Roster.....	30
1 .Purpose.....	30
2 .Scope	30
3 .Definitions and Abbreviations.....	30
4 .Policy	30
5 .Team Roles and Responsibilities	30
6 .Activation Roster and Contact Method.....	31
7 .References.....	31
8 .Attachments.....	31
SU-011 Acute Stroke Immediate Management Pathway (0–24 Hour Clinical Pathway).....	32
1 .Purpose.....	32
2 .Scope	32
3 .Policy	32
4 .Immediate Management Pathway.....	32
5 .Escalation Triggers Throughout the Pathway.....	33
6 .References.....	33
7 .Attachments.....	33
SU-012 Blood Pressure Management in Acute Ischemic and Hemorrhagic Stroke.....	34
1 .Purpose.....	34
2 .Scope	34
3 .Policy — General Principles.....	34
4 .Algorithm — Acute Ischemic Stroke.....	34
5 .Algorithm — Acute Hemorrhagic Stroke (ICH / SAH)	34
6 .Documentation.....	35
7 .References.....	35
8 .Attachments.....	35
SU-013 Critical Actions — Suspected Intracranial Hemorrhage / Systemic Bleeding After IV Thrombolytic (rtPA) Administration.....	36

1 .Purpose.....	36
2 .Scope	36
3 .Definitions	36
4 .Policy	36
5 .Critical Action Sequence.....	36
6 .Documentation.....	37
7 .References.....	37
8 .Attachments	37
SU-014 Mechanical Thrombectomy Policy	38
1 .Purpose.....	38
2 .Scope	38
3 .Definitions and Abbreviations.....	38
4 .Policy — Patient Selection.....	38
5 .Policy — Notification and Transfer Workflow	38
6 .Policy — Procedural and Post-Procedure Care	39
7 .Admission Criteria Post-Thrombectomy	39
8 .References.....	39
9 .Attachments	39
SU-015 Nursing and Staff Stroke Education Program.....	40
1 .Purpose.....	40
2 .Scope	40
3 .Policy	40
4 .Core Orientation Curriculum	40
5 .Ongoing / Annual Education.....	40
6 .Documentation.....	41
7 .References.....	41
8 .Attachments	41
SU-016 Patient and Caregiver Stroke Education Program.....	42
1 .Purpose.....	42
2 .Scope	42
3 .Policy	42
4 .Core Patient/Caregiver Education Topics.....	42
5 .Delivery Methods	43
6 .Documentation.....	43
7 .References.....	43
8 .Attachments	43
SU-017 Intensive Care Unit Admission Criteria for Stroke Patients	44
1 .Purpose.....	44
2 .Scope	44
3 .Policy — Mandatory ICU Admission Criteria	44
4 .Policy — Stroke Unit (Non-ICU Monitored) Admission Criteria	44
5 .Policy — Continuous Reassessment and Escalation.....	44
6 .Documentation.....	45
7 .References.....	45
8 .Attachments	45
Appendix A — NIH Stroke Scale (NIHSS): Full Item Scoring Reference.....	46
Scoring Summary and Severity Bands	46
Appendix B — IV Thrombolytic (Alteplase/Tenecteplase) Inclusion/Exclusion Checklist	47
Section 1 — Inclusion Criteria (all must be YES).....	47
Section 2 — Exclusion Criteria (any YES excludes thrombolysis unless overridden by documented physician judgment)	47
Section 3 — Extended-Window (3–4.5 hour) Additional Exclusion Considerations	47
Section 4 — Physician Attestation.....	48
Appendix C — Stroke Code / Acute Stroke Flow Sheet, Illustrated.....	49

Illustrated Timeline.....	49
Team Roster Recorded This Activation	50
Appendix D — Blood Pressure Management Algorithm: Acute Ischemic Stroke (Illustrated).....	51
Appendix E — Blood Pressure Management Algorithm: Acute Hemorrhagic Stroke (Illustrated).....	52
Appendix F — Critical Action Card: Suspected ICH / Systemic Bleeding After rtPA (Illustrated).....	53
Quick-Reference Contact List	53
Manual Approval	54

Manual Introduction and Scope

This manual constitutes the official Policy and Procedure framework for the Stroke Unit and Stroke Program of King Fahad Central Hospital. It is developed to ensure consistent, evidence-based, multidisciplinary stroke care across the continuum — pre-hospital notification, emergency department triage, acute treatment, inpatient management, rehabilitation, and secondary prevention — in alignment with the scientific guidelines of the American Heart Association / American Stroke Association (AHA/ASA) and the certification standards of The Joint Commission (TJC) Stroke Certification programs.

Each policy in this manual follows a standardized format — Purpose, Definitions, Policy, Procedure, References, and Attachments — with a policy identification header (policy number, effective date, review date, approving authority, and applicability) and a footer identifying the applicable accreditation standard and page numbering, to support survey readiness and document control.

This manual is reviewed and approved by the Stroke Committee and Medical Director at minimum every two years, or sooner if clinical guidelines, regulatory requirements, or internal performance data indicate a need for revision. All bracketed fields (e.g., [Effective Date], [Review Date]) are to be completed by the organization prior to implementation.

Manual Contents

SU-001 — Stroke Program Organization, Governance and Staffing
SU-002 — Stroke Code Activation — Pre-Hospital and In-Hospital
SU-003 — Acute Ischemic Stroke Management — IV Thrombolysis and Mechanical Thrombectomy
SU-004 — Spontaneous Intracerebral and Subarachnoid Hemorrhage Management
SU-005 — Transient Ischemic Attack (TIA) Evaluation and Management
SU-006 — NIHSS Assessment, Neurological Monitoring and Dysphagia/Swallow Screening
SU-007 — Stroke Rehabilitation and Early Mobilization
SU-008 — Discharge Planning, Secondary Prevention and Patient/Family Education
SU-009 — Stroke Performance Improvement, Data Registry and Peer Review
SU-010 — Code Stroke Team — Roles, Responsibilities and Activation Roster
SU-011 — Acute Stroke Immediate Management Pathway (0–24 Hour Clinical Pathway)
SU-012 — Blood Pressure Management in Acute Ischemic and Hemorrhagic Stroke
SU-013 — Critical Actions — Suspected ICH / Systemic Bleeding After IV Thrombolytic (rtPA)
SU-014 — Mechanical Thrombectomy Policy
SU-015 — Nursing and Staff Stroke Education Program
SU-016 — Patient and Caregiver Stroke Education Program
SU-017 — Intensive Care Unit Admission Criteria for Stroke Patients
Appendix A — NIH Stroke Scale (NIHSS): Full Item Scoring Reference
Appendix B — IV Thrombolytic Inclusion/Exclusion Checklist
Appendix C — Stroke Code / Acute Stroke Flow Sheet, Illustrated
Appendix D — Blood Pressure Management Algorithm: Acute Ischemic Stroke, Illustrated
Appendix E — Blood Pressure Management Algorithm: Acute Hemorrhagic Stroke, Illustrated
Appendix F — Critical Action Card: Suspected ICH / Systemic Bleeding After rtPA, Illustrated

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE PROGRAM ORGANIZATION, GOVERNANCE AND STAFFING

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-001	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Program Medical Director		
APPLIES TO	Hospital Wide — Emergency Department, Stroke Unit, ICU, Radiology, Laboratory, Pharmacy, Rehabilitation		

SU-001 Stroke Program Organization, Governance and Staffing

1. Purpose

To define the organizational structure, governance, staffing, and multidisciplinary team requirements of the Stroke Program, and to establish the administrative framework required to support consistent, evidence-based stroke care in compliance with American Heart Association/American Stroke Association (AHA/ASA) guidelines and The Joint Commission (TJC) Primary Stroke Center / Comprehensive Stroke Center certification standards.

2. Scope

This policy applies to all departments and personnel involved in the identification, triage, treatment, admission, and rehabilitation of stroke and transient ischemic attack (TIA) patients, including the Emergency Department (ED), Stroke Unit, Intensive Care Unit (ICU), Neurology, Neurosurgery, Interventional Radiology, Diagnostic Imaging, Laboratory, Pharmacy, Nursing, and Rehabilitation Services (Physical Therapy, Occupational Therapy, Speech-Language Pathology).

3. Definitions and Abbreviations

- 3.1 AHA/ASA: American Heart Association / American Stroke Association.
- 3.2 TJC: The Joint Commission.
- 3.3 PSC: Primary Stroke Center.
- 3.4 CSC: Comprehensive Stroke Center.
- 3.5 GWTG-Stroke: Get with the Guidelines®–Stroke, the AHA/ASA hospital-based quality improvement registry and program.
- 3.6 Stroke Medical Director: A physician (neurologist or physician with equivalent stroke expertise) responsible for the clinical oversight of the Stroke Program.
- 3.7 Stroke Coordinator: A registered nurse or allied health professional responsible for day-to-day program coordination, data abstraction, staff education, and performance improvement activities.
- 3.8 Multidisciplinary Stroke Team: The group of clinical disciplines required for comprehensive stroke care (see Section 5).
- 3.9 Interprofessional Committee (IPC): the multidisciplinary governance body — synonymous with the Stroke Committee referenced throughout this manual — responsible for overseeing the Stroke Program's mission, goals, scope, and organizational structure.
- 3.10 Telehealth Champion: A clinician with telehealth experience/expertise who identifies telehealth opportunities and drives adoption of telestroke and related telehealth services for the Stroke Program.
- 3.11 ASRC: Acute Stroke Ready Center — a facility providing initial stroke identification, stabilization, and, where indicated, transfer to a higher level of care.

4. Policy

- 4.1 The organization maintains a designated Stroke Program with defined governance, medical leadership, and administrative support, consistent with TJC stroke certification requirements and AHA/ASA scientific guidelines.
- 4.2 The Stroke Program operates under a Stroke Committee that meets at minimum quarterly to review performance data, complications, protocol deviations, and opportunities for improvement.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE PROGRAM ORGANIZATION, GOVERNANCE AND STAFFING

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-001	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Program Medical Director		
APPLIES TO	Hospital Wide — Emergency Department, Stroke Unit, ICU, Radiology, Laboratory, Pharmacy, Rehabilitation		

4.3 A Stroke Medical Director and a Stroke Program Coordinator are formally appointed in writing, with defined roles, responsibilities, and reporting lines to hospital administration and the Medical Staff.

4.4 All staff involved in stroke care receive stroke-specific orientation and ongoing education, with documented annual competency verification.

4.5 The organization participates in a standardized stroke data registry (e.g., GWTG-Stroke) for continuous quality measurement and public reporting where applicable.

4.6 The Stroke Unit maintains defined nurse-to-patient staffing ratios appropriate to patient acuity, with dedicated stroke-trained nursing staff on all shifts.

4.7 The Stroke Committee (IPC) develops and maintains a Stroke Program Charter documenting committee membership and roles, meeting frequency and attendance expectations, program goals, and facility demographics, reviewed and updated at minimum annually.

4.8 Program requirements are scaled to the certification level held or pursued by the organization (Acute Stroke Ready Center, Primary Stroke Center, or Comprehensive Stroke Center); where a requirement differs by certification level, this is noted within the applicable policy section.

4.9 A designated Telehealth Champion identifies and supports telestroke and related telehealth service opportunities in coordination with the Stroke Medical Director and Stroke Program Coordinator.

5. Multidisciplinary Stroke Team

The Stroke Team includes, at minimum, the following disciplines, each with defined roles documented in their respective privileging/competency files:

5.1 Stroke Medical Director (Neurologist).

5.2 Emergency Medicine Physicians.

5.3 Neurointerventionalist / Interventional Radiologist (for mechanical thrombectomy-capable and comprehensive centers).

5.4 Neurosurgery (on-call availability for hemorrhagic stroke and decompressive procedures).

5.5 Stroke Program Coordinator / Data Abstractor.

5.6 Stroke Unit Registered Nurses (NIHSS-certified).

5.7 Radiology / CT and MRI Technologists (24/7 availability).

5.8 Pharmacy (stroke order set review, thrombolytic preparation support).

5.9 Laboratory Services (STAT turnaround for stroke panel).

5.10 Physiatrist (Physical Medicine and Rehabilitation).

5.11 Physical Therapy, Occupational Therapy, and Speech-Language Pathology.

5.12 Case Management / Social Work.

5.13 Dietitian/Clinical Nutrition.

6. Staffing and Competency Requirements

6.1 All physicians and registered nurses providing direct stroke care must complete NIHSS certification (or equivalent) upon hire and recertify per organizational policy (minimum every 2 years or per certifying body requirement).

6.2 Stroke Unit nursing staff must complete a stroke-specific orientation program covering the acute stroke pathway, thrombolytic administration and monitoring, dysphagia screening, and post-stroke complication recognition prior to independent practice.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE PROGRAM ORGANIZATION, GOVERNANCE AND STAFFING

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-001	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Program Medical Director		
APPLIES TO	Hospital Wide — Emergency Department, Stroke Unit, ICU, Radiology, Laboratory, Pharmacy, Rehabilitation		

6.3 Physicians providing thrombolytic therapy must hold documented privileges for IV alteplase/tenecteplase administration.

6.4 A minimum staffing complement and on-call schedule is maintained to support 24/7/365 stroke code response, CT imaging, laboratory turnaround, and (where applicable) endovascular intervention.

7. Committee Structure and Performance Review

7.1 The Stroke Committee reviews, at minimum quarterly: door-to-CT times, door-to-needle times, door-to-groin puncture times (thrombectomy-capable/comprehensive centers), thrombolytic complication rates, GWTG-Stroke measure compliance, and mortality/morbidity outliers.

7.2 Performance data are benchmarked against AHA/ASA Target: Stroke® goals and TJC certification thresholds.

7.3 Identified gaps trigger a documented performance improvement action plan with assigned ownership and follow-up review.

8. References

8.1 American Heart Association/American Stroke Association. 2019 Guidelines for the Early Management of Patients with Acute Ischemic Stroke.

8.2 The Joint Commission. Comprehensive Stroke Center / Primary Stroke Center Certification Requirements.

8.3 AHA/ASA Get With The Guidelines®—Stroke Program Manual.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE CODE ACTIVATION — PRE-HOSPITAL AND IN-HOSPITAL

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-002	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, ICU, Inpatient Wards, Radiology		

SU-002 Stroke Code Activation — Pre-Hospital and In-Hospital

1. Purpose

To standardize the rapid identification, notification, and response process (“Stroke Code”) for any patient with suspected acute stroke, in order to minimize time from symptom onset/last known well to treatment, consistent with AHA/ASA Target: Stroke® time goals.

2. Definitions and Abbreviations

- 2.1 Last Known Well (LKW): The time the patient was last observed to be at their baseline neurological function.
- 2.2 FAST: Face, Arms, Speech, Time — a rapid stroke screening mnemonic.
- 2.3 BE-FAST: Balance, Eyes, Face, Arms, Speech, Time — an expanded screening tool incorporating posterior circulation signs.
- 2.4 Stroke Code / Code Stroke: The formal activation process that mobilizes the multidisciplinary stroke response team.
- 2.5 NIHSS: National Institutes of Health Stroke Scale.
- 2.6 EMS: Emergency Medical Services.

3. Policy

- 3.1 A Stroke Code may be activated by any nurse, physician, or EMS provider identifying signs/symptoms consistent with acute stroke using the BE-FAST screening tool, regardless of LKW time, unless clearly outside the therapeutic window for all interventions and the patient is not a candidate for any acute therapy.
- 3.2 Pre-hospital EMS providers use a validated stroke screening tool and provide advance hospital notification (“pre-notification”) for all suspected stroke patients, including LKW time, screening result, vital signs, and estimated time of arrival.
- 3.3 Pre-notified patients are met by the Stroke Team at the point of arrival and transferred directly to the CT scanner or a designated resuscitation/stroke bay.
- 3.4 Time targets (AHA/ASA Target: Stroke® Phase III goals, used as internal benchmarks):
 - 3.4.1 Door-to-physician evaluation: within 10 minutes of arrival.
 - 3.4.2 Door-to-CT initiation: within 20 minutes of arrival (goal), 25 minutes (TJC threshold).
 - 3.4.3 Door-to-CT interpretation: within 45 minutes of arrival.
 - 3.4.4 Door-to-needle (IV thrombolytic administration): within 60 minutes of arrival for ≥85% of patients (goal ≤45 minutes for ≥50% of patients).
 - 3.4.5 Door-to-groin puncture (mechanical thrombectomy): within 90 minutes of arrival (direct presentation) or 60 minutes (transfer patients), per institutional thrombectomy-capable/comprehensive stroke center goals.
 - 3.4.6 EMS-to-stroke-unit or ICU admission decision within 3 hours of arrival.

4. Procedure — In-Hospital Stroke Code Activation

- 4.1 Any staff member identifying a patient with acute focal neurological deficit(s) and LKW within the treatment window activates the Stroke Code by calling the designated hospital stroke code number/overhead page.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE CODE ACTIVATION — PRE-HOSPITAL AND IN-HOSPITAL

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-002	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, ICU, Inpatient Wards, Radiology		

4.2 The triage nurse or first responder immediately:

- 4.2.1 Notes and documents the exact LKW time (not the time the deficit was discovered).
- 4.2.2 Performs BE-FAST screening.
- 4.2.3 Obtains vital signs including finger-stick glucose.
- 4.2.4 Transfers the patient to a resuscitation/critical care bay.
- 4.2.5 Obtains IV access (18-gauge preferred) and draws STAT labs (CBC, basic metabolic panel/renal profile, coagulation studies [PT/INR, aPTT], troponin, and type & crossmatch/type & screen as per protocol).

4.3 The Stroke Code activation immediately and simultaneously notifies:

- 4.3.1 Emergency Physician / Stroke Team physician on duty.
- 4.3.2 On-call Neurologist / Stroke Medical Director.
- 4.3.3 CT Technologist and Radiology (to prioritize immediate imaging).
- 4.3.4 Stroke Unit / ICU charge nurse (bed availability).
- 4.3.5 Pharmacy (thrombolytic preparation on standby).
- 4.3.6 Laboratory (STAT stroke panel priority).

4.4 The responding physician performs a focused history and examination, including NIHSS scoring, and reviews the IV thrombolytic inclusion/exclusion checklist.

4.5 The patient is transported directly to CT for non-contrast head CT $\pm$ CT angiography (arch to vertex) $\pm$ CT perfusion, per the Acute Ischemic Stroke Management policy (SU-003).

4.6 All Stroke Code time points (activation time, physician arrival, CT arrival, CT completion, CT read, needle time, groin puncture time as applicable) are documented on the Stroke Code / Acute Stroke Flow Sheet for subsequent data abstraction and performance review.

4.7 In-hospital (“wake-up” or inpatient-onset) strokes follow the identical activation and time-target process; the LKW is the last time the inpatient was observed at baseline.

5. Procedure — Pre-Hospital Notification

5.1 EMS pre-notification calls are routed directly to the Emergency Department charge nurse/Stroke Team.

5.2 The receiving nurse relays LKW time, screening findings, vital signs, blood glucose (if obtained), anticoagulant use, and estimated arrival time to the Stroke Team.

5.3 A resuscitation bay and CT scanner are reserved in advance of patient arrival whenever feasible.

6. Documentation

All Stroke Code events are documented in the Nursing Acute Stroke Screening Form and Emergency Response Team Initial Assessment Form (see Attachments), including LKW time, activation time, and each subsequent time-stamped milestone.

7. References

- 7.1 AHA/ASA Target: Stroke® Phase III Initiative.
- 7.2 AHA/ASA 2019 Guidelines for the Early Management of Patients with Acute Ischemic Stroke.
- 7.3 The Joint Commission Stroke Certification Performance Measures.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE CODE ACTIVATION — PRE-HOSPITAL AND IN-HOSPITAL

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-002	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, ICU, Inpatient Wards, Radiology		

8. Attachments

- 8.1 Stroke Code / Acute Stroke Flow Sheet.
- 8.2 Nursing Acute Stroke Screening Form (BE-FAST).
- 8.3 EMS Pre-Notification Communication Form.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE ISCHEMIC STROKE MANAGEMENT — IV THROMBOLYSIS AND MECHANICAL THROMBECTOMY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-003	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Hospital Wide		

SU-003 Acute Ischemic Stroke Management — IV Thrombolysis and Mechanical Thrombectomy

1. Purpose

To delineate the evidence-based steps for the assessment, imaging, and treatment (intravenous thrombolysis and/or mechanical thrombectomy) of patients presenting with acute ischemic stroke, aligned with the 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke (and subsequent focused updates).

2. Definitions and Abbreviations

- 2.1 IV tPA / Alteplase: Recombinant tissue plasminogen activator, an IV thrombolytic agent.
- 2.2 Tenecteplase (TNK): Alternative IV thrombolytic agent, single-bolus administration, acceptable alternative to alteplase per AHA/ASA guidance.
- 2.3 NIHSS: National Institutes of Health Stroke Scale.
- 2.4 LKW: Last Known Well.
- 2.5 ASPECTS: Alberta Stroke Program Early CT Score, used to quantify early ischemic changes on non-contrast CT.
- 2.6 CTA: CT Angiography. CTP: CT Perfusion.
- 2.7 MT: Mechanical Thrombectomy (endovascular therapy).
- 2.8 LVO: Large Vessel Occlusion.

3. Policy

- 3.1 IV thrombolytic therapy improves functional outcome when administered within the appropriate therapeutic window and is the standard of care for eligible patients with disabling acute ischemic stroke.
- 3.2 Door-to-needle time is a primary quality metric and should be ≤60 minutes for ≥85% of treated patients (institutional goal ≤45 minutes where feasible), consistent with AHA/ASA Target: Stroke®.
- 3.3 Mechanical thrombectomy is offered to eligible patients with confirmed large vessel occlusion, in coordination with a certified neurointerventional service (on-site or via a defined transfer agreement with a Comprehensive Stroke Center / Thrombectomy-Capable Stroke Center).
- 3.4 Eligibility for both IV thrombolysis and mechanical thrombectomy is determined using current AHA/ASA inclusion/exclusion criteria, applied by a physician credentialed in acute stroke management.
- 3.5 Treatment decisions and risks/benefits are discussed with the patient or legally authorized representative and documented as informed consent, consistent with organizational policy for emergency/time-critical procedures.

4. Procedure — Initial Assessment (0–6 Hours from Onset/LKW)

- 4.1 Following Stroke Code activation (SU-002), the responding physician performs a focused history, NIHSS examination, and reviews the IV thrombolytic inclusion/exclusion checklist.
- 4.2 Airway, breathing, and circulation are stabilized; supplemental oxygen is provided to maintain SpO₂ ≥ 94%.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE ISCHEMIC STROKE MANAGEMENT — IV THROMBOLYSIS AND MECHANICAL THROMBECTOMY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-003	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Hospital Wide		

4.3 STAT labs are sent (CBC, renal profile, coagulation studies, glucose, troponin, type & screen); treatment is not delayed pending coagulation results unless the patient is on anticoagulants, has a bleeding disorder, or thrombocytopenia is suspected.

4.4 Non-contrast head CT is obtained emergently (goal: CT initiated within 20–25 minutes of arrival).

4.5 If no hemorrhage is identified and the patient meets criteria, CT angiography (arch to vertex) is obtained to identify large vessel occlusion and determine candidacy for mechanical thrombectomy.

4.6 If hemorrhage is identified, the patient is managed per the Hemorrhagic Stroke Management policy (SU-004) and neurosurgery is consulted.

5. Procedure — IV Thrombolytic Administration

5.1 Blood pressure must be <185/110 mmHg prior to thrombolytic administration; antihypertensive therapy (e.g., labetalol, nicardipine, or clevidipine per organizational protocol) is administered if needed. If blood pressure cannot be safely lowered below this threshold, thrombolytic therapy is withheld.

5.2 Alteplase is dosed at 0.9 mg/kg (maximum total dose 90 mg): 10% administered as an IV bolus over 1 minute, followed by the remaining 90% infused over 60 minutes. (If tenecteplase is used per institutional protocol, it is administered as a single IV bolus per the approved organizational order set.)

5.3 Extended window treatment (3–4.5 hours from LKW) follows the same eligibility criteria with additional exclusion considerations (e.g., age >80, oral anticoagulant use regardless of INR, baseline NIHSS >25, combined history of prior stroke and diabetes, or imaging evidence of infarction involving more than one-third of the MCA territory), per current AHA/ASA guidance and physician judgment.

5.4 During and after infusion, the patient is monitored in a high-dependency setting (ICU, Stroke Unit, or ED resuscitation area) with neurological checks and blood pressure monitoring every 15 minutes for 2 hours, then every 30 minutes for 6 hours, then hourly for 16 hours.

5.5 Blood pressure is maintained <180/105 mmHg for at least the first 24 hours post-thrombolysis.

5.6 No antiplatelet or anticoagulant agents are administered for 24 hours following thrombolysis, pending a follow-up non-contrast CT to exclude hemorrhagic transformation.

5.7 Any acute neurological deterioration, severe headache, new hypertension, nausea, or vomiting during or after infusion triggers immediate cessation of the infusion (if ongoing), STAT non-contrast head CT, and physician notification to evaluate for symptomatic intracranial hemorrhage.

6. Procedure — Mechanical Thrombectomy

6.1 Patients with confirmed anterior circulation LVO, NIHSS ≥6, pre-stroke independent functional status, and favorable imaging profile (e.g., ASPECTS ≥6) are evaluated for mechanical thrombectomy up to 24 hours from LKW using clinical-imaging mismatch criteria (CT perfusion or MRI-based core/penumbra mismatch) for the 6–24-hour window.

6.2 The neurointerventionalist/neurosurgery team is notified immediately upon identification of a thrombectomy candidate; if not available on-site, an emergent transfer to the designated thrombectomy-capable partner facility is initiated per the organization's stroke transfer agreement.

6.3 IV thrombolysis (if the patient is eligible and within window) is not withheld while arranging or performing mechanical thrombectomy.

6.4 Post-procedure patients are admitted to the ICU or Stroke Unit for neurological and access-site monitoring.

7. Procedure — Extended Window Evaluation (6–24 Hours)

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE ISCHEMIC STROKE MANAGEMENT — IV THROMBOLYSIS AND MECHANICAL THROMBECTOMY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-003	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Hospital Wide		

7.1 Patients presenting 6–24 hours from LKW are screened using BE-FAST; a positive screen prompts CT stroke protocol (non-contrast CT, CTA, CT perfusion).

7.2 Clinical-imaging mismatch is assessed using validated automated perfusion software, applying core infarct volume and NIHSS thresholds consistent with the pivotal randomized trial criteria referenced in current AHA/ASA guidelines.

7.3 Patients meeting mismatch criteria are evaluated for mechanical thrombectomy per Section 6.

8. General Supportive Care (All Acute Ischemic Stroke Patients)

8.1 Supplemental oxygen to maintain SpO₂ ≥ 94%.

8.2 Continuous cardiac monitoring for at least the first 48 hours to detect atrial fibrillation and other arrhythmias.

8.3 Treat hyperthermia (temperature >38°C) with antipyretics and identify the underlying source.

8.4 Correct hypovolemia with isotonic IV fluids; avoid fluid overload in patients with reduced ejection fraction.

8.5 Permissive hypertension in patients not receiving thrombolysis/thrombectomy: withhold antihypertensive therapy unless systolic BP >220 mmHg or diastolic BP >120 mmHg, or unless a compelling comorbid indication exists (e.g., acute coronary syndrome, aortic dissection, hypertensive encephalopathy); when treatment is indicated, lower BP by approximately 15% during the first 24 hours.

8.6 Initiate VTE prophylaxis (intermittent pneumatic compression ± prophylactic-dose subcutaneous anticoagulation per institutional VTE prophylaxis policy) as early as clinically appropriate.

8.7 Initiate aspirin within 24–48 hours of stroke onset in patients not receiving thrombolysis, or 24 hours after thrombolysis once hemorrhage is excluded, unless contraindicated.

8.8 Statin therapy is initiated or continued during hospitalization per current lipid management guidelines.

8.9 Dysphagia screening is completed prior to any oral intake, per the NIHSS/Dysphagia Screening policy (SU-006).

9. Admission Criteria

Patients are admitted to ICU, Stroke Unit, or a monitored bed based on treatment received and clinical status, per the criteria below:

9.1 ICU/high-dependency admission: patients who received IV thrombolysis and/or mechanical thrombectomy; acute stroke with depressed consciousness; stroke with seizures; hemodynamic or respiratory instability.

9.2 Stroke Unit admission: hemodynamically stable acute ischemic stroke or TIA not requiring ICU-level monitoring, per the Stroke Unit Admission/Discharge Criteria in policy SU-001/SU-005.

9.3 Any patient with new focal deficit, GCS decline, need for airway protection, or hemorrhagic transformation is escalated to ICU immediately.

10. References

10.1 Powers WJ, et al. 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke. Stroke.

10.2 AHA/ASA Target: Stroke® Best Practice Strategies.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE ISCHEMIC STROKE MANAGEMENT — IV THROMBOLYSIS AND MECHANICAL THROMBECTOMY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-003	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Hospital Wide		

10.3 The Joint Commission Comprehensive/Primary Stroke Center Certification Standards.

11. Attachments

- 11.1 IV Thrombolytic Inclusion/Exclusion Checklist.
- 11.2 Thrombolytic Therapy (Alteplase/Tenecteplase) Administration and Monitoring Form.
- 11.3 Post-tPA Neurological and Vital Sign Monitoring Flow Sheet.
- 11.4 Mechanical Thrombectomy Transfer Checklist (if applicable).

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: SPONTANEOUS INTRACEREBRAL AND SUBARACHNOID HEMORRHAGE MANAGEMENT

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-004	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Emergency Department, ICU, Stroke Unit, Neurosurgery		

SU-004 Spontaneous Intracerebral and Subarachnoid Hemorrhage Management

1. Purpose

To standardize the emergent evaluation, blood pressure management, reversal of anticoagulation, and neurosurgical coordination for patients with spontaneous intracerebral hemorrhage (ICH) or subarachnoid hemorrhage (SAH), consistent with AHA/ASA guidelines for the management of spontaneous intracerebral hemorrhage and aneurysmal subarachnoid hemorrhage.

2. Definitions

- 2.1 ICH: Intracerebral (parenchymal) hemorrhage.
- 2.2 SAH: Subarachnoid hemorrhage.
- 2.3 AVM: Arteriovenous malformation.
- 2.4 ICP: Intracranial pressure.
- 2.5 EVD: External ventricular drain.

3. Policy

- 3.1 Any hemorrhage identified on initial CT during Stroke Code evaluation triggers immediate neurosurgical notification and CT angiography of the Circle of Willis to evaluate for a secondary vascular cause (AVM, aneurysm, arteriovenous fistula).
- 3.2 Patients with ICH or SAH are managed in the ICU with continuous neurological and hemodynamic monitoring.
- 3.3 Anticoagulant reversal, when indicated, is initiated emergently without waiting for transfer or definitive imaging, per the organizational anticoagulation reversal protocol.
- 3.4 Blood pressure management follows AHA/ASA ICH guideline targets to reduce hematoma expansion risk while maintaining cerebral perfusion.

4. Procedure — Intracerebral Hemorrhage

- 4.1 Confirm ICH on non-contrast CT; obtain CT angiography (arch to vertex / Circle of Willis) to evaluate for an underlying structural lesion, particularly in younger patients, those without a history of hypertension, or with atypical hemorrhage location.
- 4.2 Notify neurosurgery immediately for evaluation of surgical candidacy (hematoma evacuation, ventriculostomy) and level of care.
- 4.3 Blood pressure management: for patients presenting with systolic BP 150–220 mmHg, acute lowering to a target systolic BP of approximately 140 mmHg is reasonable and can be safe, using IV agents such as labetalol, nicardipine, or clevidipine, with reassessment based on individual clinical context.
- 4.4 Reverse anticoagulation emergently in patients on vitamin K antagonists (e.g., with 4-factor prothrombin complex concentrate and IV vitamin K), direct oral anticoagulants (with specific reversal agents where available, or PCC), or heparin (protamine sulfate), per the organizational reversal protocol.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: SPONTANEOUS INTRACEREBRAL AND SUBARACHNOID HEMORRHAGE MANAGEMENT

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-004	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Emergency Department, ICU, Stroke Unit, Neurosurgery		

4.5 Correct severe thrombocytopenia or coagulation factor deficiencies as clinically indicated; platelet transfusion is not routinely recommended for antiplatelet-associated ICH without a specific procedural indication.

4.6 Manage elevated ICP as needed: head-of-bed elevation to 30°, analgesia/sedation as needed, osmotic therapy (mannitol or hypertonic saline) for signs of herniation or elevated ICP, and ventriculostomy for hydrocephalus per neurosurgical guidance.

4.7 Maintain normoglycemia, normothermia, and seizure precautions; prophylactic anticonvulsants are not routinely recommended in the absence of clinical or electrographic seizures.

4.8 VTE prophylaxis with intermittent pneumatic compression is initiated on admission; pharmacologic prophylaxis may be added after documented cessation of bleeding, typically 24–48 hours after onset, per physician judgment.

4.9 Decompressive craniectomy ± hematoma evacuation is considered for select patients with large hemorrhage and mass effect or clinical deterioration, per neurosurgical evaluation.

5. Procedure — Subarachnoid Hemorrhage

5.1 Suspected SAH (thunderclap headache, meningismus) with a non-diagnostic non-contrast CT is further evaluated with lumbar puncture (evaluating for xanthochromia) if clinically indicated and CT is negative but suspicion remains high.

5.2 Confirmed SAH triggers emergent CT angiography and neurosurgical/neurointerventional consultation for aneurysm identification and securement (surgical clipping or endovascular coiling), ideally within 24–72 hours.

5.3 Nimodipine 60 mg PO/NG every 4 hours is initiated for vasospasm prophylaxis, per protocol, unless contraindicated by hypotension.

5.4 Blood pressure is controlled to reduce rebleeding risk prior to aneurysm securement, balanced against maintaining adequate cerebral perfusion; specific targets are individualized with neurosurgery/neurocritical care.

5.5 Patients are monitored for delayed cerebral ischemia (vasospasm) with serial neurological assessments and, where available, transcranial Doppler monitoring, typically most pronounced days 4–14 post-hemorrhage.

5.6 Hydrocephalus is managed with external ventricular drainage per neurosurgical assessment.

5.7 Euvolemia is maintained; prophylactic hypervolemic or induced hypertensive therapy is not used routinely but may be considered for symptomatic vasospasm under neurocritical care guidance.

6. Admission and Escalation Criteria

6.1 All ICH and SAH patients are admitted to the ICU.

6.2 Escalation to neurosurgical intervention is triggered by: GCS decline ≥ 2 points, new anisocoria, hematoma expansion on repeat imaging, hydrocephalus, or signs of herniation.

7. References

7.1 Greenberg SM, et al. 2022 AHA/ASA Guideline for the Management of Patients with Spontaneous Intracerebral Hemorrhage. Stroke.

7.2 Hemphill JC, et al. AHA/ASA Guidelines for the Management of Aneurysmal Subarachnoid Hemorrhage.

7.3 The Joint Commission Comprehensive Stroke Center Certification Standards.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: SPONTANEOUS INTRACEREBRAL AND SUBARACHNOID HEMORRHAGE MANAGEMENT

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-004	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Emergency Department, ICU, Stroke Unit, Neurosurgery		

8. Attachments

- 8.1 Anticoagulation Reversal Protocol.
- 8.2 ICH Blood Pressure Management Order Set.
- 8.3 SAH / Vasospasm Monitoring Flow Sheet.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: TRANSIENT ISCHEMIC ATTACK (TIA) EVALUATION AND MANAGEMENT

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-005	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, Outpatient Neurology Clinic		

SU-005 Transient Ischemic Attack (TIA) Evaluation and Management

1. Purpose

To standardize the risk stratification, urgent evaluation, and secondary prevention management of patients presenting with transient ischemic attack (TIA), given the substantial short-term risk of subsequent stroke.

2. Definitions

2.1 TIA: A transient episode of neurological dysfunction caused by focal brain, spinal cord, or retinal ischemia, without evidence of acute infarction on imaging.

2.2 ABCD² Score: A clinical risk-stratification tool for early stroke risk after TIA (Age, Blood pressure, Clinical features, Duration of symptoms, Diabetes).

2.3 Crescendo TIA: Recurrent TIAs with increasing frequency, duration, or severity over a short period.

3. Policy

3.1 All patients presenting with suspected TIA undergo the same urgent Stroke Code evaluation pathway as acute ischemic stroke (SU-002), because clinical differentiation between TIA and acute ischemic stroke cannot reliably be made on presentation.

3.2 Patients with TIA are at significantly increased short-term risk of stroke (up to approximately 10–20% within 90 days, with the highest risk in the first 48 hours), warranting urgent risk stratification and initiation of secondary prevention.

3.3 High-risk TIA patients (including crescendo TIA, ABCD² ≥4, or TIA due to identified large-vessel disease/atrial fibrillation) are admitted for expedited work-up; lower-risk patients may be managed through an expedited outpatient TIA pathway when a same-day/next-day evaluation can be reliably arranged.

4. Procedure

4.1 Perform BE-FAST/NIHSS assessment and non-contrast head CT (or MRI with diffusion-weighted imaging where readily available) to exclude acute infarction or hemorrhage.

4.2 Calculate and document the ABCD² score to support admission/disposition decision-making, used in conjunction with overall clinical judgment.

4.3 Obtain vascular imaging (CT angiography or MR angiography of the head and neck) to evaluate for large-vessel stenosis or occlusion.

4.4 Obtain cardiac evaluation: 12-lead ECG on arrival, and continuous cardiac (telemetry) monitoring for at least 24–48 hours to detect paroxysmal atrial fibrillation; extended outpatient cardiac monitoring is arranged for patients with cryptogenic events and no arrhythmia identified during admission.

4.5 Obtain echocardiography (transthoracic, with transesophageal as indicated) in patients with suspected cardioembolic source, particularly those <60 years or with cryptogenic TIA.

4.6 Initiate antiplatelet therapy (aspirin, or short-term dual antiplatelet therapy with aspirin plus clopidogrel for 21–90 days in select high-risk minor stroke/TIA patients per current AHA/ASA guidance) unless anticoagulation is indicated for a cardioembolic source.

4.7 Initiate or optimize statin therapy and blood pressure management per secondary prevention guidelines.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: TRANSIENT ISCHEMIC ATTACK (TIA) EVALUATION AND MANAGEMENT

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-005	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, Outpatient Neurology Clinic		

4.8 For patients with symptomatic carotid stenosis $\geq 70\%$ (or 50–69% with additional risk factors), refer promptly for carotid revascularization evaluation (carotid endarterectomy or carotid artery stenting), ideally within 2 weeks of the event.

4.9 Provide stroke risk-factor education and secondary prevention counseling prior to discharge, per policy SU-008.

4.10 Arrange follow-up with neurology within 7 days of discharge for all TIA patients managed as outpatients or discharged from observation.

5. Admission Criteria

5.1 Crescendo or recurrent TIA.

5.2 ABCD² score ≥ 4 .

5.3 Identified symptomatic large-vessel stenosis or intracranial/extracranial occlusion.

5.4 Suspected cardioembolic source requiring further work-up.

5.5 Inability to reliably complete urgent outpatient evaluation within 24–48 hours.

6. References

6.1 Kleindorfer DO, et al. 2021 AHA/ASA Guideline for the Prevention of Stroke in Patients with Stroke and Transient Ischemic Attack. Stroke.

6.2 Johnston SC, et al. Validation and refinement of the ABCD² score.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: NIHSS ASSESSMENT, NEUROLOGICAL MONITORING AND DYSPHAGIA/SWALLOW SCREENING

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-006	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Program Coordinator / Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, ICU		

SU-006 NIHSS Assessment, Neurological Monitoring and Dysphagia/Swallow Screening

1. Purpose

To standardize neurological assessment using the NIH Stroke Scale (NIHSS) and to ensure all stroke patients are screened for dysphagia prior to any oral intake, reducing the risk of aspiration pneumonia — a core AHA/ASA and Joint Commission stroke performance measure.

2. Policy — NIHSS

- 2.1 All physicians and registered nurses performing stroke assessments must be NIHSS-certified prior to independently scoring the scale, with recertification per organizational policy.
- 2.2 NIHSS is documented at initial presentation, and at defined intervals thereafter (e.g., 2 hours and 24 hours post-thrombolysis, and with any acute neurological change) to detect early deterioration or improvement.
- 2.3 A sustained increase in NIHSS of ≥ 4 points, or any acute decline in level of consciousness, triggers immediate physician notification and evaluation, including consideration of an urgent non-contrast head CT to exclude hemorrhagic transformation or infarct progression.

3. Policy — Dysphagia / Swallow Screening

- 3.1 All acute stroke patients remain nil per oral (NPO), including oral medications, until a formal swallow screen is completed and passed.
- 3.2 A validated swallow screening tool (e.g., a standardized water-swallow protocol) is administered by an appropriately trained clinician (typically a trained nurse) within 4 hours of arrival, and prior to any oral intake of food, fluid, or medication.
- 3.3 Swallow screening is deferred only for patients who are NPO for other reasons (e.g., planned intervention) or who are not alert enough to participate; screening is completed as soon as the patient is alert and able to participate.
- 3.4 Patients who fail the screening remain NPO and are referred to a speech-language pathologist for comprehensive assessment, to be completed within 24 hours of referral.
- 3.5 Instrumental evaluation (fiberoptic endoscopic evaluation of swallowing [FEES] or videofluoroscopic swallow study) is performed when needed to confirm the presence/absence of aspiration and guide the treatment plan.
- 3.6 Patients with confirmed dysphagia receive individualized dietary modification, compensatory swallowing strategies, and swallowing exercises as appropriate.
- 3.7 Patients who fail repeat swallow assessment or require prolonged NPO status are considered for enteral nutrition access (nasogastric, nasoduodenal, or percutaneous endoscopic gastrostomy) to maintain nutrition and hydration.
- 3.8 Swallow screening is repeated whenever a patient's neurological or medical status changes.

4. Documentation

NIHSS scores and swallow screening results (pass/fail, time completed, screening tool used) are documented in the medical record and are subject to concurrent and retrospective audit as part of the stroke performance improvement program (SU-009), consistent with GWTG-Stroke/Joint Commission measure STK-1 (dysphagia screening) requirements.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: NIHSS ASSESSMENT, NEUROLOGICAL MONITORING AND DYSPHAGIA/SWALLOW SCREENING

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-006	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Program Coordinator / Medical Director		
APPLIES TO	Emergency Department, Stroke Unit, ICU		

5. References

- 5.1 National Institutes of Health Stroke Scale (NIHSS) Training and Certification materials.
- 5.2 AHA/ASA Get with the Guidelines®—Stroke Dysphagia Screening Measure Specifications.
- 5.3 The Joint Commission Stroke Performance Measures (STK set).

6. Attachments

- 6.1 NIHSS Scoring Form.
- 6.2 Bedside Swallow Screening Tool.
- 6.3 Speech-Language Pathology Comprehensive Dysphagia Assessment Form.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE REHABILITATION AND EARLY MOBILIZATION

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-007	—	]Effective Date[	]Review Date[
APPROVED BY	Physiatrist / Medical Director		
APPLIES TO	Stroke Unit, ICU, Rehabilitation Services		

SU-007 Stroke Rehabilitation and Early Mobilization

1. Purpose

To ensure timely, structured, multidisciplinary rehabilitation assessment and intervention for all stroke patients, consistent with AHA/ASA guidelines for stroke rehabilitation and recovery, in order to optimize functional outcomes.

2. Policy

2.1 All patients admitted with a diagnosis of stroke receive a physiatry/rehabilitation medicine consultation within 24 hours of admission.

2.2 All patients undergo a full multidisciplinary rehabilitation needs assessment (physical therapy, occupational therapy, speech-language pathology, as indicated), documented in the multidisciplinary plan of care.

2.3 Early mobilization is initiated within 24 hours of stroke onset for medically stable patients, by an appropriately trained healthcare professional with access to appropriate mobility equipment; very early (within 24 hours), high-frequency, high-dose out-of-bed mobilization is avoided per trial evidence, and mobilization intensity is individualized.

2.4 A discharge rehabilitation disposition (home with outpatient therapy, home health, inpatient rehabilitation facility, or skilled nursing facility) is determined by the multidisciplinary team prior to discharge, based on functional assessment.

3. Procedure — Rehabilitation Assessment

3.1 The physiatrist assesses the patient and documents individualized goals and a plan of care in the Physical Medicine Request Form and multidisciplinary plan of care form.

3.2 Assessment addresses: motor function (strength, tone, coordination), sensory function, swallowing (per SU-006), bladder and bowel function, skin integrity/pressure injury risk, cognition/communication, vision/hearing, nutritional status, and psychosocial needs.

3.3 A skin/pressure-injury prevention and positioning program is established, addressing contracture prevention, respiratory function, and use of specialized equipment (splints, braces) as needed.

4. Procedure — Therapeutic Interventions

4.1 Physical and occupational therapists implement a motor control/retraining program including facilitation/inhibition techniques, stretching, biofeedback, therapeutic exercise, and neuromuscular electrical stimulation as clinically indicated.

4.2 Functional mobility training addresses: bed mobility, transfers (bed, chair, wheelchair), stair negotiation, ambulation with or without assistive devices, and wheelchair mobility/propulsion, individualized to the patient's deficits and goals.

4.3 Occupational therapy addresses activities of daily living, upper-extremity function, cognitive-perceptual deficits, and home/environmental modification needs.

4.4 Speech-language pathology addresses communication deficits (aphasia, dysarthria) and cognitive-communication impairments in addition to dysphagia management (SU-006).

4.5 Psychological/neuropsychological support is arranged for patients with post-stroke depression, anxiety, or cognitive impairment affecting rehabilitation participation.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE REHABILITATION AND EARLY MOBILIZATION

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-007	—	]Effective Date[	]Review Date[
APPROVED BY	Physiatrist / Medical Director		
APPLIES TO	Stroke Unit, ICU, Rehabilitation Services		

5. Discharge Rehabilitation Planning

5.1 The multidisciplinary team, in coordination with Case Management, determines the appropriate level of post-acute rehabilitation care based on functional status, medical stability, cognitive capacity to participate in therapy, and caregiver/family support.

5.2 All stroke patients are formally assessed for and considered for rehabilitation services prior to discharge, and this assessment is documented, consistent with the GWTG-Stroke/Joint Commission rehabilitation assessment performance measure.

6. References

6.1 Winstein CJ, et al. AHA/ASA Guidelines for Adult Stroke Rehabilitation and Recovery. Stroke.

6.2 AHA/ASA Get with the Guidelines®—Stroke Rehabilitation Assessment Measure Specifications.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: DISCHARGE PLANNING, SECONDARY PREVENTION AND PATIENT/FAMILY EDUCATION

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-008	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Program Coordinator		
APPLIES TO	Stroke Unit, ICU, Case Management		

SU-008 Discharge Planning, Secondary Prevention and Patient/Family Education

1. Purpose

To ensure that every stroke and TIA patient is discharged with an evidence-based secondary prevention regimen, appropriate follow-up, and documented patient/family education, consistent with AHA/ASA secondary prevention guidelines and Joint Commission stroke discharge performance measures (STK set).

2. Policy

- 2.1 Discharge planning begins at admission and is reassessed daily by the multidisciplinary team.
- 2.2 Prior to discharge, every eligible ischemic stroke/TIA patient is prescribed antithrombotic therapy (antiplatelet or anticoagulant, as clinically indicated) unless contraindicated and documented.
- 2.3 Patients with atrial fibrillation/flutter are prescribed anticoagulation at discharge unless contraindicated and documented.
- 2.4 Patients with LDL ≥ 100 mg/dL, or with atherosclerotic disease regardless of LDL, are prescribed statin therapy at discharge unless contraindicated.
- 2.5 Smoking cessation counseling and resources are provided to all current smokers.
- 2.6 Formal stroke education is provided to the patient and/or caregiver, covering the required core topics in Section 3.
- 2.7 A stroke rehabilitation needs assessment is documented prior to discharge (SU-007).
- 2.8 Follow-up appointment(s) are scheduled prior to discharge, including neurology follow-up within 7–14 days as clinically appropriate.

3. Required Patient/Family Education Topics

- 3.1 Personal risk factors for stroke and individualized modification strategies (hypertension, diabetes, dyslipidemia, atrial fibrillation, smoking, physical inactivity, obesity).
- 3.2 Warning signs and symptoms of stroke (BE-FAST) and the importance of calling emergency services immediately if symptoms recur.
- 3.3 Prescribed medications: name, purpose, dose, schedule, and importance of adherence, including anticoagulant/antiplatelet-specific precautions.
- 3.4 Follow-up care: scheduled appointments, and who to contact with questions or concerns.
- 3.5 Activity, dietary, and lifestyle recommendations, including dysphagia diet modifications if applicable.
- 3.6 Community resources, support groups, and caregiver support services.

4. Discharge Criteria from the Stroke Unit

Discharge or transfer from the Stroke Unit is at the discretion of the treating physician in consultation with the referring physician/department, guided by the following:

- 4.1 The patient is neurologically and hemodynamically stable, with status continuously reassessed to identify patients who no longer require Stroke Unit–level care.
- 4.2 The patient's physiologic status has stabilized such that intensive monitoring is no longer required, or no further benefit is expected from continued Stroke Unit stay.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: DISCHARGE PLANNING, SECONDARY PREVENTION AND PATIENT/FAMILY EDUCATION

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-008	—	]Effective Date[	]Review Date[
APPROVED BY	Medical Director / Stroke Program Coordinator		
APPLIES TO	Stroke Unit, ICU, Case Management		

4.3 The etiology of the stroke has been identified, or a clear plan for completion of the diagnostic work-up (inpatient or outpatient) has been established.

4.4 If physiologic status has deteriorated and further active intervention is not planned (e.g., transition to comfort-focused care), transfer to a lower level of care is appropriate.

4.5 Disposition (home, home with services, inpatient rehabilitation, skilled nursing facility) is finalized in coordination with Case Management/Social Work and the patient/family.

5. Documentation

Discharge medication reconciliation, education topics covered, and follow-up plans are documented in the discharge summary and nursing discharge instructions, and are subject to audit under the stroke performance improvement program (SU-009).

6. References

6.1 Kleindorfer DO, et al. 2021 AHA/ASA Guideline for the Prevention of Stroke in Patients with Stroke and Transient Ischemic Attack.

6.2 AHA/ASA Get with the Guidelines®—Stroke Discharge Measure Specifications (STK-1 through STK-10).

6.3 The Joint Commission Stroke Certification Performance Measures.

7. Attachments

7.1 Stroke Discharge Checklist.

7.2 Patient/Family Stroke Education Booklet and Documentation Form.

7.3 Discharge Medication Reconciliation Form.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE PERFORMANCE IMPROVEMENT, DATA REGISTRY AND PEER REVIEW

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-009	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-009 Stroke Performance Improvement, Data Registry and Peer Review

1. Purpose

To establish a continuous, data-driven performance improvement process for the Stroke Program, including participation in a standardized national stroke registry, in fulfillment of AHA/ASA Get with the Guidelines®–Stroke and Joint Commission stroke certification requirements.

2. Policy

2.1 All patients with a confirmed diagnosis of ischemic stroke, hemorrhagic stroke, TIA, or subarachnoid hemorrhage are entered into the stroke data registry by the Stroke Program Coordinator/data abstractor.

2.2 Core performance measures are tracked continuously and reported to the Stroke Committee at minimum quarterly, including but not limited to:

2.2.1 STK-1: VTE prophylaxis by end of hospital day 2.

2.2.2 STK-2: Discharge antithrombotic therapy.

2.2.3 STK-3: Anticoagulation for atrial fibrillation/flutter at discharge.

2.2.4 STK-4: Thrombolytic therapy administered to eligible patients arriving within 3.5 hours of LKW/onset.

2.2.5 STK-5: Antithrombotic therapy by end of hospital day 2.

2.2.6 STK-6: Discharge statin therapy.

2.2.7 STK-8: Stroke education provided.

2.2.8 STK-10: Assessed for rehabilitation services.

2.2.9 Door-to-CT, door-to-needle, and (where applicable) door-to-groin-puncture time metrics.

2.2.10 Symptomatic intracranial hemorrhage rate following thrombolytic therapy.

2.2.11 (Comprehensive Stroke Center) Hemorrhagic transformation rate within 36 hours of treatment, overall and stratified by IV-only versus intra-arterial/mechanical reperfusion therapy.

2.2.12 (Comprehensive Stroke Center) Thrombolysis in Cerebral Infarction (TICI) post-treatment reperfusion grade of 2B or higher.

2.2.13 (Comprehensive Stroke Center) Modified Rankin Scale (mRS) ≤ 2 at 90 days post-treatment (assessment window 75–105 days).

2.2.14 (Comprehensive Stroke Center) Timeliness of reperfusion: arrival-to-TICI-2B time (≤ 150 minutes) and skin-puncture-to-TICI-2B time (≤ 60 minutes).

2.3 Cases with a protocol deviation, unexpected complication, or mortality undergo structured peer review (Stroke Morbidity & Mortality review), with findings reported to the Stroke Committee and, as applicable, the Medical Staff Quality/Peer Review Committee.

2.4 Performance below internal or national benchmark thresholds triggers a documented corrective action plan with assigned responsibility, timeline, and re-measurement.

2.5 Aggregate and case-level data are used to inform staff education, protocol revision, and resource allocation decisions.

2.6 An annual Stroke Program performance report is prepared summarizing volumes, outcome metrics, and quality improvement initiatives, for review by hospital leadership and the Medical Staff.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: STROKE PERFORMANCE IMPROVEMENT, DATA REGISTRY AND PEER REVIEW

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-009	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

3. Procedure

- 3.1 Concurrent case identification: stroke/TIA cases are flagged in real time via the Stroke Code activation log, ED diagnosis coding, and daily census review by the Stroke Coordinator.
- 3.2 Data abstraction: the Stroke Coordinator abstracts required data elements into the registry (e.g., GWTG-Stroke) within the timeframe specified by the registry's data completion requirements.
- 3.3 Quarterly Stroke Committee review of aggregate performance measures, benchmarked against AHA/ASA and Joint Commission thresholds.
- 3.4 Case-level review of all thrombolytic/thrombectomy complications, in-hospital stroke mortality, and any case flagged by nursing, physician, or risk management referral.
- 3.5 Action plans arising from performance gaps are tracked to closure and re-audited at the subsequent Committee meeting.

4. References

- 4.1 AHA/ASA Get with the Guidelines®—Stroke Program Manual and Measure Specifications.
- 4.2 The Joint Commission Stroke Performance Measures (STK Measure Set) and Certification Manual.

5. Attachments

- 5.1 Stroke Registry Data Abstraction Tool.
- 5.2 Stroke Morbidity & Mortality Review Form.
- 5.3 Quarterly Stroke Committee Performance Dashboard Template.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CODE STROKE TEAM — ROLES, RESPONSIBILITIES AND ACTIVATION ROSTER

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-010	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-010 Code Stroke Team — Roles, Responsibilities and Activation Roster

1. Purpose

To define the composition, individual roles, and activation roster of the Code Stroke Team, and to ensure that every member understands their specific responsibilities during a Stroke Code so that the team functions as a coordinated unit from activation through disposition.

2. Scope

This policy applies to all physicians, nurses, technologists, pharmacists, and allied staff who are designated members of the Code Stroke Team, and to any staff member responsible for activating or responding to a Stroke Code, in any hospital location (Emergency Department, inpatient units, ICU, Radiology, or off-site areas).

3. Definitions and Abbreviations

3.1 Code Stroke Team: The designated multidisciplinary group that responds immediately to a Stroke Code activation, as distinct from the broader Multidisciplinary Stroke Team described in SU-001, which also includes rehabilitation and longitudinal care disciplines.

3.2 Team Leader: The physician (Emergency Physician or Neurologist, per institutional assignment) who directs the Code Stroke response and has final authority for treatment decisions.

3.3 Recorder: The nurse or designated staff member responsible for real-time timestamping of all Stroke Code milestones on the Stroke Code Flow Sheet (Appendix C).

3.4 Runner: Support staff member who physically transports labs, blood products, or imaging orders when electronic/pneumatic systems are unavailable or delayed.

4. Policy

4.1 A designated Code Stroke Team responds to every Stroke Code activation (SU-002) within the defined time targets, 24 hours a day, 7 days a week, 365 days a year.

4.2 Each team role is filled by a specific position on every shift; role assignments are posted on the daily staffing assignment board and are not dependent on any single named individual.

4.3 Team members carry a dedicated Stroke Code pager/communication device or are reachable through the hospital's overhead/electronic paging system at all times while on duty.

4.4 All Code Stroke Team members complete role-specific competency validation upon assignment to the team and annually thereafter, in accordance with SU-015 (Nursing and Staff Stroke Education Program).

4.5 A debrief is conducted after every Stroke Code involving thrombolytic administration, a complication, or a significant time-target miss, to identify system issues and reinforce role performance.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CODE STROKE TEAM — ROLES, RESPONSIBILITIES AND ACTIVATION ROSTER

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-010	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5. Team Roles and Responsibilities

Role	Primary Responder	Key Responsibilities
Team Leader	Emergency Physician / Neurologist on duty	Directs the resuscitation; performs or supervises NIHSS; makes treatment eligibility decisions; reviews inclusion/exclusion checklist; obtains consent; communicates with family.
Stroke Team Nurse (Primary)	Stroke Unit / ED RN	Obtains vitals and IV access; draws STAT labs; prepares and administers thrombolytic per order; performs neuro-checks per monitoring schedule.
Recorder	Charge nurse or assigned RN	Timestamps every milestone on the Stroke Code Flow Sheet (activation, physician arrival, CT arrival/completion/read, needle time); ensures documentation is complete.
CT Technologist	Radiology on-call/on-duty tech	Prioritizes and immediately performs non-contrast CT ± CTA/CTP; notifies radiologist for urgent read.
Radiologist / Neuroradiologist	On-call Radiology	Provides urgent CT interpretation and communicates findings directly to the Team Leader by phone, not solely via written report.
Pharmacist	On-duty Pharmacy	Verifies weight-based thrombolytic dose; prepares medication on standby; double-checks dose prior to administration.
Laboratory Technologist	STAT lab on-duty	Prioritizes and expedites STAT stroke panel processing and result reporting.
Neurointerventionalist / Neurosurgery	On-call per roster	Evaluated emergently for large-vessel occlusion or hemorrhage requiring intervention (SU-003, SU-004, SU-014).
Charge Nurse (receiving unit)	Stroke Unit / ICU	Confirms bed availability and prepares for immediate patient transfer/admission.
Runner	Assigned support staff / aide	Physically expedites transport of specimens, blood products, or documents as needed.

6. Activation Roster and Contact Method

6.1 The current on-call roster for Neurology, Neurosurgery, and Neurointervention is maintained by the Stroke Program Coordinator and is available at all times through the hospital's on-call directory/paging system.

6.2 A single Stroke Code activation (call/page) simultaneously notifies all team roles listed in Section 5 through the hospital's group paging or clinical communication platform; no team member is notified sequentially or manually one-by-one.

6.3 If a team member does not acknowledge activation within 5 minutes, the charge nurse or Team Leader escalates through the on-call backup roster.

6.4 Team roster assignments (by role, not by name) are documented at the top of the Stroke Code Flow Sheet for every activation.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CODE STROKE TEAM — ROLES, RESPONSIBILITIES AND ACTIVATION ROSTER

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-010	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

7. References

- 7.1 AHA/ASA Target: Stroke® Best Practice Strategies — Organized Stroke Team Response.
- 7.2 The Joint Commission Comprehensive/Primary Stroke Center Certification Standards.

8. Attachments

- 8.1 Code Stroke Team Roster and Role Assignment Board Template.
- 8.2 Stroke Code Flow Sheet (Appendix C).
- 8.3 Stroke Code Post-Event Debrief Form.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE STROKE IMMEDIATE MANAGEMENT PATHWAY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-011	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-011 Acute Stroke Immediate Management Pathway (0–24 Hour Clinical Pathway)

1. Purpose

To provide a single, consolidated, time-sequenced clinical pathway summarizing the immediate management of any patient presenting with suspected acute stroke, from the point of first contact through the first 24 hours, integrating the detailed policies referenced throughout this manual into one quick-reference sequence for the bedside team.

2. Scope

This pathway applies to all suspected acute stroke patients, pre-hospital and in-hospital, and is intended to be used alongside — not in place of — the detailed policies SU-002 (Stroke Code Activation), SU-003 (Acute Ischemic Stroke Management), and SU-004 (Hemorrhagic Stroke Management).

3. Policy

- 3.1 Every patient with suspected acute stroke moves through the pathway below without deviation unless a specific clinical contraindication is documented by the Team Leader.
- 3.2 Pathway time targets mirror the AHA/ASA Target: Stroke® goals defined in SU-002, Section 3.4.
- 3.3 Any step that cannot be completed within its target window is documented on the Stroke Code Flow Sheet with the reason for delay, for review by the Stroke Committee (SU-009).

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE STROKE IMMEDIATE MANAGEMENT PATHWAY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-011	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

4. Immediate Management Pathway

MINUTE 0 — RECOGNITION
BE-FAST positive screen (pre-hospital or in-hospital) → ACTIVATE STROKE CODE



MINUTES 0–10 — ARRIVAL / TRIAGE
 Document exact Last Known Well (LKW) time • Vital signs + finger-stick glucose • IV access (18G preferred) • STAT labs drawn • Transfer to resuscitation/stroke bay • Team Leader begins focused history and NIHSS



MINUTES 10–25 — IMAGING
 Direct transport to CT • Non-contrast head CT ± CTA (arch to vertex) ± CTP • IV thrombolytic inclusion/exclusion checklist reviewed (Appendix B) • BP managed per algorithm (SU-012 / Appendix D or E)



DECISION POINT — CT RESULT



Hemorrhage identified	No hemorrhage — eligible for thrombolysis	No hemorrhage — not eligible / LVO suspected
Activate SU-004 pathway → Neurosurgery notified immediately → ICU admission → BP managed per hemorrhagic algorithm (Appendix E)	Proceed to thrombolytic administration within 60 minutes of arrival (goal ≤45 min) per SU-003 → monitor per post-tPA schedule	Obtain CTA/CTP if not already done → evaluate for mechanical thrombectomy (SU-014) up to 24 h from LKW



MINUTES 25–60+ — TREATMENT
 Thrombolytic administered and/or thrombectomy team activated • Continuous neuro and vital sign monitoring begins • Family/LAR informed and consent documented



HOURS 1–24 — STABILIZATION AND ADMISSION
 Admit to ICU or Stroke Unit per criteria (SU-017) • Repeat NIHSS at defined intervals • Dysphagia screen before any oral intake (SU-006) • VTE prophylaxis initiated • Continuous cardiac monitoring • Follow-up non-contrast CT at 24 h post-thrombolysis

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ACUTE STROKE IMMEDIATE MANAGEMENT PATHWAY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-011	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5. Escalation Triggers Throughout the Pathway

- 5.1 NIHSS increase ≥ 4 points or decline in level of consciousness — immediate physician notification and STAT CT.
- 5.2 Suspected symptomatic intracranial hemorrhage or systemic bleeding after thrombolytic — activate SU-013 Critical Actions immediately.
- 5.3 New large-vessel occlusion identified at any point — activate SU-014 Mechanical Thrombectomy Policy.
- 5.4 Any missed time target — document reason and notify the Stroke Coordinator for performance review (SU-009).

6. References

- 6.1 AHA/ASA 2019 Guidelines for the Early Management of Patients with Acute Ischemic Stroke.
- 6.2 AHA/ASA Target: Stroke® Phase III Initiative.

7. Attachments

- 7.1 Stroke Code Flow Sheet, Illustrated (Appendix C).
- 7.2 IV Thrombolytic Inclusion/Exclusion Checklist (Appendix B).

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: BLOOD PRESSURE MANAGEMENT IN ACUTE ISCHEMIC AND HEMORRHAGIC STROKE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-012	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-012 Blood Pressure Management in Acute Ischemic and Hemorrhagic Stroke

1. Purpose

To standardize blood pressure management targets and treatment algorithms for patients with acute ischemic stroke (with and without thrombolysis/thrombectomy) and for patients with acute hemorrhagic stroke (intracerebral or subarachnoid hemorrhage), consistent with current AHA/ASA guidelines.

2. Scope

This policy applies to all patients with confirmed or suspected acute stroke being managed in the Emergency Department, Stroke Unit, or ICU. It consolidates and standardizes the blood pressure provisions already referenced in SU-003 and SU-004 into a single dedicated policy with illustrated algorithms.

3. Policy — General Principles

3.1 Blood pressure targets differ fundamentally between ischemic and hemorrhagic stroke and by treatment received; the correct algorithm (Section 4 or 5) must be identified before any antihypertensive is administered.

3.2 Short-acting, titratable IV antihypertensive agents (labetalol, nicardipine, or clevidipine) are used in the acute phase; oral/long-acting agents are avoided until the patient is neurologically and hemodynamically stable.

3.3 Rapid, uncontrolled drops in blood pressure are avoided in all stroke subtypes, as hypotension can extend ischemic injury or reduce cerebral perfusion pressure.

3.4 All blood pressure treatment decisions and readings are documented on the Stroke Code Flow Sheet and subsequent nursing flow sheet at the frequency specified in Sections 4 and 5.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: BLOOD PRESSURE MANAGEMENT IN ACUTE ISCHEMIC AND HEMORRHAGIC STROKE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-012	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

4. Algorithm — Acute Ischemic Stroke

STEP 1: Is the patient a candidate for IV thrombolysis and/or mechanical thrombectomy?



YES — candidate for thrombolysis/thrombectomy	NO — not a candidate for acute reperfusion therapy
BEFORE treatment: lower BP to <185/110 mmHg using labetalol or nicardipine IV. If BP cannot be safely lowered below this threshold, thrombolysis is withheld. DURING/AFTER treatment (first 24 h): maintain BP <180/105 mmHg. Check BP every 15 min × 2 h, every 30 min × 6 h, then hourly × 16 h.	PERMISSIVE HYPERTENSION: withhold antihypertensive therapy unless SBP >220 mmHg or DBP >120 mmHg, or a compelling comorbid indication exists (ACS, aortic dissection, hypertensive encephalopathy, severe heart failure). If treatment is indicated, lower BP by approximately 15% during the first 24 hours — not to a normal target.



Any BP elevation with acute neurological deterioration, severe headache, or new nausea/vomiting → STAT non-contrast head CT to exclude hemorrhagic transformation before further BP treatment

5. Algorithm — Acute Hemorrhagic Stroke (ICH / SAH)

STEP 1: Confirm hemorrhage subtype on non-contrast CT — Intracerebral Hemorrhage (ICH) or Subarachnoid Hemorrhage (SAH)



Intracerebral Hemorrhage (ICH)	Subarachnoid Hemorrhage (SAH)
If SBP 150–220 mmHg: acute lowering to a target SBP of approximately 140 mmHg is reasonable and can be safe, using IV labetalol, nicardipine, or clevidipine. If SBP >220 mmHg: more aggressive reduction with continuous IV infusion and frequent (every 5–15 min) BP monitoring is reasonable, individualized to clinical context. Avoid precipitous drops; reassess and titrate continuously.	Control BP to reduce rebleeding risk prior to aneurysm securement (clipping/coiling), balanced against maintaining adequate cerebral perfusion pressure. Specific numeric targets are individualized with Neurosurgery/Neurocritical Care. Nimodipine 60 mg PO/NG every 4 hours is given for vasospasm prophylaxis (not for acute BP lowering) unless contraindicated by hypotension.



All ICH/SAH patients: continuous IV antihypertensive infusion with automated BP monitoring every 5–15 minutes during active titration, transitioning to every 30–60 minutes once at target — managed in the ICU

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: BLOOD PRESSURE MANAGEMENT IN ACUTE ISCHEMIC AND HEMORRHAGIC STROKE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-012	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

6. Documentation

Blood pressure readings, agent(s) used, and target achieved/not achieved are documented at the frequency specified above on the nursing flow sheet, and any deviation from target is flagged for physician reassessment.

7. References

- 7.1 Powers WJ, et al. 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke.
- 7.2 Greenberg SM, et al. 2022 AHA/ASA Guideline for the Management of Patients with Spontaneous Intracerebral Hemorrhage.
- 7.3 Hemphill JC, et al. AHA/ASA Guidelines for the Management of Aneurysmal Subarachnoid Hemorrhage.

8. Attachments

- 8.1 Blood Pressure Management Algorithm — Acute Ischemic Stroke, Illustrated (Appendix D).
- 8.2 Blood Pressure Management Algorithm — Acute Hemorrhagic Stroke, Illustrated (Appendix E).
- 8.3 IV Antihypertensive Agent Quick-Reference Dosing Card.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CRITICAL ACTIONS — SUSPECTED ICH / SYSTEMIC BLEEDING AFTER rtPA

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-013	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-013 Critical Actions — Suspected Intracranial Hemorrhage / Systemic Bleeding After IV Thrombolytic (rtPA) Administration

1. Purpose

To provide an immediate, standardized set of critical actions for the bedside team when a patient develops signs of symptomatic intracranial hemorrhage or systemic bleeding during or after administration of an IV thrombolytic agent (alteplase/tenecteplase), minimizing time to recognition, imaging, reversal, and specialist notification.

2. Scope

This policy applies to any patient who has received or is receiving IV thrombolytic therapy for acute ischemic stroke, in any inpatient location, and to all nursing and physician staff caring for such patients.

3. Definitions

3.1 Symptomatic Intracranial Hemorrhage (sICH): Hemorrhage on imaging associated with a clinical deterioration of ≥ 4 points on the NIHSS, or any decline attributable to the hemorrhage.

3.2 Warning Signs: Any acute neurological deterioration, severe new headache, acute hypertension, nausea, or vomiting occurring during or within 24 hours of thrombolytic infusion.

3.3 Systemic Bleeding: Clinically significant bleeding from any site (gastrointestinal, genitourinary, IV access sites, oropharyngeal, retroperitoneal) occurring during or after thrombolytic administration.

4. Policy

4.1 Any nurse identifying a Warning Sign (Section 3.2) immediately stops the thrombolytic infusion if still running, without waiting for a physician order.

4.2 A STAT non-contrast head CT and STAT coagulation panel (including fibrinogen) are obtained immediately upon recognition of a Warning Sign, without delay for other diagnostic steps.

4.3 Reversal/hemostatic therapy is initiated on physician order as soon as sICH or systemic bleeding is confirmed or strongly suspected, without waiting for full imaging report finalization when clinical suspicion is high.

4.4 Neurosurgery is notified immediately for any confirmed intracranial hemorrhage following thrombolysis.

4.5 Every activation of this policy is reviewed at the next Stroke Committee meeting as part of the thrombolytic complication rate metric (SU-009).

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CRITICAL ACTIONS — SUSPECTED ICH / SYSTEMIC BLEEDING AFTER rtPA

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-013	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5. Critical Action Sequence

RECOGNITION

Acute neuro decline • severe new headache • acute hypertension • nausea/vomiting • new bleeding at any site during/after rtPA



IMMEDIATE ACTIONS (Target: within 5 minutes of recognition)

- 1 .STOP the thrombolytic infusion if still running
- 2 .Call for physician / activate Rapid Response if not already at bedside
- 3 .Reassess airway, breathing, circulation and full vital signs
- 4 .Perform focused NIHSS to quantify any new deficit



STAT DIAGNOSTICS (Target: within 15 minutes)

STAT non-contrast head CT • STAT labs: CBC, PT/INR, aPTT, fibrinogen, type & crossmatch • Notify Neurosurgery and Stroke Team physician



REVERSAL / HEMOSTATIC THERAPY (per physician order, initiated emergently)

Cryoprecipitate (10 units) to replace fibrinogen — target fibrinogen ≥ 150 mg/dL

Tranexamic acid or ϵ -aminocaproic acid per organizational reversal protocol

Platelet transfusion if thrombocytopenic or recent antiplatelet use per physician judgment

For systemic (non-intracranial) bleeding: local hemostatic measures, direct pressure, and transfusion support as indicated



ESCALATION AND DISPOSITION

Any confirmed intracranial hemorrhage → immediate ICU admission and Neurosurgery evaluation for surgical candidacy per SU-004

Blood pressure managed per the Hemorrhagic Stroke Algorithm (SU-012, Appendix E)

Family/LAR notified promptly of the change in clinical status

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: CRITICAL ACTIONS — SUSPECTED ICH / SYSTEMIC BLEEDING AFTER rtPA

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-013	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

6. Documentation

All critical actions, timestamps, medications given, and repeat imaging results are documented on the Critical Action Card (Appendix F) and in the medical record, and the event is entered into the stroke complication log for Stroke Committee review.

7. References

- 7.1 Powers WJ, et al. 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke — Management of Thrombolytic Complications.
- 7.2 Organizational Anticoagulation/Thrombolytic Reversal Protocol.

8. Attachments

- 8.1 Critical Action Card — Suspected ICH / Systemic Bleeding After rtPA, Illustrated (Appendix F).
- 8.2 Thrombolytic Complication Reporting Form.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: MECHANICAL THROMBECTOMY POLICY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-014	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-014 Mechanical Thrombectomy Policy

1. Purpose

To establish a dedicated policy governing patient selection, notification workflow, procedural coordination, and post-procedure care for mechanical thrombectomy (endovascular therapy) in acute ischemic stroke due to large vessel occlusion (LVO), consistent with current AHA/ASA guidelines and, where applicable, an institutional transfer agreement with a thrombectomy-capable partner facility.

2. Scope

This policy applies to all patients presenting with suspected LVO acute ischemic stroke, whether treated on-site by a certified neurointerventional service or transferred to a designated thrombectomy-capable/comprehensive stroke center, and to all staff involved in identification, notification, transfer, procedural support, and post-procedure monitoring.

3. Definitions and Abbreviations

3.1 LVO: Large Vessel Occlusion — occlusion of the internal carotid artery, M1/proximal M2 middle cerebral artery, basilar artery, or other major intracranial vessel.

3.2 ASPECTS: Alberta Stroke Program Early CT Score.

3.3 Core/Penumbra Mismatch: A perfusion imaging pattern (CT perfusion or MRI) identifying a small infarct core relative to a larger volume of at-risk but salvageable tissue, used to select extended-window candidates.

3.4 Door-to-Groin-Puncture Time: The interval from hospital arrival to first arterial access for the thrombectomy procedure.

4. Policy — Patient Selection

4.1 Patients are evaluated for mechanical thrombectomy if they have: confirmed anterior circulation LVO on CTA, NIHSS ≥ 6 , pre-stroke independent functional status (modified Rankin Scale 0–1), and treatment can begin within 6 hours of LKW, OR within 6–24 hours if favorable core/penumbra mismatch criteria are met on CT perfusion or MRI.

4.2 Favorable imaging profile for the 0–6-hour window is generally defined as ASPECTS ≥ 6 ; extended-window (6–24 hour) eligibility uses validated automated perfusion software applying the core infarct volume and clinical mismatch thresholds from the pivotal randomized trials referenced in current AHA/ASA guidance.

4.3 Basilar artery occlusion is evaluated for thrombectomy on a case-by-case basis with Neurointervention/Neurosurgery, given an extended potential treatment window relative to anterior circulation LVO.

4.4 IV thrombolysis, when the patient is eligible and within window, is administered without delay and is never withheld solely to expedite transfer or thrombectomy (SU-003, Section 6.3).

5. Policy — Notification and Transfer Workflow

5.1 The Team Leader notifies the on-call Neurointerventionalist (or Neurosurgery, per institutional roster) immediately upon identification of a suspected LVO on CTA, in parallel with — not after — other Stroke Code workflow steps.

5.2 On-site capability: the neurointerventional suite/team is activated and the patient is transported directly from CT to the angiography suite once consent is obtained.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: MECHANICAL THROMBECTOMY POLICY

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-014	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5.3 No on-site capability: an emergent transfer is initiated to the designated thrombectomy-capable/Comprehensive Stroke Center per the organization's stroke transfer agreement; imaging is transmitted electronically in advance, and EMS/transfer team departure is not delayed pending non-essential paperwork.

5.4 Door-to-groin-puncture time targets: within 90 minutes of arrival for direct-presentation patients, within 60 minutes of arrival at the thrombectomy-capable center for transfer patients, consistent with SU-002, Section 3.4.5.

6. Policy — Procedural and Post-Procedure Care

6.1 Informed consent for the procedure is obtained from the patient or legally authorized representative and documented per organizational policy for emergency/time-critical procedures.

6.2 Anesthesia approach (conscious sedation vs. general anesthesia) is determined by the proceduralist based on patient stability, airway status, and ability to remain still for the procedure.

6.3 Successful reperfusion is graded using the modified Thrombolysis in Cerebral Infarction (mTICI) scale and documented in the procedure note.

6.4 Post-procedure patients are admitted to the ICU or Stroke Unit for neurological monitoring and access-site checks (groin/radial site, distal pulses, signs of hematoma) per the post-procedure monitoring schedule.

6.5 Blood pressure post-thrombectomy is managed per SU-012, individualized to reperfusion status (successful vs. incomplete) and hemorrhage risk, per Neurointervention/Neurology direction.

6.6 A follow-up non-contrast head CT is obtained at 24 hours post-procedure, or sooner if any neurological change occurs, to assess for hemorrhagic transformation.

7. Admission Criteria Post-Thrombectomy

All patients are admitted to the ICU for at least the initial post-procedure period per SU-017 (ICU Admission Criteria), with step-down to the Stroke Unit once neurologically and hemodynamically stable and the access site is confirmed intact.

8. References

8.1 Powers WJ, et al. 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke.

8.2 Nogueira RG, et al. and Albers GW, et al. — pivotal extended-window mechanical thrombectomy trials referenced in current AHA/ASA guidance.

8.3 The Joint Commission Comprehensive Stroke Center / Thrombectomy-Capable Stroke Center Certification Standards.

9. Attachments

9.1 Mechanical Thrombectomy Candidate Checklist.

9.2 Mechanical Thrombectomy Transfer Checklist and Transfer Agreement Summary.

9.3 Post-Thrombectomy Neurological and Access-Site Monitoring Flow Sheet.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: NURSING AND STAFF STROKE EDUCATION PROGRAM

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-015	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-015 Nursing and Staff Stroke Education Program

1. Purpose

To ensure that all nursing and clinical staff involved in stroke care receive structured, competency-based orientation and ongoing education, consistent with AHA/ASA guidelines and The Joint Commission requirement for documented, annual stroke-specific staff education.

2. Scope

This policy applies to all registered nurses, emergency department staff, ICU staff, and other clinical personnel who provide direct care to stroke patients, and to physicians participating in the Code Stroke Team for role-specific competency elements.

3. Policy

- 3.1 All new nursing staff assigned to the Emergency Department, Stroke Unit, or ICU complete stroke-specific orientation prior to independently caring for a stroke patient, covering the topics in Section 4.
- 3.2 All staff who perform NIHSS assessments complete formal NIHSS certification prior to independent scoring, with recertification at minimum every 2 years, per SU-006.
- 3.3 All Stroke Unit and Code Stroke Team nursing staff complete annual stroke competency validation, documented in the individual's education/competency file.
- 3.4 Education content is reviewed and updated at minimum every two years, or sooner with any significant AHA/ASA guideline update, by the Stroke Program Coordinator in coordination with Nursing Education.
- 3.5 Staff education compliance rates are tracked and reported to the Stroke Committee at minimum quarterly (SU-009).

4. Core Orientation Curriculum

- 4.1 Stroke recognition (BE-FAST) and the Stroke Code activation process (SU-002).
- 4.2 NIHSS assessment technique and certification.
- 4.3 IV thrombolytic preparation, administration, and post-administration monitoring, including recognition of and response to complications (SU-003, SU-013).
- 4.4 Blood pressure management algorithms for ischemic and hemorrhagic stroke (SU-012).
- 4.5 Dysphagia screening technique and NPO precautions (SU-006).
- 4.6 Neurological assessment and early recognition of deterioration, including hemorrhagic transformation and increased intracranial pressure.
- 4.7 VTE prophylaxis, early mobilization principles, and pressure-injury prevention specific to the stroke patient.
- 4.8 Patient/family stroke education delivery and use of the Stroke Education Booklet (SU-016).
- 4.9 Documentation requirements, including the Stroke Code Flow Sheet and registry data elements.

5. Ongoing / Annual Education

- 5.1 Annual skills validation (return demonstration or simulation) for thrombolytic administration and management of a simulated acute neurological change.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: NURSING AND STAFF STROKE EDUCATION PROGRAM

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-015	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5.2 Case-based review of de-identified complications, near-misses, or performance gaps identified through the Stroke Performance Improvement program (SU-009).

5.3 Updates on any changes to AHA/ASA guidelines, order sets, or institutional protocols.

5.4 Mock Stroke Code drills conducted at a minimum semi-annually on units with Stroke Code responsibility, with debrief and documented feedback.

6. Documentation

Completion of orientation, NIHSS certification/recertification, and annual competency validation is documented in each staff member's education file and is auditable by Nursing Education, the Stroke Program Coordinator, and TJC surveyors.

7. References

7.1 The Joint Commission Comprehensive/Primary Stroke Center Certification Standards — Staff Education Requirements.

7.2 National Institutes of Health Stroke Scale (NIHSS) Training and Certification materials.

8. Attachments

8.1 Stroke Orientation Checklist (New Staff).

8.2 Annual Stroke Competency Validation Form.

8.3 Mock Stroke Code Drill and Debrief Template.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: PATIENT AND CAREGIVER STROKE EDUCATION PROGRAM

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-016	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-016 Patient and Caregiver Stroke Education Program

1. Purpose

To ensure every stroke and TIA patient, together with their family/caregiver, receives structured, understandable, and documented education throughout the hospital stay and prior to discharge, supporting safe self-management, adherence to secondary prevention, and prompt recognition of recurrent symptoms.

2. Scope

This policy applies to all patients admitted with a diagnosis of ischemic stroke, hemorrhagic stroke, or TIA, and to their identified family members/caregivers, across the Stroke Unit, ICU, and any unit providing stroke care. It expands on the discharge education topics summarized in SU-008.

3. Policy

3.1 Stroke education begins at admission and continues throughout the hospitalization, delivered progressively rather than only at the point of discharge.

3.2 Education is delivered in the patient's/caregiver's preferred language using professional interpreter services when needed, and in a format appropriate to the patient's cognitive, visual, and communication status (accounting for aphasia or dysphagia).

3.3 All required core education topics (Section 4) are covered and documented prior to discharge using the Stroke Education Booklet and Documentation Form; when the patient cannot participate due to cognitive impairment, education is directed to the identified caregiver/family member.

3.4 Teach-back technique is used to confirm patient/caregiver understanding of medications, warning signs, and follow-up plans.

3.5 Documentation of stroke education is a required element of the discharge summary and is audited as part of the GWTG-Stroke/Joint Commission STK-8 education measure (SU-009).

4. Core Patient/Caregiver Education Topics

4.1 What happened: a plain-language explanation of the type of stroke/TIA and the affected brain territory.

4.2 Personal risk factors and individualized modification strategies (hypertension, diabetes, dyslipidemia, atrial fibrillation, smoking, physical inactivity, obesity).

4.3 Warning signs and symptoms of stroke (BE-FAST) and the critical importance of calling emergency services immediately if symptoms recur — never to "wait and see."

4.4 Prescribed medications: name, purpose, dose, schedule, common side effects, and the specific importance of adherence to antithrombotic/anticoagulant therapy.

4.5 Follow-up care: scheduled appointments, contact information for questions, and when to seek urgent versus emergency care.

4.6 Activity, dietary, and lifestyle recommendations, including dysphagia diet modifications and safe swallowing strategies if applicable.

4.7 Fall prevention and home safety modifications relevant to the patient's residual deficits.

4.8 Caregiver-specific education: safe transfer and mobility assistance techniques, recognizing caregiver strain, and available respite/support resources.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: PATIENT AND CAREGIVER STROKE EDUCATION PROGRAM

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-016	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

4.9 Community resources, stroke support groups, and how to access them.

5. Delivery Methods

5.1 One-on-one bedside teaching by the bedside nurse, reinforced by the multidisciplinary team (pharmacy for medications, dietitian for diet modification, rehabilitation team for functional strategies).

5.2 Written, plain-language Stroke Education Booklet provided to every patient/caregiver, available in the primary languages served by the organization.

5.3 Visual aids (BE-FAST posters, medication schedule cards) used to reinforce key safety messages for patients with communication or cognitive deficits.

5.4 A scheduled discharge education session, with the caregiver present whenever possible, conducted using teach-back prior to discharge.

6. Documentation

Topics covered, materials provided, language used, and teach-back confirmation are documented on the Stroke Education Booklet and Documentation Form, filed in the medical record, and summarized in the discharge instructions.

7. References

7.1 AHA/ASA Get with the Guidelines®—Stroke Education Measure Specifications (STK-8).

7.2 Kleindorfer DO, et al. 2021 AHA/ASA Guideline for the Prevention of Stroke in Patients with Stroke and Transient Ischemic Attack.

8. Attachments

8.1 Patient/Family Stroke Education Booklet and Documentation Form.

8.2 BE-FAST Warning Signs Patient Handout.

8.3 Caregiver Support and Community Resources Directory.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ICU ADMISSION CRITERIA FOR STROKE PATIENTS

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-017	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

SU-017 Intensive Care Unit Admission Criteria for Stroke Patients

1. Purpose

To consolidate, in a single dedicated policy, the criteria governing ICU admission for patients with acute ischemic stroke, hemorrhagic stroke, and post-procedural (thrombolysis/thrombectomy) patients, ensuring consistent triage between ICU, Stroke Unit, and monitored-bed levels of care.

2. Scope

This policy applies to all admission and level-of-care decisions for stroke and TIA patients and is used together with the specific admission provisions already referenced in SU-003, SU-004, SU-005, and SU-014.

3. Policy — Mandatory ICU Admission Criteria

- 3.1 Any patient who received IV thrombolysis, for at least the initial monitoring period per the post-tPA monitoring schedule (SU-003, Section 5.4).
- 3.2 Any patient who underwent mechanical thrombectomy, for at least the initial post-procedure period (SU-014, Section 7).
- 3.3 All confirmed intracerebral hemorrhage (ICH) or subarachnoid hemorrhage (SAH) patients, for the duration of active bleeding risk and neurosurgical monitoring (SU-004, Section 6.1).
- 3.4 Acute stroke with depressed level of consciousness (GCS ≤ 12 or any acute decline in consciousness).
- 3.5 Stroke with new-onset seizures.
- 3.6 Hemodynamic instability (requiring vasopressor/inotropic support or continuous IV antihypertensive titration outside standard post-tPA monitoring) or respiratory instability (requiring airway protection, non-invasive or invasive ventilation).
- 3.7 Large territory infarction with anticipated risk of malignant cerebral edema (e.g., large MCA or cerebellar infarction), for neurological monitoring and potential decompressive intervention.
- 3.8 Basilar artery occlusion pending or following intervention, given risk of rapid neurological decline.

4. Policy — Stroke Unit (Non-ICU Monitored) Admission Criteria

- 4.1 Hemodynamically and neurologically stable acute ischemic stroke not meeting any ICU criterion in Section 3.
- 4.2 Stable TIA patients admitted for expedited work-up per SU-005 admission criteria.
- 4.3 Post-tPA or post-thrombectomy patients who have completed the mandatory ICU/high-dependency monitoring period and are neurologically and hemodynamically stable, per step-down assessment by the treating physician.

5. Policy — Continuous Reassessment and Escalation

- 5.1 Level-of-care status is reassessed at minimum every nursing shift and with any change in neurological or hemodynamic status; it is not a one-time admission decision.
- 5.2 Any patient on the Stroke Unit who develops a new focal deficit, GCS decline, need for airway protection, hemorrhagic transformation, or hemodynamic instability is escalated to ICU immediately, without waiting for the next scheduled physician round.

King Fahad Central Hospital — STROKE PROGRAM

DEPARTMENT POLICY AND PROCEDURE: ICU ADMISSION CRITERIA FOR STROKE PATIENTS

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
SU-017	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide		

5.3 Conversely, ICU patients who no longer meet any criterion in Section 3 and have been stable for a physician-determined observation period are transferred to the Stroke Unit to support efficient use of critical care resources.

6. Documentation

The admitting level of care and the specific criterion/criteria met are documented in the admission note, and any escalation or de-escalation event is documented with the triggering finding and time.

7. References

- 7.1 Powers WJ, et al. 2019 AHA/ASA Guidelines for the Early Management of Patients with Acute Ischemic Stroke.
- 7.2 Greenberg SM, et al. 2022 AHA/ASA Guideline for the Management of Patients with Spontaneous Intracerebral Hemorrhage.
- 7.3 The Joint Commission Comprehensive/Primary Stroke Center Certification Standards.

8. Attachments

- 8.1 ICU / Stroke Unit Level-of-Care Triage Checklist.

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: NIH STROKE SCALE (NIHSS) SCORING REFERENCE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix A	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix A — NIH Stroke Scale (NIHSS): Full Item Scoring Reference

The NIH Stroke Scale (NIHSS) is administered by NIHSS-certified staff at the intervals defined in SU-006. Score each item in the order listed; do not go back and change scores. Score what the patient does, not what the examiner thinks the patient can do.

Item	Tests	Score / Description
1a	Level of Consciousness (LOC)	0 Alert 1 Not alert, arousable by minor stimulation 2 Not alert, requires repeated stimulation 3 Unresponsive / reflex responses only
1b	LOC Questions (month, age)	0 Both correct 1 One correct 2 Neither correct
1c	LOC Commands (open/close eyes, grip/release)	0 Both correct 1 One correct 2 Neither correct
2	Best Gaze	0 Normal 1 Partial gaze palsy 2 Forced deviation
3	Visual Fields	0 No visual loss 1 Partial hemianopia 2 Complete hemianopia 3 Bilateral hemianopia
4	Facial Palsy	0 Normal 1 Minor paralysis 2 Partial paralysis 3 Complete paralysis
5a / 5b	Motor Arm (Left / Right)	0 No drift 1 Drift 2 Some effort against gravity 3 No effort against gravity 4 No movement UN Amputation/joint fusion
6a / 6b	Motor Leg (Left / Right)	0 No drift 1 Drift 2 Some effort against gravity 3 No effort against gravity 4 No movement UN Amputation/joint fusion
7	Limb Ataxia	0 Absent 1 Present in one limb 2 Present in two limbs
8	Sensory	0 Normal 1 Mild-moderate loss 2 Severe to total loss
9	Best Language	0 No aphasia 1 Mild-moderate aphasia 2 Severe aphasia 3 Mute / global aphasia
10	Dysarthria	0 Normal 1 Mild-moderate 2 Severe UN Intubated/other barrier
11	Extinction / Inattention (Neglect)	0 No abnormality 1 Inattention/extinction to one modality 2 Profound hemi-inattention to more than one modality

Scoring Summary and Severity Bands

Total score range: 0–42 (sum of all items, treating UN items per NIHSS instructions).

0: No stroke symptoms. 1–4: Minor stroke. 5–15: Moderate stroke. 16–20: Moderate to severe stroke. 21–42: Severe stroke.

A sustained increase of ≥ 4 points from baseline, or any acute decline in level of consciousness, triggers immediate physician notification per SU-006, Section 2.3.

Full item-by-item instructions, standardized phrases, and picture cards are provided in the official NIHSS training and certification materials and are not reproduced in full here; staff must complete formal certification before independent use (SU-015).

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: IV THROMBOLYTIC INCLUSION/EXCLUSION CHECKLIST

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix B	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix B — IV Thrombolytic (Alteplase/Tenecteplase) Inclusion/Exclusion Checklist

This checklist is completed and signed by the responding physician for every patient being considered for IV thrombolytic therapy, and is filed with the Stroke Code Flow Sheet. It supplements, and does not replace, current AHA/ASA guidance and physician clinical judgment.

Section 1 — Inclusion Criteria (all must be YES)

☐	Criterion
☐	Clinical diagnosis of ischemic stroke causing a measurable neurological deficit
☐	Time of symptom onset / Last Known Well clearly established
☐	Treatment can be initiated within 4.5 hours of LKW (standard window) or patient meets extended-window imaging criteria per SU-003, Section 7
☐	Age ≥18 years

Section 2 — Exclusion Criteria (any YES excludes thrombolysis unless overridden by documented physician judgment)

☐	Criterion
☐	Evidence of intracranial hemorrhage on non-contrast CT
☐	Blood pressure remains ≥185/110 mmHg despite treatment
☐	Blood glucose <50 mg/dL with symptoms attributable to hypoglycemia rather than stroke
☐	Platelet count <100,000/mm ³
☐	Current anticoagulant use with INR >1.7, or elevated PT/aPTT, or therapeutic-dose LMWH within 24 hours
☐	Use of a direct thrombin/factor Xa inhibitor within 48 hours with elevated sensitive coagulation assays (unless assays are normal / agent held ≥48 h with normal renal function)
☐	Prior intracranial hemorrhage, known structural CNS lesion (aneurysm, AVM, neoplasm), or intracranial/spinal surgery within 3 months
☐	Active internal bleeding or acute trauma/fracture on presentation
☐	Recent (within 3 months) severe head trauma, or within 21 days GI/GU hemorrhage
☐	Major surgery or serious trauma within 14 days
☐	Suspected or confirmed infective endocarditis, or suspected aortic dissection
☐	Presentation consistent with subarachnoid hemorrhage even with a normal CT
☐	CT shows extensive regions of clear hypodensity/established infarction

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: IV THROMBOLYTIC INCLUSION/EXCLUSION CHECKLIST

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix B	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Section 3 — Extended-Window (3–4.5 hour) Additional Exclusion Considerations

Age >80 years. Oral anticoagulant use regardless of INR. Baseline NIHSS >25. Combined history of prior stroke and diabetes mellitus. Imaging evidence of infarction involving more than one-third of the MCA territory.

Section 4 — Physician Attestation

I have reviewed the above inclusion and exclusion criteria with the available clinical and imaging information. Based on this review, the patient is / is not a candidate for IV thrombolytic therapy. Risks and benefits have been discussed with the patient/legally authorized representative and informed consent has been documented.

Physician Signature		Date / Time
---------------------	--	-------------

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: STROKE CODE / ACUTE STROKE FLOW SHEET, ILLUSTRATED

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix C	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix C — Stroke Code / Acute Stroke Flow Sheet, Illustrated

This flow sheet provides a single illustrated timeline for real-time documentation of every Stroke Code milestone by the Recorder (SU-010, Section 5), from activation through disposition.


Milestone	Target	Time Recorded
Last Known Well (LKW)	—	
Stroke Code activation	T = 0	
Patient arrival (or in-hospital recognition)	—	
Physician / Team Leader at bedside	≥10 min from arrival	
Vital signs, glucose, IV access obtained	≥10 min from arrival	
STAT labs drawn	≥10 min from arrival	
Patient arrives in CT	≥20 min from arrival	
CT completed	≥25 min from arrival	
CT interpreted (verbal read)	≥45 min from arrival	
IV thrombolytic checklist completed	before needle time	
Thrombolytic bolus given ("Needle Time")	≥60 min from arrival (goal ≤45 min)	
Neurointervention/Neurosurgery notified (if LVO/hemorrhage)	immediate upon CT result	
Groin puncture (if thrombectomy)	≥90 min direct / ≤60 min transfer	
Disposition decision (ICU / Stroke Unit)	≥3 h from arrival	

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: STROKE CODE / ACUTE STROKE FLOW SHEET, ILLUSTRATED

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix C	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Illustrated Timeline



Team Roster Recorded This Activation


Role	Name / ID
Team Leader	
Stroke Team Nurse	
Recorder	
CT Technologist	
Pharmacist	

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: BP MANAGEMENT ALGORITHM — ACUTE ISCHEMIC STROKE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix D	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix D — Blood Pressure Management Algorithm: Acute Ischemic Stroke (Illustrated)




King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: BP MANAGEMENT ALGORITHM — ACUTE HEMORRHAGIC STROKE

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix E	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix E — Blood Pressure Management Algorithm: Acute Hemorrhagic Stroke (Illustrated)




King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: CRITICAL ACTION CARD — ICH / BLEEDING AFTER rtPA

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix F	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Appendix F — Critical Action Card: Suspected ICH / Systemic Bleeding After rtPA (Illustrated)

This card is posted at the bedside of every patient receiving IV thrombolytic therapy and is used as an immediate reference during the critical action sequence defined in SU-013.



Quick-Reference Contact List

Role	Contact Method
Stroke Team Physician / Team Leader	
Neurosurgery on-call	
Blood Bank / Transfusion Services	
Pharmacy	

King Fahad Central Hospital — STROKE PROGRAM

STROKE PROGRAM REFERENCE: CRITICAL ACTION CARD — ICH / BLEEDING AFTER rtPA

POLICY NUMBER	REPLACES NUMBER	EFFECTIVE DATE	REVIEW DATE
Appendix F	—	]Effective Date[	]Review Date[
APPROVED BY	Stroke Medical Director / Quality Department		
APPLIES TO	Hospital Wide — Clinical Reference / Job Aid		

Document Control and Revision History

This manual is a controlled document owned by the Stroke Program Coordinator and approved by the Stroke Committee (Interprofessional Committee) and Medical Director. It is reviewed at minimum annually, and immediately upon any material change in evidence-based guidelines, regulatory requirements, or certification standards. All revisions are version-controlled and archived; the current approved version supersedes all prior versions upon publication.

Program requirements throughout this manual are scaled to the certification level held or pursued by the organization — Acute Stroke Ready Center (ASRC), Primary Stroke Center (PSC), or Comprehensive Stroke Center (CSC) — with certification-level-specific requirements noted within the applicable policy section.

VERSION	DATE	SUMMARY OF CHANGES	APPROVED BY
1.0	]Insert Date[	Initial issue of manual.	]Name/Title[

Manual Approval

The undersigned confirm review and approval of the Stroke Unit Policy and Procedure Manual (SU-001 through SU-017, plus Appendices A–F) for implementation.

ROLE	SIGNATURE	DATE
Stroke Program Medical Director		
Chief of Neurology		
Emergency Department Medical Director		
Chief Nursing Officer		
Stroke Program Coordinator		
Director of Rehabilitation Services		
Director of Pharmacy		
Director of Radiology		
Chief Medical Officer / Medical Director		