
Figure 3. Study diagram for study 20190232



- Subjects with body weight at baseline ≤100kg receive 45mg SC; >100kg receive 90 mg SC
- At week 28, non-responders (ie, subjects not achieving PASI 50 response of better) will be considered to have completed the study and will complete the end of study procedures (ie, week 52 procedures) at week 28. Subjects missing the week 28 visit or missing the PASI assessment at week 28, will be discontinued from the study.
- *For subjects with PASI 50 response or better but less than PASI 75 response at week 28, based on investigator's discretion, subjects may increase dose frequency to every 8 weeks (Q8W). Subjects with body weight at baseline ≤ 100kg will receive 45mg Q8W and subjects > 100kg at baseline will receive 90mg Q8W at weeks 28, 36, and 44. These subjects will continue on original treatment and will NOT be re-randomized.

PASI = Psoriasis Area and Severity Index; Q8W = every 8 weeks; Q12W = every 12 weeks; SC = subcutaneous

Objectives

The primary objective of the study was to compare the efficacy of ABP 654 with ustekinumab in subjects with moderate to severe Ps.

The secondary objective was to assess the safety and immunogenicity of ABP 654 compared with ustekinumab.

Null Hypothesis (H_0): Mean difference in PASI percent improvement from baseline at week 12 between ABP 654 and ustekinumab is outside an equivalence margin of (-15, 15),

versus

Alternative Hypothesis (H_A): Mean difference in PASI percent improvement from baseline at week 12 between ABP 654 and ustekinumab is within an equivalence margin of (-15, 15).

Outcomes/endpoints

The primary efficacy endpoint is the Psoriasis Area and Severity Index (PASI) percent improvement from baseline to week 12.

Secondary Endpoints