
PK Similarity Assessment

PK similarity between ABP 654, US-Stelara and EU-Stelara has been demonstrated in a 3-arm, parallel PK similarity study in healthy subjects (Study 20190230). PK samples were collected pre-dose on Day 1, and post-dose at 8 hr and 24 hr and on Days 3, 7, 9, 11, 13, 21, 28, 35, 49, 56, 70, 98 and 112. The point estimates and 90% CIs of the GMRs of PK parameters ($AUC_{0\text{-}inf}$, C_{max} and AUC_{last}) were all within the pre-defined criteria of 0.8 to 1.25 (Table 8). The mean serum concentration-time profiles were similar between the ABP 654, US-Stelara and EU-Stelara treatment groups (Figure 2). The PK parameters following a single dose of ABP 654, US-Stelara and EU-Stelara are summarized in Table 8. In general, PK parameters and inter-subject variability were comparable across the three treatment groups.

Figure 2. Mean (±SD) Concentrations Over Time by Treatment (Study 20190230)



Source: FDA Analysis

Table 8. Summary of PK Parameters (Study 20190230)

PK Parameters	Geometric Mean (Geometric CV%)		
	ABP 654 (n=78)	US-Stelara (n=79)	EU-Stelara (n=80)
C_{max} (ng/mL)	11546 (36.1)	11258 (33.8)	12085 (32.8)
AUC_{last} (h*ng/mL)	10175553 (39.6)	9784499 (38.5)	10684286 (33.5)
$AUC_{0\text{-}inf}$ (h*ng/mL)	10859684 (40.9)	10592020 (38.5)	11310320 (35.3)
T_{max} (h)[a]	144 (24, 288)	192 (48, 648)	192 (48, 648)