

Study CT-P43 1.1 Part 2

The analyses for maximum serum concentration ($C_{\max}$) and area under the concentration-time curve from time zero to infinity ($AUC_{0-\infty}$) by post-treatment ADA status regardless of pre-dose result are presented for the PK Set in the below Table.

The $AUC_{0-\infty}$ was lower in ADA positive subjects compared to ADA negative subjects in all treatment groups, while there was little impact of ADA positivity on $C_{\max}$.

Table 15. Impact of ADA positivity (post-treatment ADA status regardless of pre-dose result) on $C_{\max}$ and $AUC_{0-\infty}$ in Study CT-P43 1.1 Part 2 (PK Set)

Parameter Statistic	CT-P43 (N=80)		EU-Stelara (N=81)		US-Stelara (N=77)	
	ADA Negative	ADA Positive	ADA Negative	ADA Positive	ADA Negative	ADA Positive
$C_{\max}$ ($\mu\text{g/mL}$)						
n	67	13	57	24	43	34
Mean	5.37	5.29	4.48	4.78	4.85	4.88
Min	2.1	2.5	2.2	2.1	1.1	2.3
Max	9.6	9.5	7.2	7.6	8.3	8.3
$AUC_{0-\infty}$ ($\text{day} \cdot \mu\text{g/mL}$)						
n	66	13	57	24	43	34
Mean	209.70	177.66	167.05	153.27	177.22	151.48
Min	90.5	100.2	84.8	65.4	61.4	39.3
Max	363.6	290.8	256.7	277.8	325.3	287.4

In addition, the post-hoc analyses for $C_{\max}$ and $AUC_{0-\infty}$ by ADA titre separated by ADA high and low, are presented in Table below. Highest post-treatment ADA titres of individual subjects were used in the analyses. Impact of ADA titre on $AUC_{0-\infty}$ was observed while there was no notable impact on $C_{\max}$.