

	Bmab 1200 (N=191) n (%)	Stelara (N=193) n (%)	Total (N=384) n (%)
Smoker Status			
Never Smoked	108 (56.5)	122 (63.2)	230 (59.9)
Ex-smoker	22 (11.5)	19 (9.8)	41 (10.7)
Smoker	61 (31.9)	52 (26.9)	113 (29.4)
Unknown	0	0	0
Alcohol Status			
Non-drinker	116 (60.7)	119 (61.7)	235 (61.2)
Ex-drinker	1 (0.5)	0	1 (0.3)
Current drinker	74 (38.7)	73 (37.8)	147 (38.3)

Abbreviations: FAS, full analysis set; N, number of patients in the treatment group; n, number of patients with available data; Q1, 1st quartile; Q2, 2nd quartile; Q3, 3rd quartile.

Note: Percentages are based on the number of patients in the treatment group (N).

Note: Baseline is defined as the last non-missing value before the first dose of study treatment.

^a. Percentages (%) are based on the number of female patients in each treatment group.

^b. Percentages (%) are based on the number of current drinkers in each treatment group.

Baseline Disease Characteristics

The mean (SD) PASI score, sPGA, BSA, and DLQI was 23.2 (9.2), 3.6 (0.7), 29.9 (16.3), and 14.4 (6.7), respectively. The majority of patients had an sPGA score of 3 (205 patients; 53.4%) or 4 (125 patients; 32.6%). sPGA scores of 4 or 5 were observed in slightly more patients in the Bmab1200 group than the Stelara group. The majority of patients did not have previous exposure to biologic-based therapies (331 patients; 86.2%) and did not have concomitant psoriatic arthritis (322 patients; 83.9%).