

Light drinker (less than 2 drinks per day)	73 (38.2)	71 (36.8)	144 (37.5)
Moderate drinker (2-4 drinks per day)	1 (0.5)	2 (1.0)	3 (0.8)
Ex-drinker	1 (0.5)	0	1 (0.3)
Non drinker	116 (60.7)	119 (61.7)	235 (61.2)
Unknown	0	1 (0.5)	1 (0.3)
Previous exposure to biologic-based therapies			
No prior exposure to biologics	165 (86.4)	166 (86.0)	331 (86.2)
Prior exposure to biologics	26 (13.6)	27 (14.0)	53 (13.8)
Concomitant psoriatic arthritis			
Concomitant psoriatic arthritis	30 (15.7)	32 (16.6)	62 (16.1)
No Concomitant psoriatic arthritis	161 (84.3)	161 (83.4)	322 (83.9)

Source: Reviewer's analysis

SD = Standard Deviation.

Other Product-Specific Safety Concerns

Deaths

There were no deaths reported in BM12H-PSO-03-G-02.

Treatment Emergent Adverse Events

Serious adverse events (SAE)s

Overall, 7 subjects (1.8%) experienced 9 serious treatment emergent adverse events (TEAEs) through the study.

In TP1, four subjects reported 5 SAEs, none of which was considered related to study drug treatment.

Table 33. SAEs in Subjects with Psoriasis in TP1 (Up to Week 16)

Preferred Term	Bmab 1200	EU-Stelara
	N = 191 n (%)	N = 193 n (%)
Any SAE	3 (1.6)	1 (0.5)
Endometrial adenocarcinoma	1 (0.5)	0 (0.0)
Squamous cell carcinoma of the tongue	1 (0.5)	0 (0.0)
Acute myocardial infarction	1 (0.5)	0 (0.0)
Cardiac failure	1 (0.5)	0 (0.0)
Cholecystitis acute	0 (0.0)	1 (0.5)

Source: Reviewer's analysis