

Demographic Characteristics	TP1		TP2	
	EU-Stelara (n=253)	CT-P43 (n=256)	EU-Stelara (n=125)	CT-P43 (n=377)
Mean	42.18	42.55	42.99	42.13
Standard Deviation	19	19	20	19
Sex				
F	80 (31.6%)	95 (37.1%)	39 (31.2%)	133 (35.3%)
M	173 (68.4%)	161 (62.9%)	86 (68.8%)	244 (64.7%)
Baseline Body Weight (kg)				
Mean	86	85	86	85
Standard Deviation	1.2	1.2	1.7	1.0
Race				
Asian	23 (9.1%)	25 (9.8%)	10 (8.0%)	34 (9.0%)
White	230 (90.9%)	231 (90.2%)	115 (92.0%)	343 (91.0%)
Ethnicity				
Hispanic or Latino	1 (0.4%)	1 (0.4%)	1 (0.8%)	1 (0.3%)
Not Hispanic or Latino	252 (99.6%)	255 (99.6%)	124 (99.2%)	376 (99.7%)
Country				
Poland	188 (74.3%)	189 (73.8%)	91 (72.8%)	283 (75.1%)
Ukraine	36 (14.2%)	37 (14.5%)	21 (16.8%)	52 (13.8%)
Korea, South	23 (9.1%)	25 (9.8%)	10 (8.0%)	34 (9.0%)
Estonia	6 (2.4%)	5 (2.0%)	3 (2.4%)	8 (2.1%)

Source: Clinical Reviewer

Clinical Reviewer's Comment: The demographic and disease characteristics, in terms of age, sex, race, ethnicity, and mean baseline body weight, were similar across the two treatment arms, EU-Stelara and CT-P43, for Treatment Period I (TP1) and for Treatment Period II (TP2).

Overall AE profile

The phase 3 clinical study, Study CT-P43 3.1, provides the main comparative data for the safety review. During TP1, the overall rate of TEAEs was higher for the CT-P43 treatment group (37.1% - 155 events in 95 subjects), compared to the EU-Stelara treatment group (29.6% - 108 events in 75 subjects) (Table 20). The overall rates of SAEs were similar 1.6% (4 subjects) for the CT-P43 treatment and 1.6% (4 subjects) for the EU-Stelara treatment groups. No subjects discontinued treatment due to an AE during TP1.

During TP2, TEAEs were reported in 34.0% (144 events in 86 subjects) for the CT-P43 maintenance, in 40.8% (78 events in 51 subjects) for the EU-Stelara maintenance and in 41.9% (95 events in 52 subjects) for the Switched to CT-P43 treatment groups. The overall rates of SAEs were similar in the CT-P43 maintenance, EU-Stelara maintenance (2.4%), and Switched to CT-P43 (1.6%) groups. However, slightly more subjects discontinued due to an AE in the CT-P43 maintenance group (2.0%) compared to the EU-Stelara maintenance (0.8%) and Switched to CT-P43 (TP2) (0.8%) groups (Table 36). The details for the discontinuations are discussed further in the section **Dropouts and Discontinuations** below.